

Synthesis and crystal structures of *cis*-palladium(II) and *cis*-platinum(II) complexes containing dipyridyl ligands

Noemí Andrade-López^{*}, José G. Alvarado-Rodríguez, Simplicio González-Montiel, María Elena Páez-Hernández, Carlos Andrés Galán-Vidal, and Araceli Jiménez-Pérez

Centro de Investigaciones Químicas. Universidad Autónoma del Estado de Hidalgo, Ciudad Universitaria, Carretera Pachuca-Tulancingo Km. 4.5. Colonia Carboneras, Mineral de la Reforma. C. P. 42184, México

E-mail: nandrade@uaeh.edu.mx

Dedicated to Professor Rosalinda Contreras Theurel, an exceptional researcher

Abstract

Coordinative behavior of the di-(2-pyridyl)hydroxymethane (**1**) and di-(2-pyridyl)-2-pyridylsulfanylmethane (**2**) ligands towards Pd(II) and Pt(II) is described. The compounds **3-5** were characterized by IR, mass spectrometry and NMR spectroscopy. Crystal structures of [Pt{(C₅H₄N)₂CH(OH)}Cl₂] **3** and [Pd{(C₅H₄N)₂CH(2-C₅H₄NS)}Cl₂] **4** as DMSO solvates displayed the formation of neutral mononuclear complexes. In the complexes **3** and **4** is observed the formation of six-membered chelate rings. The local geometry of the metallic atoms can be described as square planar in the title complexes.

Keywords: Pd(II) and Pt(II) complexes, di-(2-pyridyl)-2-pyridylsulfanylmethane, di-(2-pyridyl)hydroxymethane, NMR, X ray diffraction

Introduction

Several nitrogenated heterocyclic compounds as dipyridines have been widely used in coordination chemistry.¹⁻⁶ These compounds act as versatile ligands towards transition metals which yield five- and six-membered chelate rings in a bidentate coordination mode⁷⁻¹⁶ Synthesis of compounds containing five-membered rings can be envisaged by the use of bipyridine (unbridged) derivatives (Ia, Figure 1).^{7, 8} When the dipyridinic ligands have flexible bridging groups as methylene, hydroxymethyl, amine, hydrazone and amide groups, they mainly form six-membered rings (Ib).⁹⁻¹⁶ In other dipyridines with less flexible bridging groups, the formation of five-membered rings (Ic) has been reported.^{17, 18} Recently, we described the synthesis of complexes of Li(I)¹⁹ and Pd(II)²⁰ and the formation of five-membered rings by the use of the di-(2-pyridyl)-*N*-methylimine ligand; the formation of these rings has been attributed to the rigidity

of the C=N iminic bond. Continuing with our research, here we report the coordinative behavior of the flexible dipyridines **1** and **2** towards heavier tenth group Pd(II) and Pt(II) metals. Synthesis and characterization by NMR spectroscopy, mass spectrometry and IR of *cis*-dichloro[di-(2-pyridyl)hydroxymethane]platinum(II) **3** and *cis*-dichloro-[di-(2-pyridyl)-2-pyridylsulfanylmethane]palladium(II) **4** and *cis*-dichloro-[di-(2-pyridyl)-2-pyridylsulfanylmethane]platinum(II) **5** complexes are described. Crystal structures of the DMSO solvates **3** and **4** are reported.





Figure 1. Complexes containing five- and six-membered chelate rings derived from dipyridines.

Results and Discussion

Synthesis of ligands **1** and **2**


Di-(2-pyridyl)hydroxymethane **1** was synthesized from the reduction of the di-(2-pyridyl)ketone.²¹ Di-(2-pyridyl)-2-pyridylsulfanylmethane **2** was prepared from the reaction of di-(pyridyl)chloromethane²¹ with 2-mercaptopyridine in equimolar ratio, see Scheme 1.



Scheme 1. Synthesis of ligands **1** and **2**.

Synthesis of palladium(II) and platinum(II) complexes **3-5**

Complexes **3-5** were synthesized from the reaction of **1** and **2** with PdCl_2 or PtCl_2 in equimolar ratio in hot acetonitrile, see Scheme 2.



Scheme 2. Synthesis of the *cis*-dichloro complexes **3-5**.

NMR Spectra

^1H and ^{13}C NMR data and the atom numbering for compounds **2**, **3**, **4** and **5** are showed in Table 1.

3 [Pt{(C ₅ H ₄ N) ₂ CH(OH)}Cl ₂]	DMSO- <i>d</i> ₆	8.92 (d, 2H, <i>J</i> = 5.12 Hz; H6)	157.7 (C2)
		8.15 (dd, 2H, <i>J</i> = 7.72, 7.52 Hz; H4)	153.2 (C6)
		7.95 (d, 1H, <i>J</i> = 5.12 Hz; OH)	141.0 (C4)
		7.85 (d, 2H, <i>J</i> = 7.72 Hz; H3)	125.8 (C5)
		7.50 (dd, 2H, <i>J</i> = 7.52, 5.12 Hz; H5)	122.2 (C3)
		6.37 (d, 1H, <i>J</i> = 5.12 Hz; H7)	74.4 (C7)
4 [Pd{(C ₅ H ₄ N) ₂ CH(2-SC ₅ H ₄ N)}Cl ₂]	DMSO- <i>d</i> ₆	8.92(d, 2H, <i>J</i> = 5.48 Hz; H6)	155.6 (C9)
		8.61 (d, 1H, <i>J</i> = 4.76 Hz; H11)	154.3 (C2)
		8.11(dd, 2H, <i>J</i> = 7.72, 6.60 Hz; H4)	154.2 (C6)
		8.05 (d, 2H, <i>J</i> = 7.72 Hz; H3)	150.2 (C11)
		7.77 (dd, 1H, <i>J</i> = 7.72, 7.68 Hz; H13)	141.3 (C4)
		7.55 (m, 3H, H5, H14)	138.2 (C13)
		7.26 (dd, 1H, <i>J</i> = 4.76, 7.68 Hz; H12)	126.7 (C3)
		6.95 (s, 1H, H7)	125.6 (C5)
			122.9 (C14)
	121.8 (C12)		
	54.4 (C7)		
5 [Pt{(C ₅ H ₄ N) ₂ CH(2-SC ₅ H ₄ N)}Cl ₂]	DMSO- <i>d</i> ₆	9.03 (d, 2H, <i>J</i> = 5.84 Hz; H6)	155.2 (C9)
		8.62 (d, 1H, <i>J</i> = 4.40 Hz; H11)	154.8 (C6)
		8.13 (dd, 2H, <i>J</i> = 7.72, 7.68 Hz; H4)	154.0 (C2)
		8.02 (d, 2H, <i>J</i> = 7.72 Hz; H3)	150.1 (C11)
		7.71 (dd, 1H, <i>J</i> = 7.90, 7.68 Hz; H13)	141.1 (C4)
		7.54(dd, 2H, <i>J</i> = 7.68 Hz, 5.88 Hz; H5)	138.4 (C13)
		7.47 (d, 1H, <i>J</i> = 7.90, Hz; H14)	126.4 (C3)
		7.24 (dd, 1H, <i>J</i> = 7.68 Hz, 4.40 Hz; H12)	126.1 (C5)
		6.93 (s, 1H, H7)	122.3 (C14)
	121.7 (C12)		
	54.1 (C7)		

¹H NMR spectrum in DMSO-*d*₆ solution of **2** showed seven signals in the 8.47-7.07 ppm region and a singlet at lower frequency (6.47 ppm) for the proton of the CH group, see Table 1. In this ligand, the H3-H6 and H11-H14 protons were observed in a 2:1 ratio. Its ¹³C{¹H} spectrum showed ten signals for the pyridine rings; the bridging C7 atom was observed at 54.2 ppm. Complexes **3**, **4** and **5** are slightly soluble in acetonitrile and dimethylsulfoxide and insoluble in chloroform and benzene. ¹H spectra in DMSO-*d*₆ solutions for **3**, **4** and **5** showed four, seven and eight signals corresponding to the pyridine ring protons, respectively. In **4** and **5**, the signal intensities for the di-2-pyridylmethyl and the 2-pyridylsulfanyl fragment protons were

observed in a 2:1 ratio, as in the compound **2**. In **3**, the OH and H7 protons were observed as doublets at 7.95 and 6.37 ppm, respectively, whereas the H7 proton of the CH bridging group in **4** and **5** was observed as a singlet at 6.95 and 6.93 ppm, respectively, Table 1. ^1H NMR spectra of complexes **3-5** did not show significant variations in the chemical shift with respect to the free ligands **1** and **2**.

On the other hand, $^{13}\text{C}\{^1\text{H}\}$ spectrum of complex **3** showed five signals for the pyridine rings and the spectra of **4** and **5** displayed ten signals. The bridging C7 carbon in the complexes **3**, **4** and **5** was observed at 74.4, 54.4 and 54.1 ppm, respectively. In these complexes, the metallic coordination affects mainly the chemical shifts of the neighboring carbon atoms to the coordinate nitrogen atoms (C2 is shielded and C6 is deshielded).

The comparison of ^1H and ^{13}C NMR data indicates that the complexes **4** and **5** have analogous structures, where the palladium(II) and platinum(II) cations cause similar chemical shifts. The same situation occurs when the NMR spectra of complex **3** are compared with those of the palladium(II) analogue.²⁰ These data and the magnetic equivalence of the pyridine rings of the dipyridine fragment suggest the formation of symmetrical palladium and platinum complexes with six-membered chelate rings.

X-ray crystallography

The structures in crystalline solid state of compounds **3** and **4** were determined by single-crystal X-ray diffraction analyses; they crystallized as DMSO solvates. Crystallographic data are given in Table 2 and selected bond lengths and angles are given in Table 3.

Molecular structures of complexes **3**·DMSO and **4**·DMSO are showed in Figures 2 and 3.

The X-ray structures of **3**·DMSO and **4**·DMSO showed that the pseudo-anthracenic tricyclic systems are not planar; the angles between the pyridine rings are 110.9° and 123.5° for **3** and **4**, respectively. When the ligands **1** and **2** coordinate to the M(II) central atom, they behave as bidentate ligands, and leave the hydroxyl and 2-pyridylsulfanyl groups uncoordinated. In both complexes, the chelate ring is six membered, as proposed by the NMR analyses, and the local geometry of the M(II) central atom is square planar.

Table 2. Selected crystallographic data for complexes **3**·DMSO and **4**·DMSO

Compound	3 ·DMSO	4 ·DMSO
Molecular Formula	C ₁₃ H ₁₆ Cl ₂ N ₂ O ₂ PtS	C ₁₈ H ₁₉ Cl ₂ N ₃ OPdS ₂
<i>M_r</i> [g/mol]	530.33	534.78
Crystal size [mm]	0.58 x 0.35 x 0.23	0.33 x 0.30 x 0.12
F(000)	504	536
Crystal system	Triclinic	Triclinic
Space group	<i>P1bar</i>	<i>P1bar</i>
<i>a</i> [Å]	9.7309(9)	8.2035(8)
<i>b</i> [Å]	10.0992(9)	8.5934(8)
<i>c</i> [Å]	10.9146(10)	15.5817(15)
α [°]	65.232(2)	76.956(2)
β [°]	89.001(2)	81.603(2)
γ [°]	61.555(2)	85.196(2)
<i>V</i> [Å ³]	833.80(13)	1057.16(18)
<i>Z</i>	2	2
$\rho_{\text{calcd.}}$ [Mg/m ³]	2.112	1.680
μ [mm ⁻¹]	8.863	1.341
<i>GoF</i>	1.094	1.090
Temperature [K]	298(2)	298(2)
θ range [°]	2.11 to 26.01	1.35 to 26.04
Reflections collected	10132	13067
Unique reflections	3278	4164
Absorption correction	SADABS	SADABS
Solution method	Patterson	Patterson
<i>R</i> _{int}	0.0286	0.0392
<i>R</i> ₁ ; <i>wR</i> ₂ [<i>I</i> >2σ(<i>I</i>)]	0.0216, 0.0556	0.0353, 0.0818
<i>R</i> ₁ ; <i>wR</i> ₂ [all data]	0.0242, 0.0565	0.0514, 0.0976
Largest residuals [e·Å ⁻³]	0.543/-0.765	0.553/ -0.348

As can be observed from the listed data in Table 3, the N-M(II) distances in these compounds are essentially the same and similar to other previously reported for dipyridine and pyridine complexes containing N-M bonds (M= Pd, Pt).^{2, 4, 11, 14, 22, 23} The close similitude of the Pt-Cl and Pd-Cl bond distances in complexes **3** and **4** can be attributed to the lanthanide contraction effect present in the 5d platinum(II) cation.

With regard to the conformational features displayed by the complexes **3** and **4** where six-membered chelate rings were formed, the central ring in both complexes is in a boat conformation. The position of the exocyclic ligands attached to the bridging carbon that joins the two pyridine rings deserves a more detailed discussion. The predominant conformers in solid state display the -OH group in **3** in a bowsprit position, as in the palladium analogue,²⁰ and the -

SC₅H₄N group in **4** in a flagpole position. This difference can be attributable to an intramolecular contact between the soft acid palladium and the soft base sulfur (Pd---S = 3.054 Å) that is shorter than the van der Waals radii sum ($\Sigma r_{\text{vdw}}(\text{Pd},\text{S}) = 3.43 \text{ \AA}$) but longer than the covalent radii sum ($\Sigma r_{\text{cov}}(\text{Pd},\text{S}) = 2.33 \text{ \AA}$).²⁴ This interaction can be also responsible of the large folding in the tricyclic system observed in the complex **4** (interplanar angle 123.5° in **4** respect to 110.9° in **3**).

Table 3. Selected bond lengths [Å] and bond angles [°] of complexes

[M{(C ₅ H ₄ N) ₂ G}Cl ₂].DMSO		
Compound	3 ·DMSO	4 ·DMSO
	M=Pt, G =CH(OH)	M=Pd, G =CH(SC ₅ H ₄ N)
M-Cl(1)	2.3020(12)	2.3003(11)
M-Cl(2)	2.2962(12)	2.3037(11)
M-N(1)	2.026(3)	2.036(3)
M-N(2)	2.015(3)	2.034(3)
C(11)-Y(1) ^a	1.401(5)	1.838(4)
Cl(1)-M-Cl(2)	91.03(5)	92.25(4)
N(1)-M-N(2)	86.76(13)	87.76(12)
Cl(1)-M-N(1)	90.94(10)	90.05(9)
Cl(1)-M-N(2)	177.68(9)	177.81(9)
Cl(2)-M-N(1)	177.37(9)	177.34(9)
Cl(2)-M-N(2)	91.27(10)	89.93(9)
C(5)-C(11)-C(6)	108.5(3)	112.9(3)
N(1)-C(5)-C(11)-Y(1) ^a	-176.7(3)	-70.3(4)
N(2)-C(6)-C(11)-Y(1) ^a	176.1(3)	68.6(4)

a) For M=Pt, Y1 = O1; for M= Pd, Y1 = S1.




Figure 2. ORTEP diagram of 3·DMSO with labeled atoms and 40% probability displacement ellipsoids (DMSO molecule is omitted for clarity).




Figure 3. ORTEP diagram of 4·DMSO with labeled atoms and 40% probability displacement ellipsoids (DMSO molecule is omitted for clarity).

Experimental Section

General Procedures. Unless otherwise noted, all reagents used in this work are commercially available (Aldrich) and were used without further purifications. Compounds di-(2-pyridyl)chloromethane and di-(2-pyridyl)hydroxymethane **1** were prepared according to the literature procedure.²¹ Solvents were dried and distilled by standard methods before use. Melting points were determined with a Mel-Temp II instrument and are uncorrected. Spectra were recorded with the following instruments. Elemental analyses: Perkin–Elmer Series II CHNS/O Analyzer. IR spectra: 4000–400 cm⁻¹ range, Perkin–Elmer System 2000 FTIR spectrometer, as KBr pellets. Mass spectra: EI- and FAB-mass spectra: Jeol JMS-AX505HA with a Xe beam at 6 keV, and nitrobenzyl alcohol as matrix (FAB). NMR spectra: Jeol Eclipse 400 spectrometer, residual solvent signal used as reference for ¹H NMR spectra; ¹³C{¹H} NMR spectra referenced through the solvent peaks; chemical shifts quoted on the δ scale (downfield shifts positive) relative to tetramethylsilane (¹H, ¹³C{¹H}); spectra recorded at about 20°C; ¹H, 399.78 MHz; ¹³C{¹H}, 100.53 MHz. Assignments of all compounds were performed by means of homonuclear correlation (cosy) and heteronuclear correlation (hetcor) experiments.

Suitable single crystals of **3** and **4** were obtained by slow evaporation of a DMSO solution in a NMR tube. X-ray diffraction data on **3** and **4** were collected at about 25°C with a CCD Smart 6000 diffractometer through the use of Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$, graphite monochromator). Data were integrated, scaled, sorted and averaged using the SMART software package. Structures were solved by direct methods, using SHELXTL NT Version 5.10²⁵ and refined by full-matrix least squares against F^2 . An empirical absorption correction based on the multiple measurement of equivalent reflections was applied by using the program SADABS. The displacement parameters of non-hydrogen atoms were refined anisotropically. The positions of the hydrogen atoms were kept fixed with a common isotropic displacement parameter. Crystallographic data are given in Table 2.

Di-(2-pyridyl)-2-pyridylsulfanylmethane (**2**)

A mixture of di-(2-pyridyl)chloromethane (200 mg, 0.98 mmol) and 2-mercaptopyridine (110 mg, 0.98 mmol) in 20 mL of dry CHCl₃ was refluxed for 16 h. Removal of the solvent under vacuum was followed by addition of 20 mL of water and acidified with HCl 2 M and stirred for 10 min. The aqueous layer was neutralized with K₂CO₃ and extracted with 25 mL of chloroform; this solution was dried by means of a column of Celite and Na₂SO₄. The slow evaporation of the solvent yielded a brown solid. Yield: 92 %, (245 mg). m. p.: 77-79°C. - IR: ν (cm⁻¹) = 1618 (C=N), 1579, 1555 (C=C). - Anal. Calcd for C₁₆H₁₃N₃S; C, 68.79; H, 4.69. Found: C, 68.07, H, 4.74. EI: m/z 279 [M⁺].

Cis-dichloro[di-(2-pyridyl)hydroxymethane]platinum (II) (**3**)

To a solution of PtCl₂ (250 mg, 0.94 mol) in 40 mL of hot acetonitrile were added 175 mg (0.94 mmol) of **1**. The resulting brown solution was refluxed by 24 h and cooled to room temperature to give a brown suspension. The cold suspension was filtered and the yellow solid obtained was

dried in vacuum. Yield: 86% (366 mg). Dec.: 275°C. - IR: ν (cm⁻¹) = 1606 (C=N), 1565 (C=C). - Anal. Calcd for C₁₁H₁₀Cl₂N₂OPt•(CH₃)₂SO; C, 29.44; H, 3.04. Found: C, 29.32, H, 2.92. FAB+: m/z 452 [M⁺], 417 [M⁺-Cl].

Cis-dichloro-[di-(2-pyridyl)-2-pyridylsulfanylmethane]palladium(II) (4)

To a solution of PdCl₂ (64 mg, 0.35 mmol) in 25 mL of hot acetonitrile were added 100 mg (0.35 mmol) of **2**. The brown solution obtained was refluxed by 24 h and cooled to room temperature to give a brown suspension. The cold suspension was filtered and the brown solid obtained was dried in vacuum. Yield: 90% (147 mg). m. p.: 243°C. - IR: ν (cm⁻¹) = 1602, 1576, 1565 (C=C). - Anal. Calcd for C₁₆H₁₃Cl₂N₃SPd•(CH₃)₂SO; C, 40.42; H, 3.58. Found: C, 41.00, H, 3.23. FAB+: m/z 522 [M⁺-Cl].

Cis-dichloro-[di-(2-pyridyl)-2-pyridylsulfanylmethane]platinum(II) (5)

Complex **5** was prepared using the method described for the formation of the complexes **3** and **4**, with PtCl₂ (95 mg, 0.35 mol) and 100 mg (0.35 mmol) of **2** in 25 mL of acetonitrile. **5** Was obtained as a yellow solid. Yield: 85% (166 mg). Dec.: 253°C. - IR: ν (cm⁻¹) = 1590, 1571 (C=C). FAB+: m/z 545 [M⁺], 510 [M⁺-Cl].

Supplementary Information Available

Supplementary data are available from the Cambridge Crystallographic Data Centre, with the deposition numbers CCDC649169 (**3**) and CCDC649168 (**4**). Copies of this information are available free of charge from the Director, CCDC, 12 Union Road, Cambridge, CB2 1EZ, UK (Fax: +44 1223 336 033; e-mail: deposit@ccdc.cam.ac.uk or <http://www.ccdc.cam.ac.uk>).

Acknowledgements

A.J.P. fully acknowledges to CONACyT (44009) the scholarship for supporting their studies. We wish to thank Dra. Verónica García Montalvo (UNAM) for the mass spectra and Mrs. Yolanda Marmolejo (UAEH) for the vibrational spectra and elemental analyses.

References

1. Vedernikov, A. N.; Fettingner, J. C.; Mohr, F. *J. Am. Chem. Soc.* **2004**, *126*, 11160.
2. Steel, J.; Sumby, C. J. *Dalton Trans.* **2003**, 4505.
3. Katsoulakou, E.; Lalioti, N.; Raptopoulou, C. P.; Terzis, A.; Manessi-Zoupa, E.; Perlepes, S. P. *Inorg. Chem. Commun.* **2002**, 719.
4. Schareina, T.; Hillebrand, G.; Fuhrmann, H.; Kempe, R. *Eur. J. Inorg. Chem.* **2001**, 2421.

5. Carr, J. D.; Coles, S. J.; Hursthouse, M. B.; Light, M. E.; Munro, E. L.; Tucker, J. H. R.; Westwood, J. *Organometallics* **2000**, *19*, 3312.
6. Bernhard, P. V.; Comba, P.; Mahu-Rickenbach, A.; Stebler, S.; Steiner, S.; Várnagy, K.; Zehnder, M. *Inorg. Chem.* **1992**, *31*, 4194.
7. Godard, C.; Duckett, S. B.; Parsons, S.; Perutz, R. N. *Chem. Comm.* **2003**, *18*, 2332.
8. Milani, B.; Mestroni, G.; Zangrando, E. *Croat. Chem. Acta* **2001**, *4*, 851.
9. Garcia, A. M.; Manssur, J. *Acta Crystallogr., Sect. C.* **1994**, 1882.
10. Gornitzka, H.; Stalke, D. *Organometallics* **1994**, *13*, 4398.
11. Crowder, K. N.; Garcia, S. J.; Burr, R. L.; North, J. M.; Wilson, M. H.; Conley, B. L.; Fanwick, P. E.; White, P. S.; Sienerth, K. D.; Granger, R. M. *Inorg. Chem.* **2004**, *43*, 72.
12. Wen-Rui, Y.; Ze-Hua, L.; Qian-Feng Z. *Acta Cryst.* **2003**, *C59*, m139.
13. Kin-Ying, H.; Wing-Yiu, Y.; Kung-Kai, C.; Chi-Ming, C. *Chem. Commun.* **1998**, 2101.
14. Mock, C.; Puscasu, I.; Rauterkus, M. J.; Tallen, G.; Wolf, J. E. A.; Krebs, B. *Inorg. Chim. Acta* **2001**, *319*, 109.
15. Bakir, M.; Brown, O. *J. Mol. Struct.* **2002**, *641*, 183.
16. Silberg, J.; Schareina, T.; Kempe, R.; Wurst K.; Buchmeiser, M. R. *J. Organomet. Chem.* **2001**, *622*, 6.
17. Grewe, J.; Hagenbach, A.; Stromburg, B.; Alberto, R.; Vazquez-Lopez, E.; Abram, U. Z. *Anorg. Allg. Chem.* **2003**, *629*, 303.
18. Canty, A. J.; Chaichit, N.; Gatehouse, B. M.; George, E. E.; Hayhurst, G. *Inorg. Chem.* **1981**, *20*, 2414.
19. Flores-Chávez, B.; Martínez-Ortega, B. A.; Alvarado-Rodríguez, J.G.; Andrade-López, N. *J. Chem. Crystallogr.* **2005**, *35*, 435.
20. Andrade-López, N.; Alvarado-Rodríguez, J. G.; González-Montiel, S.; Rodríguez-Méndez, M. G.; Hernández-Páez, M. E.; Galán-Vidal, C. A. *doi:10.1016/j.poly.2007.06.006*.
21. La Crois, R. Ph. D. thesis, University Library of Groningen, Netherlands, **2002**.
22. Annibale, G.; Canovese, L.; Cattalini, L.; Natile, G.; Biagini-Cingi, M.; Manotti-Lanfredi, A.-M.; Tiripicchio, A. *J. Chem. Soc., Dalton Trans.* **1981**, 2280.
23. Canty, A. J.; Minchin, N. J.; Engelhardt, L. M.; Skelton, B. W.; White, A. H. *J. Chem. Soc. Dalton Trans.* **1986**, 645.
24. Porterfield, W. W., *Inorganic Chemistry: A Unified Approach*, 2nd ed., Academic Press, Inc. USA, **1993**, 214.
25. SHELXTL 5.10 Bruker AXS, Inc., Madison, WI, USA, **1998**.