

Reagents for new heteroannellation reactions part V: 2-(methylthio)-2-imidazoline

Johannes Frohlich*, Fritz Sauter, A. Z. M. Shaifullah Chowdhury[#], and Christian Hametner*

*Institute of Organic Chemistry, Vienna University of Technology
Getreidemarkt 9, A-1060 Vienna, Austria.*

E-mail: jfroehli@pop.tuwien.ac.at

(received 07 Mar 00; accepted 20 Aug 00; published on the web 28 Aug 00)


Abstract

Double annellation of an imidazo[1,2-a]pyrimidine moiety was achieved in a one-pot process by reacting heteroaromatic 2-aminoesters and 2-aminonitriles with 2-(methylthio)-2-imidazoline, obtaining a number of mostly novel tetracyclic hetero-systems.


Keywords: Methylthioimidazoline, aminoesters, aminonitriles

Introduction


Within a long-term project dealing with the synthesis of novel fused heterocyclic systems we have shown that compounds of a N-[bis(methylthio)methylene]-amino (BMMA) type are versatile reagents for a one-pot annellation of a pyrimidine ring to 2-aminoesters:



Furthermore we have extended this methodology towards cyclic analogs of the BMMA reagents, using methylthio-substituted thiazole and thiazine derivatives for double-annellation reactions¹:




In the present paper we report on the utilization of 2-(methylthio)-2-imidazoline (1) for double annelation reactions, expanding the BMMA strategy towards the construction of N,N-heterocycles.




Results and Discussion

In contrast to the cyclizations of the thiazole and thiazine reagents, heating in dry acetic acid turned out to be unsuccessful in case of 1. Thus in a large number of experiments optimal conditions concerning solvent and temperature had to be revealed. Finally, heating the starting materials in HMPA at 160 °C for several hours proved to yield best results, and a variety of heteroaromatic substrates was reacted with 2-(methylthio)-2-imidazoline to obtain the desired tetracyclic fusion products:




Annulation of 1 with 2-aminoesters produced oxo compounds 2,



Compd	X
2a	CH-CH ₃
2b	N-CH ₂ -Ph
2c	S

whereas 2-aminonitriles gave the analogous imino derivatives 3 and 4.



Compd	X	Y	n
3a	CH ₂	S	1
3b	CH ₂	S	2
3c	CH-CH ₃	S	2
3d	CH ₂	S	3
3e	N-CH ₃	S	2
3f	N-CH ₂ -Ph	S	2
3g	S	S	2
4	CH ₂	O	2

Some compounds derived from parent systems described here have already been synthesized by alternative, multi-step pathways. However, the method reported here allows smooth access to complex fused products starting from easily obtainable substrates (e.g. by *Gewald* reaction) in one step.

Experimental Section

General Procedures. Melting points were determined on a Kofler hot stage apparatus and are uncorrected. ^1H and ^{13}C NMR spectra were recorded on a Bruker AC 200 spectrometer (TMS as internal standard, DMSO- d_6 as solvent, δ -values in ppm). Elementary analyses were performed at the Microanalytical Laboratory, Institute of Physical Chemistry, University of Vienna (Mag. J. Theiner).

2-(Methylthio)-2-imidazoline (1) was prepared *via* a two step procedure from ethylene diamine, CS_2 , and methyl iodide. The starting aminoesters and aminonitriles were prepared according to literature procedures.

General procedure for the reaction of *I* with 2-aminoesters. Synthesis of compounds *2a-c*

2-(Methylthio)-2-imidazoline (3 mmol) and the aminoester (2 mmol) were dissolved in hexamethylphosphoric acid triamide (3 mL) and heated to 160 °C for 3 h. After cooling to room temperature, crushed ice was added and the mixture stirred for 1 h. The separated product was collected by filtration and crystallized from an appropriate solvent.

2,3,6,7,8,9-Hexahydro-8-methyl-[1]benzothieno[2,3-*d*]imidazo[1,2-*a*]pyrimidin-5(1H)-one (*2a*) From ethyl 2-amino-4,5,6,7-tetrahydro-6-methylbenzo[*b*]thiophene-3-carboxylate; yield: 76%; m.p.: 290 °C (methanol); $\text{C}_{13}\text{H}_{15}\text{N}_3\text{OS}$ (261.34); $^1\text{H-NMR}$ δ 7.70 (s, 1H), 4.00 (t, 2H), 3.60 (t, 2H), 2.90-1.20 (m, 7H), 1.00 (d, 3H); $^{13}\text{C-NMR}$ δ 166.00 (s), 156.95 (s), 155.43 (s), 129.50 (s), 124.81 (s), 113.82 (s), 41.81 (2t), 32.37 (t), 30.16 (t), 28.90 (d), 24.82 (t), 21.18 (q).

2,3,6,7,8,9-Hexahydro-8-(phenylmethyl)-imidazo[1,2-*a*]pyrido[4',3':4,5]thieno-[2,3-*d*]pyrimidin-5(1H)-one (*2b*) From ethyl 2-amino-4,5,6,7-tetrahydro-6-(phenylmethyl)-thieno[2,3-*c*]pyridine-3-carboxylate; yield: 73%; m.p.: 235 °C (acetone); $\text{C}_{18}\text{H}_{18}\text{N}_4\text{OS}$ (338.43); calc.: C 63.88%, H 5.36%, N 16.55%; found: C 63.67%, H 5.23%, N 16.56%; $^1\text{H-NMR}$: δ 7.70 (s, 1H), 7.40-7.20 (m, 5H), 4.00 (s, 2H), 4.00 (s, 2H), 3.60 (s, 2H), 3.50 (s, 2H), 2.80-2.60 (m, 4H).

1,2,3,6,7,9-Hexahydro-5H-imidazo[1,2-*a*]thiopyrano[4',3':4,5]thieno[2,3-*d*]pyrimidin-5-one (2c). From ethyl 2-amino-4,7-dihydro-5H-thieno[2,3-*c*]thiopyran-3-carboxylate; yield: 73%; m.p.: 245 °C (methanol); $\text{C}_{11}\text{H}_{11}\text{N}_3\text{OS}_2$ (265.36); calc.: C 49.78%, H 4.18%, N 15.84%; found: C 49.38%, H 4.04%, N 16.01%; $^1\text{H-NMR}$: δ 7.90 (s, 1H), 4.00 (t, 2H), 3.80 (s, 2H), 3.60 (t, 2H), 3.00 (s, 2H), 2.80 (s, 2H); $^{13}\text{C-NMR}$: δ 165.25 (s), 156.78 (s), 155.87 (s), 129.69 (s), 121.20 (s), 113.90 (s), 41.98 (t), 41.98 (t), 27.17 (t), 24.68 (t), 24.24 (t).

General procedure for the reaction of *I* with 2-aminonitriles

Synthesis of compounds *3a-g* and *4* 2-Methylthio-2-imidazoline (5 mmol) and the aminonitrile (3 mmol) were dissolved in hexamethylphosphoric acid triamide (3 mL) and heated to 160 °C for

a given period of time. After cooling to room temperature, crushed ice was added and the mixture stirred for 1 h. The separated product was collected by filtration and crystallized from an appropriate solvent.

1,2,3,6,7,8-Hexahydro-5*H*-cyclopenta[4,5]thieno[2,3-*d*]imidazo[1,2-*a*]pyrimidin-5-imine

(3a). From 2-amino-5,6-dihydro-4*H*-cyclopenta[*b*]thiophene-3-carbonitrile; reaction time: 3 h; yield: 56%; m.p.: >320 °C (methanol); C₁₁H₁₂N₄S (232.30); ¹H-NMR: δ 7.50 (s, 1H), 6.50 (s, 1H), 3.90 (t, 2H), 3.60 (t, 2H), 2.90 (t, 2H), 2.80 (t, 2H), 2.30 (m, 2H); ¹³C-NMR: δ 166.45 (s), 155.79 (s), 154.97 (s), 137.90 (s), 129.82 (s), 109.40 (s), 42.84 (t), 42.84 (t), 28.68 (t), 28.58 (t), 27.32 (t).

2,3,6,7,8,9-Hexahydro-[1]benzothieno[2,3-*d*]imidazo[1,2-*a*]pyrimidin-5(1*H*)-imine (3b).

From 2-amino-4,5,6,7-tetrahydrobenzo[*b*]thiophene-3-carbonitrile; reaction time: 10 h; yield: 61%; m.p.: 265 °C (methanol); C₁₂H₁₄N₄S (246.33); ¹H-NMR: δ 7.40 (s, 1H), 6.60 (s, 1H), 3.90 (t, 2H), 3.80 (t, 2H), 3.60 (t, 2H), 2.50 (t, 2H), 1.80 (m, 4H); ¹³C-NMR: δ 160.44 (s), 155.02 (s), 152.43 (s), 129.37 (s), 124.60 (s), 111.96 (s), 42.54 (t), 39.65 (t), 25.61 (t), 24.34 (t), 22.39 (t), 22.08 (t).

2,3,6,7,8,9-Hexahydro-8-methyl-[1]benzothieno[2,3-*d*]imidazo[1,2-*a*]pyrimidin-5(1*H*)-imine

(3c). From 2-amino-4,5,6,7-tetrahydro-6-methylbenzo[*b*]thiophene-3-carbonitrile; reaction time: 4 h; yield: 59%; m.p.: 266 °C (methanol); C₁₃H₁₆N₄S (260.36); ¹H-NMR: δ 7.40 (s, 1H), 6.50 (s, 1H), 3.90 (t, 2H), 3.60 (t, 2H), 2.70 (m, 2H), 2.20 (m, 2H), 1.80 (m, 2H), 1.20 (m, 1H), 1.00 (d, 3H).

1,2,3,6,7,8,9,10-Octahydro-5*H*-cyclohepta[4,5]thieno[2,3-*d*]imidazo[1,2-*a*]pyrimidin-5-imine

(3d). From 2-amino-5,6,7,8-tetrahydro-4*H*-cyclohepta[*b*]thiophene-3-carbonitrile¹¹; reaction time: 3 h; yield: 57%; m.p.: 237 °C (methanol); C₁₃H₁₆N₄S (260.36); ¹H-NMR: δ 7.40 (s, 1H), 6.40 (s, 1H), 3.90 (t, 2H), 3.50 (t, 2H), 3.20 (m, 2H), 2.70 (m, 2H), 1.80 (m, 2H), 1.60 (m, 4H); ¹³C-NMR: δ 158.71 (s), 154.63 (s), 151.35 (s), 135.82 (s), 128.19 (s), 113.47 (s), 48.55 (t), 42.25 (t), 31.68 (t), 28.58 (t), 27.80 (t), 27.54 (t), 26.74 (t).

2,3,6,7,8,9-Hexahydro-8-methylimidazo[1,2-*a*]pyrido[4',3':4,5]thieno[2,3-*d*]pyrimidin-5(1*H*)-imine (3e).

From 2-amino-4,5,6,7-tetrahydro-6-methylthieno[2,3-*c*]pyridine-3-carbonitrile; reaction time: 3 h; yield: 46%; m.p.: 262 °C (ethyl acetate); C₁₂H₁₅N₅S (261.34); ¹H-NMR: δ 7.50 (s, 1H), 6.40 (s, 1H), 3.90 (t, 2H), 3.60 (t, 2H), 3.40 (s, 2H), 2.90 (t, 2H), 2.60 (t, 2H), 2.30 (s, 3H); ¹³C-NMR: δ 160.90 (s), 155.20 (s), 152.20 (s), 127.67 (s), 122.21 (s), 111.71 (s), 53.09 (t), 51.01 (t), 45.06 (t), 42.51 (t), 39.71 (t), 26.09 (q).

2,3,6,7,8,9-Hexahydro-8-(phenylmethyl)-imidazo[1,2-*a*]pyrido[4',3':4,5]thieno[2,3-*d*]pyrimidin-5(1*H*)-imine (3f). From 2-amino-4,5,6,7-tetrahydro-6-(phenylmethyl)-thieno[2,3-*c*]pyridine-3-carbonitrile; reaction time: 3 h; yield: 53%; m.p.: 205 °C (ethyl acetate); C₁₈H₁₉N₅S (337.44); ¹H-NMR: 7.50 (s, 1H), 7.40-7.20 (m, 5H), 6.40 (s, 1H), 3.90 (t, 2H), 3.70 (s, 2H), 3.60 (t, 2H), 3.40 (s, 2H), 2.80 (t, 2H), 2.70 (t, 2H); ¹³C-NMR: δ 161.05 (s), 155.15 (s), 152.15 (s), 138.22 (s), 128.74 (2d), 128.20 (2d), 127.93 (s), 126.99 (s), 122.22 (s), 111.62 (s), 60.89 (t), 51.10 (t), 49.19 (t), 42.53 (2t), 25.94 (t).

1,2,3,6,7,9-Hexahydro-5*H*-imidazo[1,2-*a*]thiopyrano[4',3':4,5]thieno[2,3-*d*]pyrimidin-5-imine (3g). From 2-amino-4,7-dihydro-5*H*-thieno[2,3-*c*]thiopyran-3-carbonitrile¹³; reaction time: 12 h; yield: 56%; m.p.: >320 °C (methanol); C₁₁H₁₂N₄S₂ (264.36); ¹H-NMR: δ 7.50 (s, 1H), 6.50 (s, 1H), 3.90 (t, 2H), 3.70 (t, 2H), 3.60 (t, 2H), 3.10 (m, 2H), 2.90 (m, 2H); ¹³C-NMR: δ 160.21 (s), 155.10 (s), 151.56 (s), 129.51 (s), 120.79 (s), 112.21 (s), 42.54 (2t), 27.72 (t), 25.02 (t), 24.65 (t).

2,3,6,7,8,9-Hexahydrobenzofuro[2,3-*d*]imidazo[1,2-*a*]pyrimidin-5(1*H*)-imine (4). From 2-amino-4,5,6,7-tetrahydrobenzofuran-3-carbonitrile; reaction time: 3 h; yield: 58%; m.p.: 278 °C (methanol); C₁₂H₁₄N₄O (230.27); ¹H-NMR: δ 7.60 (s, 1H), 6.20 (s, 1H), 3.90 (t, 2H), 3.60 (t, 2H), 2.70-2.50 (m, 4H), 1.80-1.60 (m, 4H); ¹³C-NMR: δ 162.81 (s), 156.71 (s), 153.55 (s), 144.53 (s), 113.78 (s), 95.64 (s), 49.06 (t), 42.98 (t), 22.73 (t), 22.65 (t), 22.56 (t), 21.37 (t).

References

1. Frohlich, J.; Sauter, F.; Shaifullah Chowdhury, A.Z.M.; Hametner, C. *Sci. Pharm.* **1997**, *65*, 83.
2. Sauter, F.; Frohlich, J.; Blasl, K.; Gewalt, K. *Heterocycles* **1995**, *40*, 851.
3. Sauter, F.; Frohlich, J.; Chowdhury, A.Z.M.S.; Hametner, C. *Monatsh. Chem.* **1997**, *128*, 503.
4. Urleb, U.; Stanovnik, B.; Tisler, M. *J. Heterocycl. Chem.* **1990**, *27*, 643.
5. Hofmann, J. *Chem. Ber.* **1872**, *5*, 242.
6. Aspinall, S.R.; Bianco E.J. *J. Am. Chem. Soc.* **1951**, *73*, 602.
7. Perrissin, M.; Luu Doc, C.; Narcisse, G.; Bakri-Logeais, F.; Huguet, F. *Eur. J. Med. Chem. Chim. Ther.* **1980**, *15*, 413.
8. Kato, Y. *Yakugaku Zasshi* **1972**, *93*, 397.
9. Ahmed, E.K. *Monatsh. Chem.* **1995**, *126*, 953.
10. Bogdanowicz-Szwed, K.; Nagraba, K. *Org. Mass. Spectrom.* **1984**, *19*, 528.
11. Gewalt, K.; Schinke, E.; Böttcher, H. *Chem. Ber.* **1966**, *99*, 94.
12. Manhas, M.S.; Rao, V.V.; Amin, S.G. *J. Heterocycl. Chem.* **1976**, *13*, 821.

13. Rosowsky, A.; Chaykovsky, M.; Chen, K.K.N.; Lin, M.; Modest, E.J. *J. Med. Chem.* **1973**, *16*, 185.
14. Elslager, E.F.; Jacob, P.; Werbel, L.M. *J. Heterocycl. Chem.* **1972**, *9*, 775.
15. Ducker, J.W.; Gunter, M.J. *Aust. J. Chem.* **1974**, *27*, 2229.