

Synthesis and structure of some (*E*)-ferrocene-methylenecycloalkanones and their benzylidene analogues¹

Pál Sohár, Antal Csámpai and Pál Perjési*

Research Group for Structural Chemistry and Spectroscopy, Hungarian Academy of Sciences –
Department of General and Inorganic Chemistry, Eötvös Loránd University, P.O. Box 32, H-
1518 Budapest-112, Hungary

*Institute of Pharmaceutical Chemistry, University of Pécs, P.O. Box 99,
H-7602 Pécs, Hungary

E-mail: sohar@para.chem.elte.hu

Dedicated to Professor Gábor Bernáth on his 70th birthday
(received 17 Feb 03; accepted 26 May 03; published on the web 27 May 03)

Abstract

E-2-ferrocenemethylene- and *E,E*-diferrocenemethylenecycloalkanones as well as partial benzylidene analogues of the latter were synthesized by base-catalyzed Claisen-Schmidt condensation of ferrocenecarboxaldehyde with cyclopentanone, cyclohexanone, cycloheptanone and *E*-2-(4'-X-benzylidene)cycloalkanones. The stereostructure (configuration and conformation) and the electronic properties (conjugation of the enone moiety and the aromatic rings) of the compounds were studied by IR, ¹H and ¹³C NMR methods. The spectroscopic properties of the ferrocene derivatives were compared with those of the respective benzylidene analogues.

Keywords: Ferrocene, cyclic enones, synthesis, spectroscopy

Introduction

Arylidencycloalkanones are versatile starting materials for the synthesis of saturated and partially saturated heterocyclic ring systems.¹ Previously, we described the preparation of saturated and partially saturated 1,3-thiazines using *E*-2-arylidencycloalkanones as starting materials.^{2,3} Some diarylidencycloalkanones have been reported to possess cytoprotective^{4,5} as well as HIV-1 integrase⁶ and factor Xa inhibitor activities⁷. These effects were found to depend on the ring size (stereochemistry) and the substituents of the aromatic rings.

¹Study on ferrocenes, Part 11

*Correspondance in relation to syntheses.

Replacement of a benzene ring by a ferrocene moiety influences significantly both the chemical and the biological properties of aromatic compounds.^{8,9} Recently, we have synthesized some *E*-2-ferrocenemethylenebenzocyclokanones and compared their electronic and stereochemical characteristics with the respective benzylidene analogues.^{10,11} As a continuation of this work here we report on the synthesis and the spectroscopic investigation of some *E*-2-ferrocenemethylene- **3** and *E,E*-diferrocenemethylenecycloalkanones **4** as well as their complete **5,6** and partial **7-11** benzylidene analogues (Figure 1).




Figure 1

Experimental Section

General Procedures. Cycloalkanones **1a-c**, ferrocenecarboxaldehyde (**2**) were obtained from Sigma-Aldrich, Hungary. Compounds **5a-c** (and their substituted derivatives) and **6a-c** were prepared according to the methods described in the literature.^{12,13,14} The isolated **4a** and **4b** had the melting points (250 °C (decomp.) and 179-181 °C, respectively) corresponding to the previously reported values.¹⁵ All the compounds were purified by column chromatography over Kieselgel 60 (Merck, 0.063-0.2 mm) using toluene or toluene/methanol (99/1)- as eluents. Melting points were determined on a Boetius apparatus at a heating rate of 4 °C/min and are uncorrected.

The ^1H and ^{13}C NMR spectra were recorded in CDCl_3 solution in 5 mm tubes at RT, on a Bruker DRX-500 spectrometer at 500.13 (^1H) and 125.76 (^{13}C) MHz, with the deuterium signal of the solvent as the lock and TMS as internal standard. The standard Bruker microprogram NOEMULT to generate NOE¹⁶ and to get DIFFNOE spectra^{17a,18} was used with a selective pre-irradiation time. DEPT spectra¹⁹ were run in a standard manner,²⁰ using only a $\Theta=135^\circ$ pulse to separate the CH/CH_3 and CH_2 lines phased 'up' and 'down,' respectively. The 2D-COSY,^{21a,22a} HMQC^{21b,22b} and HMBC^{23,24} spectra were obtained by using the standard Bruker pulse programs COSY-45, INV4GSSW and INV4GSLRNDWS, respectively. IR spectra were run in KBr disks on a Bruker IFS-55 FT-spectrometer controlled by Opus 3.0 software.

Reaction of ferrocenecarboxaldehyde (2) with cycloalkanones (1a-c). 1 mmol ferrocenecarboxaldehyde (2) and 5 mmol cycloalkanone (1a-c) were dissolved in 5-mL of MeOH and the solution was saturated with O_2 -free N_2 -gas for 5 min. In 5 mL MeOH saturated with O_2 -free N_2 , 1 mmol KOH was dissolved and the solution was added to the MeOH solution of 2 and the respective 1a-c. The flask was sealed and kept at RT for 3-5 days. Then the mixture was placed in a refrigerator overnight and the resulting crystals (4a, 4b) were filtered and purified by column chromatography. The filtrate (or the homogeneous reaction mixture) was evaporated at 40°C , the residue was dissolved in 100 mL CHCl_3 , washed with distilled water (to remove all KOH), dried over anhydrous Na_2SO_4 and the solvent was evaporated. The residue, as well as the separated crystals, were subjected to column chromatography on Kieselgel 60 (Merck, 0.063-0.2 mm) adsorbent using toluene followed by toluene/MeOH (99/1) as eluent. The purity of the collected fractions were checked by TLC (Merck, Kieselgel 60 F₂₅₄ aluminum sheets; developing solvents: dichloromethane and/or toluene/EtOH (4/1). The combined chromatographically pure fractions were evaporated to yield 3a, 3b, 3c, 4a, and 4b as red or deep-red (4a,4b) crystals. The products obtained were further purified by crystallization from hexane or toluene-hexane (4a,4b) mixtures. The physical and analytical data of the previously unpublished compounds follow:

3a: 71-72 $^\circ\text{C}$; Yield 25%, Analysis: Found: C 68.47, H 5.68, Calculated ($\text{C}_{16}\text{H}_{16}\text{FeO}$, MW: 280.15) C 68.60, H 5.76%.

3b: 115-116 $^\circ\text{C}$; Yield 51%, Analysis: Found: C 69.32, H 6.23, Calculated ($\text{C}_{17}\text{H}_{18}\text{FeO}$, MW: 294.18) C 69.41, H 6.17%.

3c: 64-66 $^\circ\text{C}$; Yield 16%, Analysis: Found: C 69.98, H 6.31, Calculated ($\text{C}_{18}\text{H}_{20}\text{FeO}$, MW: 308.20) C 70.15, H 6.54%.

Reaction of ferrocenecarboxaldehyde (2) with *E*-2-benzylidenecycloalkanones (5a-c, X=H) and their substituted derivatives. 1 mmol ferrocenecarboxaldehyde (2) and 1.2 mmol *E*-2-benzylidenecycloalkanone (5a-c, X=H) were dissolved in 10 mL of MeOH and the solution was saturated with O_2 -free N_2 -gas for 5 min. In 5-mL MeOH saturated with O_2 -free N_2 , 1 mmol KOH was dissolved and the solution was added to the MeOH solution of 2 and the respective 5a-c. The flask was sealed and kept at RT for 2 days (5a), or heated at reflux for 2 hours (5b) or 8 hours (5c) under an argon atmosphere. (Similar conditions were applied when the corresponding *E*-2-(4'-X-benzylidene)cycloalkanones [5a: X = H, CH_3 , OCH_3 , $\text{N}(\text{CH}_3)_2$; 5b: X= H, CH_3 ,

OCH₃, NO₂; **5c**: X= H, CH₃, OCH₃, N(CH₃)₂] were used instead of the unsubstituted **5a-c**). Then the mixture was placed in a refrigerator overnight and any resulting crystals (**7a**, **8a**, **9a**, **11a**) were filtered and purified by column chromatography. The filtrate (or the homogeneous reaction mixture) was evaporated at 40 °C and the residue was treated as described above. Column chromatography of the hydroxide-free residues and the separated crystals yielded **7a**, **8a**, **9a**, **11a**, **7b**, **8b**, **9b**, and **10b** as deep-red crystals. The products obtained were further purified by crystallization from hexane (**7b**, **8b**, **9b**) or toluene-hexane mixtures. The physical and analytical data of the obtained compounds follow:

7a: M.p. 173-176 °C, Yield 67%, Analysis: Found: C 74.93, H 5.38, Calculated (C₂₃H₂₀FeO, MW: 368.26) C 75.02, H 5.47%.

8a: M.p. 198-199 °C, Yield 81%, Analysis: Found: C 75.18, H 5.83, Calculated (C₂₄H₂₂FeO, MW: 382.29) C 75.41, H 5.80%.

9a: M.p. 226-228 °C, Yield 73%, Analysis: Found: C 72.41, H 5.41, Calculated (C₂₄H₂₂FeO₂, MW: 398.28) C 72.38, H 5.57%.

11a: M.p. 212-215 °C; Yield 85%, Analysis: Found: C 72.86, H 5.93, Calculated (C₂₅H₂₅FeNO, MW: 411.33) C 73.00, H 6.13%.

7b: M.p. 128-130 °C; Yield 58%, Analysis: Found: C 75.24, H 5.77, Calculated (C₂₄H₂₂FeO, MW: 382.29) C 75.41, H 5.80%.

8b: M.p. 153-155 °C; Yield 64%, Analysis: Found: C 75.65, H 6.01, Calculated (C₂₅H₂₄FeO, MW: 396.31) C 75.77, H 6.10%.

9b: M.p. 181-183 °C; Yield 62%, Analysis: Found: C 72.78, H 5.92, Calculated (C₂₅H₂₄FeO₂, MW: 412.31) C 72.83, H 5.87%.

10b: M.p. 182-184 °C. Yield 78%, Analysis: Found: C 67.24, H 4.81, Calculated (C₂₄H₂₁FeNO₃, MW: 427.28) C 67.46, H 4.95%.

Results and Discussion

Reaction of ferrocenecarboxaldehyde (**2**) with cycloalkanones **1a-c** yielded the respective mixtures of *E*-ferrocenemethylenecycloalkanones **3a-c** and *E,E*-diferrocenemethylenecycloalkanones **4a** and **4b**. While most of the formed **4a** separated from the reaction mixture, no precipitate formation was observed in the other two reactions. The soluble mixtures of unreacted **2**, **3** and **4** could be separated by column chromatography on Kieselgel 60 using toluene and toluene/MeOH (99/1) as eluent. Contrary to the reactions with cyclopentanone (**1a**) and cyclohexanone (**1b**), in which two products (**3a,b** and **4a,b**) could be isolated, reaction of **2** with cycloheptanone (**1c**) resulted in the formation of only the respective monoferrocenemethylene derivative **3c** in a rather low yield. TLC analysis of the reaction mixture showed the formation of a second product as well; however, the amount of the formed compound was insufficient for its isolation and structural characterization.

The reason for the low yield is supposed to be the relatively high stability of the intermediate cycloheptanolate, which results in a less reactive intermediate to give a good yield in an aldolic reaction.²⁵ For spectroscopic comparisons we have also prepared the corresponding *E*-2-benzylidenecycloalkanones (**5a-c**)^{12,13,14} and *E,E*-dibenzylidenecycloalkanones (**6a-c**).¹²

In order to synthesize ferrocene analogues (**7-11**) of *E,E*-dibenzylidenecycloalkanones **6a-c** we treated *E*-2-benzylidenecycloalkanones (**5a-c**) with ferrocenecarboxaldehyde (**2**) in the presence of KOH as a catalyst. Since the expected compounds (**7a-c**) are of interest from both chemical and biological points of view, we performed the reactions with some substituted *E*-2-(4'-X-benzylidene)cycloalkanones as well. In these reactions the expected mixed type ferrocene analogues of **6a** and **6b** were formed (Figure 1). In case of the reactions of **2** with **5c** and its substituted derivatives, low quantities of product were formed even after prolonged heating of the reaction mixtures, although formation of a new product could be detected on TLC. Also, no structural characterization of the products could be performed.

Because of the biological interest in the seven-membered ring derivatives, we are planning to develop modified reaction conditions that provide the expected compounds in higher yields in order to perform spectroscopic investigations and biological testing.

The ¹H and ¹³C NMR data and characteristic IR frequencies of the new compounds are given in Tables 1 and 2 and they prove the expected structures straightforwardly. Only a few additional remarks are necessary.

The expected *E*-configuration of the compounds was based on comparisons of their ¹H and ¹³C NMR data with previously published data of **5** and **6**^{5,13} and those of the respective *E*-2-benzylidene- and *E*-2-ferrocenmethylenebenzocycloalkanones.^{10,11} The stereohomogeneous *E,E*-configuration of **4a** and **6a-c** is proved by the symmetrical pattern of their NMR spectra. The change in substituents bonded to the cycloalkanone ring (arylmethylene or ferrocenylmethylene) has no significant influence on the structures and spectral characteristics in the new compounds.

In accordance with our earlier investigations the stereostructure (conformation) of *E*-2-benzylidene- and *E*-2-ferrocenmethylenebenzocycloalkanones depend on the size of the cycloalkanone ring.^{10,11} Similar to the respective benzocycloalkanones, in the series **6** (dibenzylidenecycloalkanones) the molecule with 5-membered ring (**6a**) is almost fully planar: in the solid state the two benzylidene moieties subtend an angle of 10°. ²⁶ In the cyclohexanone (**6b**) and the cycloheptanone (**6c**) derivatives the molecules are more distorted and the conformers with out of plane aryl group to the C=C double bonds are preferred.^{27,6} The stereostructure is practically the same in the **4** and the **7-11**-type dienones.

In the monosubstituted cycloalkanones **5** the cinnamoyl moiety is distorted even in the molecules of **5a** with 5-membered ring: in the solid state the plane of the aryl ring deviates from that of the C=C double bond by 23°. ²⁸ Similar conclusions could be drawn from the ¹³C NMR investigation of a series of *E*-(4'-X-benzylidene)cyclohexanones.¹³ The similar spectral characteristics of **5** and **3** reflect a similar stereostructure for the corresponding **3**.

In all the molecules the chemical equivalence of H/C-2',5' and H/C-3',4' atom pairs of the substituted Cp rings as well as the H/C-2,6 and H/C-3,5 atom pairs in the aryl groups proves a

fast interchange of coplanar conformations – the free internal rotation around the C–C bond binding the cyclic enone moiety to the ferrocenyl or aryl substituent.

The electron-reservoir character of the ferrocenyl moiety compensates the electron deficiency due to enone-polarisation around the C- β atom, and as a consequence, the chemical shift of =C(Fc) atom is higher in **b**- and **c**-type compounds where this effect is not present in the non-planar conformations.

In accordance with our previous results with the respective benzocyanones,^{10,11} the characteristic higher C=O IR frequency²⁹ and ¹³C NMR chemical shift^{17b} of the cyclopentanones are observed in the series **a** (see Tables 1 and 2).

Acknowledgements

We express our thanks to the Hungarian Research Found (OTKA, grant T-29651), and to the Ministry of Education (grant FKFP 0200/2000) for financial support.

References

1. Fülöp, F.; Bernáth, G.; Pihlaja, K. *Adv. Heterocycl. Chem.* **1998**, *69*, 349.
2. Perjési, P.; Földesi, A.; Batta, G.; Tamás, J. *Chem. Ber.* **1989**, *122*, 651.
3. Perjési, P.; Sohár, P. *Monatsh. Chem.* **1991**, *122*, 1047.
4. Markaverich, B. M.; Schauweker, T. H.; Gregory, R. R.; Varma, M.; Kittrell, F. S.; Medina, D.; Varma, R. S. *Cancer Res.* **1991**, *52*, 2482.
5. Dinkova-Kostova, A. T.; Abeygunawardana, C.; Talalay, P. *J. Med. Chem.* **1998**, *41*, 5287.
6. Artico, M.; Di Santo, R.; Costi, R.; Novellino, E.; Greco, G.; Massa, S.; Tramontano, E.; Marongiu, M. E.; De Montis, A.; La Colla, P. *J. Med. Chem.* **1998**, *41*, 3948.
7. Shaw, K. J.; Guilford, W. J.; Dallas, J. L.; Koovakkat, S. K.; McCarick, M. A.; Liang, A.; Light, D. R.; Morrissey, M. M. *J. Med. Chem.* **1998**, *41*, 3551.
8. Herberhold, M.; Nuyken, O.; Pöhlmann, T. *J. Organomet. Chem.* **1995**, *501*, 13.
9. Top, S.; Vessieres, A.; Cabestaing, C.; Laios, I.; Leclercq, G.; Provot, C.; Jaouen, G. *J. Organomet. Chem.* **2001**, *637-639*, 500.
10. Perjési, P.; Nusser, T.; Tarczay, G.; Sohár, P. *J. Mol. Struct.* **1999**, *479*, 13.
11. Sohár, P.; Perjési, P.; Törnroos, P. W.; Husebye, S.; Vértes, A.; Vankó, G.; Bozak, R. E. *J. Mol. Struct.* **2000**, *524*, 297.
12. Lóránd, T.; Szabó, D.; Neszmélyi, A. *Acta Chim. Acad. Sci. Hung.* **1997**, *93*, 51.
13. Geribaldí, S.; Girault, Y.; Perjési, P.; Tortajada, J. *Spectrochim. Acta Part A - Molecular Spectroscopy* **1992**, *48A*, 879.

14. Dimmock, J. R.; Kandepu, N. M.; Nazarali, A. J.; Kowalchuk, T. P.; Motaganahalli, N.; Quail, J. W.; Mykytiuk, P.; Audette, G. F.; Prasad, L.; Perjési, P.; Allen, T. M.; Santos, C. L.; Szydłowski, J.; De Clercq, E.; Balzarini, J. *J. Med. Chem.* **1999**, *42*, 1358.
15. Bozak, R. E.; Hicks, R. J.; Rennels, R.; Husebye, S.; Maartmann-Moe, K. *CPS: orgchem/0010003*. (<http://preprint.chemweb.com/orgchem/0010003>) (2000).
16. Noggle, J. H.; Schirmer, R. E. *The Nuclear Overhauser Effect*. 1971, Academic Press, New York.
17. Sohár, P. *Nuclear Magnetic Resonance Spectroscopy. Vol. 1-3*. 1983, CRC Press, Boca Raton, Florida, Vol. 2, p. (a) 89; (b) 197.
18. Sanders, J. K. M.; Mersch, J. D. *Prog. Nucl. Magn. Reson.* **1982**, *15*, 353.
19. Pegg, D. T.; Doddrell, D. M.; Bendall, M. R. *J. Chem. Phys.* **1982**, *77*, 2745.
20. Bendall, M. R.; Doddrell, D. M.; Pegg, D. T.; Hull, W. E. *High Resolution Multipulse NMR Spectrum Editing and DEPT*. 1982, Bruker, Karlsruhe.
21. Ernst, R. R.; Bodenhausen, G.; Wokaun, A. *Principles of Nuclear Magnetic Resonance in one and Two Dimensions*. 1987, Clarendon Press, Oxford, U.K. p. a) 400. (b) 471.
22. Sanders, J. K. M.; Hunter, B. K. *Modern NMR Spectroscopy. A Guide for Chemists*. **1987**, University Press, Oxford, U. K. p. (a) 108. (b) 94, 100.
23. Bax, A.; Morris, G. *J. Magn. Res.* **1981**, *42*, 501.
24. Kessler, H.; Griesinger, C.; Zarboch, J.; Loosli, H. *J. Magn. Res.* **1984**, *57*, 331.
25. Simeon, F.; Ghourand, F.; Barré, L. *CPS: orgchem/0009008*. (<http://preprint.chemweb.com/orgchem/0009008>) (2000).
26. Hopkin, S. E.; Muir, M.; Theocharis, C. R. *J. Chem. Soc., Perkin Trans. 2* **1991**, 1131.
27. Dimmock, J. R.; Arora, V. K.; Wonko, S. L.; Hamon, N. W.; Quail, J. W.; Jia, Z.; Warrington, R. C.; Fang, W. D.; Lee, J. S. *Drug Design and Delivery* **1990**, *6*, 183.
28. Kearsley, S. K.; Desiraju, G. R. *Proc. R. Soc. London* **1985**, *A397*, 157.
29. Holly, S.; Sohár, P. *Theoretical and Technical Introduction to the Series "Absorption Spectra in the Infrared Region."* (Ed.: Láng, L., Collaborators: Holly, S.; Sohár, P.). 1975, Akadémiai Kiadó, Budapest, pp. 93 and 98.