

A new aryl-naphthalene type lignan from *Cordia rufescens* A. DC. (Boraginaceae)

Sâmia Andrícia Souza da Silva,^a Augusto Lopes Souto,^a Maria de Fátima Agra,^a Emídio Vasconcelos Leitão da-Cunha,^{a,b} José Maria Barbosa-Filho,^a Marcelo Sobral da Silva,^{a*} and Raimundo Braz-Filho^c

^a Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, Cx. Postal 5009, 58051-970, João Pessoa, PB, Brasil. ^b Departamento de Farmácia e Biologia, Universidade Estadual da Paraíba, CCBS, 58100-000, Campina Grande, PB, Brasil. ^c Setor de Química de Produtos Naturais- LCQUI-CCT- Universidade Estadual do Norte Fluminense, 28013-600, Campos, RJ, Brasil

E-mail: marcelosobral@ltf.ufpb.br

Dedicated to Prof. Otto Richard Gottlieb

(received 28 Jan 04; accepted 06 July 04; published on the web 30 July 04)

Abstract

A new aryl-naphthalene type lignan named rufescidride, the first containing an unusual anhydride moiety, has been isolated from the stem and branches of *Cordia rufescens*. Its structure was elucidated on the basis of spectral data (IR, MS and NMR), mainly 1D and 2D NMR.

Keywords: *Cordia rufescens*, Boraginaceae, aryl-naphthalene lignan, rufescidride

Introduction

The Boraginaceae family comprises about 130 genera and 2600 species distributed in temperate and tropical zones.¹ Terpenes, pyrrolizidine alkaloids, flavonoids and naphthoquinones have been frequently reported in this family. However, only 15 lignoids have been reported: lithospermic acid, lithospermic acid B, radosiin,² arnebia lignan caffeate 3, arnebia lignan caffeate 4, arnebia caffeate 5,³ epi-radosiin,⁴ iso-radosiin,⁵ magnesium lithospermic acid,⁶ buddlenol B, ehletianol C, ehletianol D, icariside E-5,⁷ iso-salvianolic acid and salvianolic acid F.⁸ Among all the lignoids above, only the magnesium salt of lithospermic acid was isolated from a species of the genus *Cordia*. *Cordia rufescens* A. DC. (Syn: *C. piauiensis* Fresen) is a shrub popularly known in Northeastern Brazil as “ramela de velho”. Some plants of the genus *Cordia* have been used in popular medicine as abortive,⁹ anti-inflammatory,¹⁰ and to treat dysmenorrhea and dyspepsia.¹¹ Only 3β-O-[α-L-rhamnopyranosyl-(1→2)-β-D-glucopyranosyl]

ursolic acid 28-*O*-[β -D-glucopyranosyl-(1 \rightarrow 6)- β -D-glucopyranosyl] ester were reported from *C. rufescens*.¹² As a part of the study of the genus *Cordia* carried out by our group, this paper reports the isolation of a new aryl-naphthalene type lignan, named rufescidride, from the stem and branches of *C. rufescens*.

Results and Discussion

The dried and ground stem and branches of *C. rufescens* were exhaustively extracted with EtOH at room temperature. The crude EtOH extract was fractionated with hexane, CHCl₃ and AcOEt. The chloroform fraction was subjected to column chromatography over silica gel yielding 3 β -*O*- β -D-glucopyranosyl- β -sitosterol. The AcOEt fraction was subjected to successive chromatographic procedures on Sephadex LH-20 to yield the new aryl-naphthalene lignan (1).




Figure 1. Rufescidride, rufescidride triacetate and the fragment attributed to the peak at *m/z* 264 observed in the EIMS.

The molecular formula C₁₈H₈O₇ for 1 was deduced by analysis of the EIMS, ¹H NMR and APT-¹³C NMR spectra (Table 1). The presence of carbonyl functions belonging to the anhydride group was revealed by two absorptions at ν_{\max} 1798 and 1737 cm⁻¹ observed in the IR spectrum, in agreement with the signals at δ_C 163.67 (correlated in the HMBC spectrum with the signal of H-7 at δ_H 8.14) and 163.38 in the ¹³C NMR spectra (Table 1). This deduction was supported by the presence of the base peak at *m/z* 264, attributed to fragment 1a derived from the molecular ion at *m/z* 336 by loss of CO₂ + CO (C₂O₃ = 72). The ¹H NMR spectrum (200 MHz) of 1 (Table 1) showed the presence of three singlet signals at δ_H 8.81 (s, H-6'), 8.14 (s, H-7) and 6.73 (s, H-3') and two doublets at δ_H 7.68 (d, *J* = 8.8 Hz, H-6) e 7.47 (d, *J* = 8.8 Hz, H-5). The existence of three hydroxyl groups was confirmed by ¹H NMR of its acetyl derivative 2, which revealed the presence of three acetyl signals at δ_H 2.44 (s, 3H), 2.35 (s, 3H) e 2.33 (s, 3H). Comparative analysis of the {¹H}- and APT-¹³C NMR spectra (Table 1) showed eighteen signals: five for

methine aromatic carbons and thirteen for non-hydrogenated carbons [including five oxygenated: δ_C 138.55 (C-3), 143.79 (C-4), 151.58 (C-2'), 147.30 (C-4') and 142.15 (C-5')].

The 2D HMQC and HMBC spectra were also used to attribute the structure 1 and to complete ^1H and ^{13}C chemical shift assignments, summarized in Table 1. The cross correlations of C-1' (δ_C 109.06) with H-3' (δ_H 6.73, $^3J_{\text{CH}}$) and C-2' (δ_C 151.58) with both H-3' (δ_H 6.73, $^2J_{\text{CH}}$) and H-6' (δ_H 8.81, $^3J_{\text{CH}}$) observed in the HMBC spectrum suggested the location of the hydrogens H-3' and H-6' in the same aromatic ring. Analogously way, the correlations between hydrogen H-7 (δ_H 8.14) and both carbon atoms C-2 (δ_C 110.82, $^3J_{\text{CH}}$) and C-9 (δ_C 163.67, $^3J_{\text{CH}}$) were used to locate the anhydride function. All the HMBC correlations are summarized in Figure 2.

The analysis of the spectral data (IR, MS and 1D and 2D NMR), comparison with literature values described for yunnaneic acid¹³ and the significant deshielding revealed by the signals corresponding to H-5 ($\Delta\delta_H = 7.56 - 7.47 = 0.09$), H-7 ($\Delta\delta_H = 8.24 - 8.14 = 0.10$), H-3' ($\Delta\delta_H = 7.22 - 6.73 = 0.49$) and H-6' ($\Delta\delta_H = 9.56 - 8.81 = 0.75$) in the ^1H NMR spectrum of 2 established the structure 1, a new lignan of the arylnaphthalene type named rufescidride.




Figure 2. Heteronuclear correlations $^2J_{\text{CH}}$ and $^3J_{\text{CH}}$ (HMBC) for rufescidride.

Table 1. ^1H (200 MHz) and ^{13}C (50 MHz) NMR (1D and 2D) spectral data for compound 1

C	δ_C	HMQC	HMBC	
		δ_H	$^2J_{\text{CH}}$	$^3J_{\text{CH}}$
1	128.03			H-5
2	110.82			H-7
3	138.55			H-5
4	143.79			H-6
5	121.08	7.47 (d, $J = 8.8$ Hz)		
6	122.83	7.68 (d, $J = 8.8$ Hz)		
7	124.38	8.14 (s)		H-6
8	123.77			
9	163.67			H-7
1'	109.06			H-3'

Table 1. Continued

2'	151.58		H-3'	H-6'
3'	103.06	6.73 (s)		
4'	147.30		H-3'	H-6'
5'	142.15		H-6'	H-3'
6'	114.82	8.81 (s)		H-6'
7'	133.35			H-7
8'	125.38			
9'	16338			

Experimental Section

General Procedures. IR spectra were recorded on a PERKIN-ELMER FT-IR. ^1H and ^{13}C NMR were measured on a MERCURY VARIAN spectrometer at 200 MHz using DMSO- D_6 (**1**) or CDCl_3 (**2**) as solvent and TMS as internal standard or by reference to the solvent signal ($\text{CD}_2\text{HSOCD}_3$ at δ_{H} 2.50 or CHCl_3 at δ_{H} 7.24 and CD_3SOCD_3 at δ_{C} 39.5 or CDCl_3 and at δ_{C} 77.00). EIMS were obtained at 70 eV on a Shimadzu QP-2000 spectrometer.

Plant material. Stem and branches of *C. rufescens* were collected in August 2002 in the city Cruz do Espírito Santo, State of Paraíba, Brazil. A voucher specimen has been deposited at the Herbarium Prof. Lauro Pires Xavier (JPB) in the Universidade Federal da Paraíba.

Extraction and isolation. The stem and branches of *C. rufescens* (7000g), air-dried and powdered, were exhaustively extracted with EtOH at room temperature. The crude extract was taken up in MeOH:H₂O (7:3) and extracted successively with hexane, CHCl_3 and EtOAc (20 g). 10 g of the AcOEt fraction was subjected to column chromatography over Sephadex LH-20 using MeOH, resulting in 27 fractions. Fractions 6-8 (0,200 g) were reunited and subjected to successive CC over Sephadex LH-20 to yield compound **1** (0,017 g) as a red amorphous powder with melting point at 327-330 °C.

Acetyl derivative (2, rufescidride triacetate). Acetylation of **1** (Ac_2O , py) yielded **2** as a yellow amorphous powder (0,010 g) with melting point at 236-238 °C. δ_{H} (CDCl_3 , 200 MHz) 2.33 (3H, s), 2.35 (3H, s), 2.44 (3H, s), 7.22 (1H, s), 7.56 (1H, d, J = 8.8 Hz), 7.68 (1H, d, J = 8.8 Hz), 8.24 (1H, s), 9.56 (1H, s).

Acknowledgements

The authors express their gratitude to Vicente Carlos de Oliveira Costa and Wellington de Lima Navarro, Laboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, for providing the NMR and IR spectra, respectively, and CNPq/IMSEAR and FAPERJ for financial support. Sincere thanks are also due to NAPRALERT.

References

1. Brummitt, R.K. *Vascular Plant Families and Genera*; Royal Botanic Garden: Kew, 1992.
2. Yamamoto, H.; Inoue, K.; Yazaki, K. *Phytochemistry* **2000**, *53*, 651.
3. Kashiwada, Y.; Nshizawa, M.; Yamagishi, T.; Tanaka, T.; Nonaka, G.; Cosentino, L.M.; Zinder, J.V.; Lee, K.H. *J. Nat. Prod.* **1995**, *58*, 392.
4. Nishizawa, M.; Tsuda, M.; Hayashi, K. *Phytochemistry* **1990**, *29*, 2645.
5. Kashiwada, Y.; Bastow, K.F.; Lee, K.H. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 905.
6. Matsuse, I.T.; Lim, Y.A.; Hattori, M.; Correa, M.; Gupta, M.P. *J. Ethnopharmacol.* **1999**, *64*, 15.
7. Yoshikawa, K.; Kageyama, H.; Arihara, S. *Phytochemistry* **1995**, *39*, 659.
8. Lin, Y.L.; Chang, Y.Y.; Kuo, Y.H.; Shiao, M.S. *J. Nat. Prod.* **2002**, *65*, 745.
9. Yu, G.D. *Chung Yao Tung Pao* **1982**, *7*, 6.
10. Sertie, J.A. A.; Basile, A. C.; Panizza, S.; Oshiro, T. T.; Azzolini, C. P.; Penna, S. C. *J. Ethnopharmacol.* **1991**, *31*, 239.
11. Asprey, G.F.; Thornton, P. *West Indian Med. J.* **1955**, *4*, 69.
12. Santos, R.P.; Viana, F.A.; Lemos, T.L.G.; Silveira, E.R.; Braz-Filho, R.; Pessoa, O.T.L. *Magn. Reson. Chem.* **2003**, *41*, 735.
13. Tanaka, T.; Nishimura, A.; Kouno, I.; Nonaka, G.; Yang, C.R. *Chem. Pharm. Bull.* **1997**, *45*, 1596.