

A bile acid derived potassium ion sensor

Uday Maitra* and Suvadeep Nath

*Department of Organic Chemistry, Indian Institute of Science,
Bangalore 560 012, India*

** Also at the Chemical Biology Unit, JNCASR, Bangalore 560 064, India*

E-mail: maitra@orgchem.iisc.ernet.in

Dedicated to Dr. A. V. Rama Rao on the occasion of his 70th birthday

(received 15 Oct 04; accepted 14 Dec 04; published on the web 15 Dec 04)

Abstract

A chenodeoxycholic acid based K⁺ ion sensor has been designed using a modular approach in which a fluorophore and a cation receptor are attached to the bile acid backbone. In the absence of K⁺ the fluorescence of the molecule is quenched because of through-space, photo-induced electron-transfer from the aza-crown unit. Fluorescence enhancement was observed upon titration with K⁺ (and other alkali metal ions too). In methanol, good selectivity towards the sensing of K⁺ has been observed.

Keywords: Photo-induced electron transfer (PET), fluorescence enhancement, aza-crown, bile acid, cation-sensor


Introduction

The design of ion sensors based on photo-induced electron-transfer (PET) mechanism are attracting interest in recent years for their applications in fields like analytical, medical science, and molecular electronics.¹ Such a sensor molecule combines a fluorophore and a receptor module, and the sensing of metal ion relies typically on the enhancement of fluorescence when the cation binds to the receptor module. In an azacrown ether - spacer - fluorophore system, the nitrogen lone pair quenches the fluorescence of the fluorophore. When a metal ion binds, the nitrogen lone pair is engaged through coordination to the cation causing fluorescence enhancement. The efficiency of PET process depends on the distance between the quencher moiety and the fluorophore. The effect of spacer length on the emission properties of a pyrene fluorophore upon alkali metal ion complexation by 1-aza-18-crown-6 has been studied extensively, which showed that the efficiency of the PET process was higher for shorter spacers.²

Bile acids contain a varying number of OH groups on the α -face of the molecule positioned at distances appropriate for the design of a cation sensor.³ We have designed a bile acid based sensor molecule **3** in which the pyrene fluorophore and the aza-18-crown-6 receptor are linked formally *via* a 1, 5-diol. The rigidity of the steroid backbone allows the fluorophore and the receptor module to be in close proximity.

Results and Discussion

Methyl chenodeoxycholate was selectively acylated with pyrene-1-carboxylic acid at the 3 α position to yield **1** (43%), which was converted to its chloroacetyl derivative **2** (65%), and finally reacted with 1-aza-18-crown-6 to form the desired product **3** in 55% yield (**Scheme 1**).



Scheme 1

Binding constants for the association of different alkali metal cations and NH_4^+ with **3** in CHCl_3 at 20 °C were determined by Cram's picrate extraction method,⁵ and the estimated

binding constants are, $\log K_a$ (K^+) 7.47; $\log K_a$ (NH_4^+) 6.75; $\log K_a$ (Na^+) 6.11; $\log K_a$ (Li^+) 4.49. This order is in accordance with published data.⁶

Fluorescence titration experiments were subsequently carried out using 5 μM of **3** in 4:1 toluene/acetonitrile. $KClO_4$, NH_4PF_6 , $NaBF_4$, and $LiClO_4$ were used as analytes in varying concentrations. In all the cases the fluorescence intensity increased with the guest concentrations and reached a saturation value at $\sim 5 \mu M$ suggesting 1:1 complexation (**Figure 1**).




Figure 1. Fluorescence titration of 5 μM soln. of **3** (4:1 PhMe/MeCN) with $KClO_4$.

The increase in the fluorescence intensity differed slightly with the cation. There was a small difference in the initial slopes of the curves when the fluorescence enhancement was plotted against the salt concentration. There was a gradual decrease in the slopes as the following order $K^+ \sim NH_4^+ > Na^+ > Li^+$ (**Figure 2**).




Figure 2. Increase in fluorescence intensity of **3** in 4:1 toluene/acetonitrile with added salt, showing 1:1 complexation I, I_0 are fluorescence intensities in the presence and absence of salt.

To explore the difference in the fluorescence enhancement for different guests, the same experiment was performed at a lower sensor concentration ($0.2 \mu\text{M}$). As expected, different slopes for the increment of the fluorescence intensity with the concentration of the cations was observed with K^+ showing a higher slope compared to Li^+ because of its higher binding affinity (**Figure 3**).




Figure 3. Increase in fluorescence intensity of **3** with added salt in 4:1 PhMe/MeCN.

Since the difference in the slopes in 4:1 toluene/ acetonitrile for different cations was not significantly different, the same experiment was performed in MeOH to enhance the selectivity.

When a 1 μM solution of **3** in MeOH was titrated with cations a sharp increase in the fluorescence intensity was observed with K^+ , while Na^+ showed significantly less enhancement and Li^+ didn't show any enhancement upto 10 μM of LiClO_4 (**Fig. 4**).




Figure 4. Increase in fluorescence intensity of **3** with various guests in MeOH, and the sensitivity is high when the guest is K^+ .

To verify the fluorescence enhancement behavior for a mixture of K^+ and Na^+ ions, competition experiments were done by varying the concentrations of Na^+ and K^+ in MeOH. At low $[\text{K}^+]$ a small fluorescence enhancement due to the addition of Na^+ could be observed. On the other hand, at higher potassium concentration the effect of sodium was insignificant because of the saturation of the binding sites (**Figure 5**). When the fluorescence enhancement was plotted against $[\text{K}^+]$ at different constant $[\text{Na}^+]$, a lowering of the slope was observed at higher $[\text{Na}^+]$ which indicated that the influence of added Na^+ was predominant only at low K^+ concentration (**Figure 6**). The data from Fig. 5 and 6 are presented in a 3D format in **Figure 7**. It is noteworthy that 1 μM of K^+ could be sensed in the presence of 40 μM of Na^+ .




Figure 5. Change in the fluorescence intensities of **3** with Na^+ keeping K^+ concentration constant in MeOH.




Figure 6. Change in the fluorescence intensities of **3** with K^+ keeping Na^+ concentration constant in MeOH.




Figure 7. Relative fluorescence enhancement of **3** in MeOH for mixtures of K⁺ and Na⁺.

The increase in the fluorescence intensity upon cation binding can be explained on the basis of a PET mechanism (**Figure 8**).




Figure 8. Quenching of fluorescence due to PET and enhancement of pyrene fluorescence upon binding to cations.

To confirm the phenomena as a through-space PET process, analog **4**³ (**Figure 9**) was tested under identical conditions. A methanolic solution of **4** at 1 μM showed a *higher* fluorescence intensity compared to **3** at the same conc. and didn't increase much upon the addition of KClO_4 (**Figure 10**) or NaClO_4 . Unlike **3**, the distance between the two modules is larger in **4**, and thus the pyrene fluorescence was not quenched. Another control experiment with non-covalently linked fluorophore and the aza-crown receptor (each at 1 μM) showed no change in the fluorescence intensity in the presence of alkali metal ions, suggesting that an appropriate geometry and distance is a prerequisite for the 'through space photo induced electron transfer' process.




Figure 9. Structural analog of **3**.




Figure 10. The fluorescence enhancement with $[\text{K}^+]$ for **3** is higher as compared to that of **4**, while a 1:1 mixture of methyl pyrene-1-carboxylate and 1-aza-18-crown-6 does not show any enhancement with K^+ .

Conclusions

In conclusion, we have synthesized a bile acid based PET sensor for alkali metal ions, where PET occurs through space and the fluorescence quenching process is inhibited upon binding to alkali metal ions. Using this sensor, $<0.2 \mu\text{M}$ of K^+ can be determined in 4:1 toluene/acetonitrile, and K^+ can selectively be sensed in MeOH. Currently we are exploring the synthesis of polymer bound analogs of the sensor to examine the detection of ions in aqueous fluids. We believe that since the synthesis of the sensor is modular, one can envision designing other sensors by using different receptor and/or sensor modules to attach to the 3 α and 7 α positions of chenodeoxycholic acid.

Experimental Section

General Procedures. All melting points were checked in Bñchi B-540 melting point apparatus. TLC was done on pre-coated silica gel plates (Merck) and stained with Liebermann Buchard reagent or observed under long/shortwave UV or in iodine vapor. Column chromatographic purifications were carried out on 100-200 mesh silica gel (Acme) using gravity columns. UV-Vis, IR and fluorescence spectra were recorded on Shimadzu UV2100, JASCO-70 FT-IR and Perkin-Elmer LS-50B spectrometers, respectively. NMR spectra were recorded on a 300 MHz (JEOL Lambda-300) and 500 MHz (BRUKER DRX 500) instruments in deuterated solvents as indicated; TMS or the residual solvent peaks were used as internal standards. Optical rotations were measured at 589 nm at 24°C on a JASCO DIP-370 digital polarimeter. Micro analyses were done on a Carlo Erba Strumentazione CHNS Analyser-model1106 and Flash EA 1112. MALDI-TOF-MS was done in KRATOS KOMPACT MALDI 4. LRMS and HRMS were recorded on Micromass Q-Tof micro.

Methyl 3 α -(1-pyrenecarbonyloxy)-7 α -hydroxy-5 β -cholan-24-oate (1). To a dichloromethane (24 mL) suspension of pyrene-1-COOH (1.16 g, 4.72 mmol), oxalyl chloride (1 mL, 11.5 mmol) was added and stirred at room temperature for 45 min. Excess oxalyl chloride was removed from the reaction mixture *in vacuo*. To a solution of methyl 3 α , 7 α -dihydroxy-5 β -cholan-24-oate (1.73 g, 4.25 mmol) in dry dichloromethane (15 mL), triethylamine (3 mL), DMAP (0.22 g, 1.8 mmol), and the freshly prepared solution of pyrene-1-carbonyl chloride in dichloromethane (20 mL) was added and the mixture was stirred at room temperature for 7 h. Solvent was removed from the reaction mixture and the residue was dissolved in chloroform (100 mL) and washed with satd. NaHCO_3 (100 mL), water (100 mL) and finally with brine (100 mL). The clear organic layer was dried over anhyd. Na_2SO_4 , and solvent was removed *in vacuo*. The crude product was chromatographed on a silica gel column (100-200 mesh, 3.1 cm x 17 cm) with 0-2% ethyl acetate/ CHCl_3 to obtain 1.17 g (43%) of the pure product; mp 128-131 °C; $[\alpha]_{\text{D}}^{24}$ 31 (c. 1.0, CHCl_3); IR (KBr, cm^{-1}) 3558, 2937, 2867, 1737, 1689, 1251, 1231; $^1\text{H-NMR}$ (300 MHz, CDCl_3)

δ 0.69 (s, 3H); 0.94 (d, $J = 6.0$ Hz, 3H); 1.00 (s, 3H); 1.17- 2.62 (m, steroidal CH and CH₂); 3.67 (s, 3H); 3.91 (s, 1H); 5.05 (m, 1H); 8.04-8.25 (m, 7H, aromatic H of pyrene); 8.63 (d, $J = 8.1$ Hz); 9.26 (d, $J = 9.1$ Hz). ¹³C-NMR (75 MHz, CDCl₃) δ 11.66; 18.18; 20.54; 22.70; 23.59; 26.99; 28.06; 30.89; 30.96; 32.80; 34.46; 35.09; 35.12; 35.28; 35.48; 39.30; 39.46; 41.35; 42.59; 50.35; 51.44; 55.66; 68.38; 75.31; 124.00; 124.13; 124.33; 124.70; 124.90; 125.95; 126.04; 126.12; 127.09; 128.36; 129.16; 129.31; 130.30; 130.86; 130.91; 133.98; 167.62; 174.71. HRMS: Calcd for (M⁺+ Na) 657.35558; Found: 657.3556; *Anal.* Calcd for C₄₂H₅₀O₅: C 79.45; H 7.94. Found: C 79.40; H 7.91.

Methyl 3 α -(1-pyrenecarbonyloxy)-7 α -chloroacetyloxy-5 β -cholan-24-oate (2). To a solution of **2** (1.03 g, 1.62 mmol) in toluene (15 mL), CaH₂ (0.20 g, 4.72 mmol), nBu₄N⁺I⁻ (0.17 g, 0.47 mmol) and chloroacetyl chloride (700 μ L, 8.79 mmol) were added and the mixture was refluxed for 12 h. The reaction mixture was filtered through celite and the residue was washed with ethyl acetate (100 mL). The filtrate was washed with satd. NaHCO₃ solution (100 mL), water (100 mL), brine (100 mL) and finally dried over anhyd. Na₂SO₄ and filtered. The solvent was removed *in vacuo* to yield the crude product (1.23 g), which was purified by column chromatography on silica gel (100-200 mesh, 30 cm x 3 cm, 1.23 g) using 4-10% EtOAc/hexanes as the eluent. The pure product weighed 1 g (65%); mp 163.1-166.4 °C; [α]_D²⁴ 7 (c. 1.0, CHCl₃); IR (KBr, cm⁻¹) 3443, 2942, 2869, 1735, 1695, 1255, 1133; ¹H-NMR (300 MHz, CDCl₃) δ 0.68 (s, 3H); 0.93 (d, $J = 6$ Hz, 3H); 1.02 (s, 3H); 1.12- 2.4 (m, steroidal CH and CH₂); 3.66 (s, 3H); 4.03 (d, $J = 15$ Hz, 1H); 4.05 (d, $J = 15$ Hz, 1H); 5.05 (br. m, 2H); 8.04- 8.31 (m, aromatic protons of pyrene, 7H); 8.60 (d, $J = 8.1$ Hz, 1H); 9.210 (d, $J = 9.3$ Hz, 1H). ¹³C-NMR (75 MHz, CDCl₃) δ 11.51; 18.13; 20.46; 22.49; 23.42; 26.89; 27.83; 30.76; 30.81; 31.19; 33.93; 34.67; 34.79; 34.82; 35.10; 37.87; 39.20; 40.76; 41.17; 42.55; 50.07; 51.35; 55.52; 73.74; 74.78; 123.95; 123.98; 124.16; 124.59; 124.69; 125.91; 126.04; 126.07; 126.96; 128.09; 129.10; 129.28; 130.15; 130.69; 130.79; 133.93; 166.31; 167.44; 174.48. LRMS Calcd for (M⁺+ Na) 733.3; found: 733; Calcd for (M⁺+ K) 749.3; Found: 749. *Anal.* Calcd for C₄₄H₅₁ClO₆: C 74.29; H 7.23. Found: C 74.24; H 7.07.

Methyl 3 α -(1-pyrenecarbonyloxy)-7 α -((1-((1-aza-4,7,10,13,16-pentaoxa)-cyclooctadecanyl)-methyl)carbonyloxy)-5 β -cholan-24-oate (3). To a mixture of **3** (0.35 g, 0.49 mmol) and Na₂CO₃ (0.36 g, 3.42 mmol) in dry CH₃CN, 1-aza-18-crown-6 (0.21 g, 0.8 mmol) was added and the mixture was refluxed under nitrogen for 3 d. The reaction mixture was filtered through celite, washed with EtOAc (80 mL) and the filtrate was washed with satd. NaHCO₃ solution (120 mL), dried over anhyd. Na₂SO₄, filtered and solvent was removed *in vacuo* to get the crude product (0.50 g), which was purified by column chromatography on silica gel (100-200 mesh) using 1-10% MeOH/CHCl₃ as the eluent followed by PTLC using 20% MeOH/EtOAc as the eluent to obtain 0.18 g (51%) pure product; mp 99.0-104.2 °C; [α]_D²⁴ 4.5 (c. 2.0, CHCl₃); IR (KBr, cm⁻¹) 3435, 2930, 2869, 1735, 1706; ¹H-NMR (500 MHz, CDCl₃) δ 0.70 (s, 3H); 0.96 (d, $J = 6.5$ Hz, 3H); 1.04 (s, 3H); 1.09- 2.41 (m, steroidal CH and CH₂); 2.91 (s, 3H); 3.31-3.54 (m, aza-crown CH₂ and acetyl CH₂); 3.69 (s, 3H); 4.98 (s, 1H); 5.08 (m, 1H); 8.08- 8.30 (m, aromatic protons from pyrene, 7H); 8.62 (d, $J = 8.5$ Hz, 1H); 9.25 (d, $J = 9.5$ Hz, 1H). ¹³C-NMR (75 MHz, CDCl₃)

δ 11.46; 18.06; 20.45; 22.49; 23.36; 26.86; 27.77; 30.71; 31.24; 33.92; 34.64; 34.76; 34.89; 35.04; 37.72; 39.18; 40.79; 42.42; 50.07; 51.24; 53.62; 55.50; 55.81; 69.66; 70.12; 74.78; 123.91; 123.98; 124.51; 125.89; 126.00; 126.07; 126.88; 127.98; 129.11; 129.26; 130.07; 130.64; 130.69; 133.85; 167.26; 171.03; 174.38; HRMS Calcd for (M^+ + H) 938.5418; found: 938.5420; *Anal.* Calcd for $C_{56}H_{75}NO_{11} \cdot 5H_2O$: C 65.41; H 8.33; N 1.36. Found C 65.23; H 7.87; N 1.42.

Acknowledgements

This work was supported by a research grant from the Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore.

References

- (a) de Silva, A. P.; Gunaratne, H. Q. N.; Gunnlaugsson, T.; Huxley, A. J. M.; McCoy, C. P.; Rademacher, J. T.; Rice, T. E. *Chem. Rev.* **1997**, *97*, 1515. (b) Xu, X.; Xu, H.; Ji, H. *Chem. Commun.* **2001**, 2092. (c) He, H.; Mortellaro, M.A.; Leiner, M.J.P.; Young, S.T.; Fraatz, R.J.; Tusa, J.K. *Anal. Chem.* **2003**, *75*, 549. (d) Benco, J.S.; Nienaber, H.A.; Dennen, K.; McGimpsey, W.G. *J. Photochem. Photobiol. A Chem.* **2002**, *152*, 33. (e) Benco, J.S.; Nienaber, H.A.; McGimpsey, W.G. *Sens. Actuators, B* **2002**, *85*, 126. (f) Gunnlaugsson, T.; Nieuwenhuyzen, M.; Richard, L.; Thoss, V. *Tetrahedron Lett.* **2001**, *42*, 4725. (g) Ghosh, P.; Shukla, A.D.; Das, A. *Tetrahedron Lett.* **2002**, *43*, 7419. (h) Xiao, Y.; Qian, X. *Tetrahedron Lett.* **2003**, *44*, 2087. (i) Pearson, A.J.; Xiao, W. *J. Org. Chem.* **2003**, *68*, 5361. (j) Pearson, A.J.; Xiao, W. *J. Org. Chem.* **2003**, *68*, 5369. (k) Xia, W.S.; Schmehl, R.H.; Li, C.J. *Eur. J. Org. Chem.* **2000**, 387. (l) de Silva, S.A.; Zavaleta, A.; Baron, D.E.; Allam, O.; Isidor, E.V.; Kashimura, N.; Percarpio, J.M. *Tetrahedron Lett.* **1997**, *38*, 2237. (m) Gunnlaugsson, T.; McCoy, C.P.; Morrow, R.J.; Phelan, C.; Stomeo, F. *ARKIVOC* **2003**, (vii), 216.
- Ji, H. F.; Dabestani, R.; Brown, G. M.; Hettich, R. L. *Photochem. Photobiol.* **1999**, *69*, 513.
- Maitra, U.; D'Souza, L. J.; Vijay Kumar, P. *Supramol. Chem.* **1998**, *10*, 97.
- Becker, V.; Streeck, C. *Liebigs Ann. Chem.* **1937**, *531*, 108.
- (a) Timko, J.M.; Moore, S.S.; Walba, D.M.; Hiberty, P.C.; Cram, D.J., *J. Am. Chem. Soc.* **1977**, *99*, 4207. (b) Moore, S.S.; Tarnowski, T. L.; Newcomb, M.; Cram, D. J., *J. Am. Chem. Soc.* **1977**, *99*, 6398. (c) Kyba, E. P.; Helgeson, R. C.; Madan, K.; Gokel, G. W.; Moore, S.S.; Tarnowski, T. L.; Cram, D. J., *J. Am. Chem. Soc.* **1977**, *99*, 2564.
- Gokel, G.W.; Hernandez, J.C.; Viscariello, A.M.; Arnold, K.A.; Campana, C.F.; Echegoyen, L.; Fronczek, F. R.; Gandour, R.D.; Morgan, C.R.; Trafton, J.E.; Miller, S.R.; Minganti, C.; Eiband, D.; Schultz, R.A.; Tamminen, M. *J. Org. Chem.* **1987**, *52*, 2963.