

# Montmorillonite impregnated with bismuth nitrate: a versatile reagent for the synthesis of nitro compounds of biological significance

Susanta Samajdar, Frederick F. Becker, and Bimal K. Banik\*

The University of Texas, M. D. Anderson Cancer Center, Department of Molecular Pathology,  
Box 89, 1515 Holcombe Blvd., Houston, TX 77030, USA.

*E-mail:* [banik@mdanderson.org](mailto:banik@mdanderson.org)

**Dedicated to Professor T. R. Govindachari on the occasion of his 85<sup>th</sup> birthday**  
(received 10 May 01; accepted 08 Aug 01; published on the web 16 Aug 01)

---

## Abstract

Montmorillonite impregnated with bismuth nitrate has been found to be an excellent reagent for the synthesis of several nitro compounds of biological significance in high yield.

**Keywords:** Montmorillonite, bismuth nitrate, surface mediated reactions, nitrating agents

---

## Introduction

Surface-mediated organic reactions are undergoing extensive investigation especially because they are ecologically friendly.<sup>1</sup> The usefulness of clay mediated organic synthesis has been documented in a large number of recent publications.<sup>2</sup> For example, montmorillonite which is commercially available has shown considerable promise as a solid support in carrying out different chemical reactions.<sup>2</sup> Based on our own work on bismuth<sup>3</sup> salts, we became interested in exploring trivalent bismuth nitrate as the nitrating agent under solid support. These experiments culminated in a facile synthetic method of aromatic nitration of several aromatic hydrocarbons by bismuth nitrate supported on montmorillonite which takes taking only a few minutes.<sup>4</sup> In continuation of bismuth nitrate mediated aromatic nitration, we report here the synthesis and characterization of several biologically active aromatic nitro compounds not only with polyaromatic hydrocarbons, but also with steroids and  $\beta$ -lactams.

## Results and Discussion

### Polyaromatic Nitro Compounds

Polyaromatic nitro compounds are widespread in the environment. We have been engaged in the


development of aromatic compounds as anti-tumor agents and have shown structure-activity relationships with several diamides and diamines to which a polycyclic aromatic ring was bound.<sup>5</sup> The nitro compounds required for the synthesis of these types of derivatives were prepared by the conventional nitric acid-sulfuric acid or nitronium tetrafluoroborate methods. We became concerned about the disposal of the large amount of acid-waste that resulted. To avoid this complication and the associated hazards, we set out to test the nitrating abilities of nitrate-salts under solid phase conditions in a projected route to develop a simple synthesis of the aromatic nitro compound in a relatively short period of time.

Montmorillonite clay impregnated with anhydrous cupric nitrate termed as “claycop” was used for aromatic nitration reaction.<sup>6</sup> A large excess of acetic anhydride was required when claycop was used as the nitrating agent. The actual nitrating species was believed to be acetyl nitrate. Similarly, a clay with ferric nitrate termed as “clayfen” was used as the reagent for the nitration of estrone, though yields were poor.<sup>7</sup> In addition, extreme precautions were necessary for the preparation of this reagent.<sup>8</sup> Sulfuric acid supported on silica gel that catalyzed nonselective nitration of simple aromatic compounds has also been reported.<sup>9</sup>


Mixing the starting materials with bismuth nitrate and montmorillonite with tetrahydrofuran and evaporation of the solvent under rotavapor comprise the reaction conditions for successful, regiospecific nitration with several aromatic compounds including multicyclic ring systems. The nitration of aromatic compounds by our method is shown in Scheme 1. A comparison with other solid supports, such as silica gel, acidic alumina, and ground molecular sieves was also carried out. Anisole **1**, naphthalene **2**, fluorene **3**, chrysene **4**, and dibenzofluorene **5** were regiospecifically converted to the corresponding nitro compounds **6** through **10** by bismuth nitrate-montmorillonite in excellent yield. Interestingly, the site of the electrophilic attack by this reagent was found to be identical to the conventional nitric acid or acetyl nitrate mediated nitration reaction (Scheme 1).

The reaction of naphthalene **2** using silica gel as support under identical conditions as described with montmorillonite failed to produce the nitro derivative although the same mixture upon microwave irradiation<sup>10</sup> afforded the 1-nitronaphthalene **7** in 70% yield. Acidic alumina and molecular sieves alone failed to give any product with or without acid. The results indicated that montmorillonite is the solid support of choice for aromatic nitration.

**Steroids:** At the beginning of the nitration study on steroid, a model reaction was investigated. Phenol **11** produced a mixture of 2- and 4-nitrophenol **12** and **13** in a ratio of 1:3. The formation of the mixture of nitro derivatives **12** and **13** with phenolic substrates indicated lesser selectivity under this condition (Scheme 2).




Scheme 1




Scheme 2





Scheme 4

Our results indicate that the nitration reaction depends strongly on the nature of the solid support. Further, two solid adsorbents of identical structure with different surface area (particle size) can give different product distribution. Based on a recent publication,<sup>15</sup> we believe that the binding of bismuth nitrate to the free hydroxyl group of the multi-metallic montmorillonite is important and that this complex produces the nitronium ion. Clearly, the failure of bismuth nitrate to produce any product with alumina and only 70% with silica gel (under microwave irradiation), despite the presence of surface hydroxyl groups, strengthens the importance of the composition of the solid support.



Scheme 5

In summary, we have shown a simple, rapid and convenient method for the aromatic nitration with bismuth nitrate in the presence of montmorillonite at room temperature. The superiority of this method over others includes a very rapid reaction (approximate 30 minutes instead of 24 hour) with readily available reagents. Further, our method requires no strong acid,<sup>16</sup> no acetic anhydride and it can be applied to a wide range of substrates and results in high overall yield.

## Experimental Section

**General Procedures.** All reactions described in this paper were carried out under a well-ventilated hood. IR spectra were recorded on a Perkin-Elmer Spectrum 2000 instrument. NMR spectra were recorded on Bruker 300 MHz spectrometer. Chemical shifts were reported as  $\delta$  values in parts per million downfield from t e t r a m e t h y l silane as the internal standard in  $\text{CDCl}_3$ . Elemental analysis was performed by Schwarzkopf Microanalytical Laboratory, Inc., New York. Melting points were taken in open capillary tube and are not corrected. Column chromatography was carried out with Aldrich silica gel (230 mesh). TLC was run with precoated silica gel plate.

The compound to be nitrated (1 mmol) and Montmorillonite KSF (500 mg, Aldrich) were added to a suspension of bismuth nitrate (1 mmol) in THF (10 mL). The solvent was then evaporated

under reduced pressure and dried in the vacuum pump for 5 min (in some cases the mixture needed microwave irradiation for 2 min to complete the reaction). The mixture was then repeatedly washed with dichloromethane (ca 25 mL) and it was concentrated to afford the crude product. The pure product was isolated after column chromatography and by crystallization.

Compounds **6** to **9**, **12** and **13** are known (Aldrich Chemical Company) and these gave identical physico-chemical data with respect to authentic compounds.

The procedure for the nitration of  $\beta$ -lactams is different than the method described above.

Analytical data of **10**, **15** and **16** are reported in earlier publications.<sup>5b,7</sup>

**Cis-1-N-(2-Nitro-4-methoxy)-phenyl-3-acetoxy-4-phenyl-2-azetidinone (18) and Cis-1-N-(3-Nitro-4-methoxy)-phenyl-3-acetoxy-4-phenyl-2-azetidinone (19).** The compound **17** (311 mg, 1 mmol) and Montmorillonite KSF (500 mg, Aldrich) were added to a suspension of bismuth nitrate pentahydrate (485 mg, 1 mmol) in THF (10 mL). The solvent was then evaporated under reduced pressure and irradiated in a kitchen microwave for 2 min (Using 50% power level). The mixture was then repeatedly washed with dichloromethane (ca 25 mL) and it was concentrated to afford a mixture of two regio-isomeric mono nitro derivatives. The pure products were isolated after column chromatography. The compound obtained in the earlier fraction (10% ethyl acetate-90% hexane) was found to be **18** (110 mg, 31%); mp 150 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.48 (1H, d,  $J$  = 8.97 Hz), 7.37–7.33 (6H, m), 7.08 (1H, dd,  $J$  = 2.89 Hz, 8.95 Hz), 5.99 (1H, d,  $J$  = 4.8 Hz), 5.51 (1H, d,  $J$  = 4.9 Hz), 3.82 (3H, s), 1.72 (3H, s); IR (neat) 1756, 1538, 1505, 1222 cm<sup>-1</sup>. Anal. Calcd for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>: C, 60.7%; H, 4.5%; N, 7.9%. Found: C, 60.33; H, 4.56; N, 7.90. The compound obtained in the latter fraction was found to be **19** (225 mg, 67%); mp 152 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.76 (1H, d,  $J$  = 2.65 Hz), 7.56 (1H, dd,  $J$  = 2.68 Hz, 8.96 Hz), 7.37–7.29 (5H, m), 7.01 (1H, d,  $J$  = 9.08 Hz), 5.98 (1H, d,  $J$  = 4.93 Hz), 5.39 (1H, d,  $J$  = 4.93 Hz), 3.90 (3H, s), 1.69 (3H, s); IR (neat) 1756, 1534, 1505, 1222 cm<sup>-1</sup>. Anal. Calcd for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>: C, 60.7%; H, 4.5%; N, 7.9%. Found: C, 60.30; H, 4.45; N, 7.89.

**Trans-1-N-(2-Nitro-4-methoxy)-phenyl-3-acetoxy-4-phenyl-2-azetidinone. (21) and Trans-1-N-(3-Nitro-4-methoxy)-phenyl-3-acetoxy-4-phenyl-2-azetidinone (22).** Following the same procedure as above the compound **20** was converted to a mixture of **21** and **22**. **21** (23%); mp 96 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.44–7.32 (7H, m), 7.06 (1H, dd,  $J$  = 2.9 Hz, 8.9 Hz), 5.54 (1H, d,  $J$  = 2.0 Hz), 5.17 (1H, d,  $J$  = 2.0 Hz), 3.80 (3H, s), 2.19 (3H, s); IR (neat) 1757, 1535, 1505, 1221 cm<sup>-1</sup>. Anal. Calcd for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>: C, 60.7%; H, 4.5%; N, 7.9%. Found: C, 60.40; H, 4.39; N, 7.87. **22** (55%); mp 92 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.72 (1H, d,  $J$  = 2.6 Hz), 7.51 (1H, dd,  $J$  = 2.6 Hz, 9.04 Hz), 7.44–7.32 (5H, m), 7.01 (1H, d,  $J$  = 9.06 Hz), 5.44 (1H, d,  $J$  = 1.54 Hz), 4.99 (1H, d,  $J$  = 1.3), 3.91 (3H, s), 2.21 (3H, s); IR (neat) 1756, 1533, 1505, 1219 cm<sup>-1</sup>. Anal. Calcd for C<sub>18</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>: C, 60.7%; H, 4.5%; N, 7.9%. Found: C, 60.42; H, 4.42; N, 7.84.

## Acknowledgements

We gratefully acknowledge the funding support received for this research project from the

Golden Family Fund for cancer research and NIH Cancer Center Support Grant, 5-P30-CA16672-25, in particular the shared resources of the Pharmacology and Analytic Center Facility.

## References and Notes

1. For example, see: (a) Cornelis, A.; Delaude, L.; Gertmans, A.; Laszlo, P. *Tetrahedron Lett.* **1988**, 29, 5657. (b) Laszlo, P.; Cornelis, A. *Aldrichim. Acta* **1988**, 21, 97. (c) Laszlo, P.; Vandormael, J. *Chem Lett.* **1988**, 1843. (d) Laszlo, P.; Cornelis, A.; Gerstmans, A.; Laszlo, P. *Chem Lett.* **1988**, 1839.
2. Varma, R. S. *Green Chem.* **1999**, 43 and references cited therein.
3. (a) Banik, B. K.; Venkatraman, M. S.; Mukhopadhyay, C.; Becker, F. F. *Tetrahedron Lett.* **1998**, 39, 7243. (b) Banik, B. K.; Ghatak, A.; Venkatraman, M. S.; Becker, F. F. *Synth. Commun.* **2000**, 30, 2701. (c) Banik, B. K. Ghatak, A.; Mukhopadhyay, C.; Becker, F. F. *J. Chem. Res. (S)*, **2000**, 108. (d) Samajdar, S., Becker, F. F., Banik, B. K. *Synth. Commun.* **2001**, in press.
4. Samajdar, S.; Becker, F. F.; Banik, B. K. *Tetrahedron Lett.* **2000**, 41, 8017.
5. (a) Becker, F. F.; Banik, B. K. *Bioorganic & Med. Chem.* **1998**, 50, 2877. (b) Becker, F. F.; Mukhopadhyay, C.; Hackfeld, L.; Banik, I.; Banik, B. K. *Bioorg. & Med. Chem.* **2000**, 8, 2693. (c) Banik, B. K.; Becker, F. F. *Current Medicinal Chemistry* **2001**, in press. (d) Banik, B. K.; Becker, F. F. *Bioorganic & Med. Chem.* **2001**, 9, 593.
6. For nitration with claycop, see: Gigantee, B.; Prazeres, A. O.; Marcelo-Curto, M. J.; Cornelis, A.; Laszlo, P. *J. Org. Chem.* **1995**, 60, 3445 and references cited therein.
7. Cornelis, A.; Laszlo, P.; Pennetreau, P. *J. Org. Chem.* **1983**, 48, 4771.
8. Cornelis, A.; Laszlo, P. *Synthesis* **1985**, 909.
9. Riego, J.; M. Sedin, Z.; Zaldivar, J. M.; Marziano, N. C.; Tortato, C. *Tetrahedron Lett.* **1996**, 37, 513
10. Microwave irradiation has been used in organic synthesis, for some examples, see: (a) Caddick, S. *Tetrahedron* **1995**, 51, 10403. (b) Bose, A. K.; Banik, B. K.; Lavlinskaia, N.; Jayaraman, M.; Manhas, M. S. *Chemtech* **1997**, 27, 18. (c) Mukhopadhyay, C.; Becker, F. F. Banik, B. K. *J. Chem. Res.* **2001**, 28.
11. Suffness, M. *Taxol Science and Applications.*, CRC Press, Boca Raton, 1995.
12. Banik, B. K.; Becker, F. F. *Tetrahedron Lett.* **2000**, 41, 6551.
13. Just, G.; Vgolini, A.; Zamboni, R. *Synth. Commun.* **1979**, 9, 117.
14. We thank Professor A. K. Bose for sharing their results with us.
15. Kropp, J. P.; Breton, G. W.; Fields, J. D.; Tung, J. C.; Loomis, B. R. *J. Am. Chem. Soc.* **2000**, 122, 4280.
16. For some examples of aromatic nitration with mixed acids see: (a) Olah, G. A.; Malhotra, R.; Narang, S. C. In *Nitration. Methods and Mechanisms*, VCH Publishers: New York, 1989. (b) Schofield, K. In *Aromatic Nitration*, Cambridge University Press: Cambridge, 1980.