

Nucleophilic ring opening of 1,2-epoxides in aqueous medium

David Amantini, Francesco Fringuelli*, Oriana Piermatti, Simone Tortoioli, and Luigi Vaccaro

Dipartimento di Chimica, Università di Perugia via Elce di Sotto, 8 06100 Perugia, Italy

E-mail: frifra@unipg.it

Dedicated to Professor Domenico Spinelli on the occasion of his 70th birthday

(received 10 Oct 02; accepted 28 Feb 03; published on the web 08 Mar 03)

Abstract

Nucleophilic ring opening of 1,2-epoxides in aqueous medium in the presence and absence of metal salts is reviewed. Azidolysis, hydrolysis, iodolysis and thiolysis are the reactions mainly investigated. The pH of the reaction medium controls the reactivity and regioselectivity of the process. By working at suitable pH values, even salts such as AlCl₃, SnCl₄ and TiCl₄ are active catalysts.

Keywords: 1,2-Epoxides, azidolysis, hydrolysis, thiolysis, lewis-acid catalysts

Introduction

Sustainable development requires redesigning many organic chemical processes, most of which are often based on technology developed in the first half of the 20th century, and inventing new reactions that use and produce safer chemicals under more environmentally benign conditions.¹ Considerable interest has developed in the use of water in organic synthesis either at “normal” temperature (3-150 °C, 1-5 atm), or elevated temperature (250-350 °C, 40-170 atm) or in supercritical conditions (400 °C, 250-500 atm).² At elevated temperature and pressure the spatial structures of water changes significantly and consequently the physicochemical properties such as dielectric constant, density, solubility parameter and dissociation constant change dramatically.³ Under these severe conditions water can act either as a reagent or as an acid or basic catalyst and many non-polar organic substrates and gases are soluble in the aqueous phase facilitating or making organic reactions possible that were previously thought to occur only in the presence of a strong acid or base or in organic solvent.

Working at elevated temperature and pressure requires high investment costs because the reactions must be carried out in an autoclave (superheated water) or in complex apparatus

(supercritical water) and because special materials are required to overcome problems of corrosion.

The water used in aqueous organic reactions carried out under “normal” conditions of temperature and pressure, acts mainly as a reaction medium; a special, active reaction medium. Liquid water has physiochemical properties that are indeed very different from those of organic solvents: its molecular volume is small, the cohesive pressure is the highest and the internal pressure is the lowest, the surface tension is very large and the heat capacity, heat fusion and heat vaporization are high.⁴ These properties are the consequence of intermolecular forces between closed-shell molecules.

The aqueous medium offers notable advantages with respect to the organic solvent: (i) it is abundant, cheap, non-toxic, non-inflammable, (ii) because of its high heat capacity, it is a heat sink, (iii) the protection-deprotection of functional groups such as $-OH$, $-NH_2$ and $-COOH$ may be unnecessary, (iv) water-soluble compounds can be used directly without derivatization, (v) the reaction products can sometimes be isolated simply by decantation or filtration, (vi) salts, surfactants and cyclodextrins can be used, and (vii) the pH of the reaction medium can be controlled which strongly affects the rate and selectivity of the reaction and allows the reaction to be carried out *one-pot* by domino or consecutive procedures.⁵

Reactions in water often proceed faster than in organic solvents even if one or more reagents and products seem to be insoluble. The use of a co-solvent, to favor the solubility of reagents, does not always favor the reactivity and selectivity of the process.

To date pericyclic, condensation, oxidation and reduction reactions are routinely carried out in aqueous medium.⁶ The use of Lewis-acids such as lanthanide triflates $Bi(OTf)_3$, $Sc(OTf)_3$, $Y(OTf)_3$, $InCl_3$, $InBr_3$, $In(OTf)_3$ has revolutionized the organometallic chemistry.^{6b} Recently we have shown that Lewis acid catalysis in water is strongly dependent on the pH of the aqueous medium and, by maintaining the pH at a suitable level, it is possible to use Lewis acids such as $AlCl_3$, $TiCl_4$ and $SnCl_4$, for which anhydrous conditions are usually recommended, in water.⁷

To date the underlying reasons of the role of the water as reaction medium are still not clear. The inner structure of the liquid water is complex and none of the proposed models completely describes its physiochemical properties. Hydrophobic and hydrogen bonding interactions and polarity are the main factors that influence the reactivity and selectivity of the process.^{4,8}

The 1,2-epoxide functionality is largely present in nature, is biologically important and is a powerful building block in organic synthesis.⁹ Recently Sharpless,¹⁰ following the chemical lead of mother nature, proposed to term “click chemistry” the synthetic approach that generates substances “by joining small units together with heteroatom links (C-X-C)” and defined the criteria that a process must meet to be useful in this context. A “click reaction” that uses this strategy is the nucleophilic ring-opening of 1,2-epoxides. Moreover Sharpless notes that “many of the reactions that meet the click chemistry standard often proceed better (faster and more selectivity) in water than in organic solvents. Nucleophilic additions to epoxide electrophiles are favored by solvents best able to respond continuously to the demanding range of hydrogen-

bonding situations that arise during these processes. In this respect water is unique as reaction medium.”


An overview of nucleophilic ring opening reactions in aqueous media of the 1,2-epoxide functionality present in various compounds, is presented.

Reactions in the absence of metal salts

Azidolysis of 1,2-epoxides is a widely investigated organic reaction because 1,2-azidoalcohols are precursors of vicinal aminoalcohols and are building blocks for carbohydrates and nucleosides.¹¹


The classical protocol uses NaN₃ (5 mol/eq) as reagent in the presence of NH₄Cl (2.3 mol/eq) as coordinating salt in alcohol-water at 70-80 °C. Some examples are illustrated in Table 1¹² and Scheme 1.^{13,14}

Table1. Azidolysis of 1,2-epoxides by sodium azide in aqueous methanol




Entry	Epoxide	t (h)	C-β/C-α	Yield (%)
1		36		92
2		48	73/27	94
3		18	99/1	89
4		16	17/83	85
5		24	95/5	94
6		20	21/79	94
7		12	99/1	95




Under these conditions the reaction is completely *anti*-stereoselective and generally requires a long reaction time. The attack of nucleophile on substituted oxirane ring occurs mainly on the least substituted carbon except when the substituent is an aryl group.




1	t (h)	2/3	Yield (%)
X= CH ₂	18	93/7	95
X= O	60	84/16	96



4	t (h)	5/6	Yield (%)
X= CH ₂	18	99/1	91
X= O	60	99/1	94



Scheme 1. Azidolysis of 1,2-epoxides by sodium azide in aqueous organic media.

**Scheme 2.** Azidolysis of diepoxycyclohexane.

In this case the nucleophilic attack generally occurs mainly on the benzylic carbon (Table 1, entries 4 and 6). In the absence of a specific substituent on the α - and β -carbons of the oxirane ring, the nucleophile preferentially attacks the carbon which is less influenced by unfavorable effects of electron-withdrawing functionalities present in the molecule (Scheme 1).

Table 2. Azidolysis in water of 1,2-epoxides by sodium azide at 30 °C

Entry	Epoxide	pH	t (h)	C- β /C- α	Yield ^a (%)
1		9.5	12		90
		4.2	0.5		90
2		9.5	12	65/35	50
		4.2	0.5	21/79	65
3		9.5	60	65/35	55 ^c
		4.2	48	14/86	70 ^b
4		9.5	20	97/3	77
		4.2	1.5	30/70	58 ^b
5		9.5	24	96/4	88 ^b
		4.2	4	70/30	
6		9.5	28	98/2	90 ^b
		4.2	3	15/85	70
7		9.5	13	3/97	90
		4.2	0.3	3/97	92


^a. Yield of the main isolated product. ^b. With CTABr (1 mol/eq). ^c. Reaction at 100 °C.

Azidolysis of *cis*- and *trans*- 1,4-diepoxyhexanes **7** and **9** (Scheme 2) with hydrazoic acid, generated in situ from NaN₃ and *p*-toluenesulfonic acid, carried out in 1:1 DMSO/H₂O mixture at 70°C, gave the azidoalcohols **8** and **10**, respectively, as sole products which were converted to corresponding aminocyclitols with good yields by catalytic reduction.¹⁵

The reason for using an aqueous-organic medium for the azidolysis of 1,2-epoxides is to carry out the reaction under homogeneous conditions solubilizing both the sodium azide (water) and the epoxide (organic solvent). Recently we have shown^{5c} that the azidolysis of 1,2-epoxides can be suitably performed in water alone under heterogeneous conditions. The nucleophilic addition was totally *anti*-diastereoselective and the reactivity and regioselectivity of the process and the competition of the azido ion with the water or with the hydroxide ion were controlled by working at suitable pH values. Some results are reported in Table 2. The conversion of 1,2-epoxide into azidoalcohol was quantitative. With highly hydrophobic epoxides, the azidolysis was accelerated by carrying out the reaction in the presence of cetyltrimethylammonium bromide (CTABr).


At pH 9.5 the attack of azide ion preferentially occurred, as expected, on the less substituted β-carbon of all epoxides with the exception of styrene oxide (Table 2 entry 7) in which the nucleophile predominately attacked the more substituted benzylic α-carbon. Under acidic conditions (pH 4.2), the reaction was strongly accelerated and a reversed regioselectivity, or an increment of α-attack, was observed for all epoxides except styrene oxide.

Table 3. α/β Cyclization products of 2'-hydroxychalcone epoxides



R ₁	R ₂	12/13 (α/β)
H	H	100:0
6'-OMe	H	14:86
6'-OiPr	H	16:84
4',6'-(OMe) ₂	H	15:85
H	4-Cl	10:90
H	2-CF ₃	0:100

The regioselectivity under acidic conditions is explained by considering that the attack of the azido ion on the more substituted α -carbon arises from the prior protonation of the epoxide, which produces a considerably more positive charge on the tertiary α -carbon than on the secondary or primary one. Coupling these results with the epoxidation of alkenes and the reduction of azides in water,^{16,17} 1,2-azidoalcohols from alkenes and aminoalcohols from epoxides can be prepared by a one-pot procedure in water alone as reaction medium.




Scheme 3. Reactions of benzene oxide-oxepin and naphthalene 1,2-oxide with various nucleophiles.

Nucleophilic attack on a phenyl-substituted epoxide is not always controlled by the benzylic nature of the α -carbon atom. The intramolecular cyclization of 2'-hydroxychalcones **11** is an

example.¹⁸ Under neutral or weakly basic conditions in 1:1 MeCN/H₂O mixture, the unsubstituted chalcone **11** (R₁ = R₂ = H) gave the expected 3-hydroxyflavone **12** (R₁ = R₂ = H), while the *erythro*-aurone-hydrates **13** were the main reaction products when alkoxy substituents were present in the benzene rings in conjugative position (Table 3). The prevalence of β -cyclization was explained on the basis of stereoelectronic factors.

Another example is the azidolysis of styrene oxide carried out in the presence of β -cyclodextrin (β -CD) in water at room temperature.¹⁹ Thus using LiN₃ and 2 mol/eq of β -CD, the reaction conversion after 17 h was 72%, the α/β ratio 50:50 and the *ee* of the adduct coming from the β -attack was 78 % in favor of the (R) enantiomer.

Due to the importance of substituted 1,3-cyclohexadienes in nature and the role of arene oxide-oxepin system in their formation, the ring opening of benzene oxide-oxepin **14** and naphthalene-1,2-oxide **15** were investigated.²⁰ The oxide **14** reacted with various nucleophiles in water under neutral-basic conditions to give 5,6-*trans*-substituted-1,3-cyclohexadienes (Scheme 3).



Scheme 4. Hydrolysis of 1,2-epoxides of polycyclic substrates.

In principle, the products can be obtained by direct attack at C-1 or C-2 of epoxide ring or by conjugative addition resulting from an attack at C-6. Reactions carried out by using benzene-oxepin-3,6-deuterated indicate that the nucleophilic addition occurs only by 1,2-addition.

Reaction of **15** with sodium azide and thioethanol resulted in sole attack at C-2 producing *trans*-1-hydroxy-2-azido- and 1-hydroxy-2-ethylthio-1,2-dihydronaphthalenes (Scheme 3).

The acid-catalyzed hydrolysis of the 7,8-dihydroxy-9,10-epoxide metabolite **16** gives *cis* and *trans* tetrols **18a** and **18b** in 92:8 ratio, whereas its stereoisomer **17**, under the same reaction conditions, gives **19a** and **19b** in 5:95 ratio (Scheme 4). These different ratios of *cis*- and *trans*-tetrols formed from the hydrolysis of **16** and **17** was rationalized by a mechanism involving a favorable axial attack of water on the more stable conformation of the benzylic carbocation.²¹ Recently,²² the rates of hydrolysis of **16** and **17** in water and water-dioxane mixtures have been investigated over a wide range of pH showing that the reaction yields and the mechanism of nucleophilic attack are pH-dependent.

Table 4. Hydrolysis of aryl 1,2-epoxides catalyzed by APSG/I₂

Ar	t (h)	Yield (%)
Ph	1	86
<i>p</i> Cl-C ₆ H ₄	0.75	91
PhOCH ₂	5	90

Iodine (5-10 mol%) supported on aminopropyl silica gel (APSG), prepared by reaction of activated silica gel with aminopropyl triethoxy silane, efficiently catalyzed the hydrolysis of aryl 1,2-epoxides²³ (Table 4).

Table 5. Nucleophilic addition of phenols to (S)-glycidol

Ar	t (h)	Yield (%)
<i>m</i> Pr ^{<i>i</i>} -C ₆ H ₄	14	78
<i>o</i> Pr ^{<i>i</i>} -C ₆ H ₄	12	61
<i>p</i> Cl-C ₆ H ₄	6	86
<i>p</i> MeO-C ₆ H ₄	14	73
<i>p</i> Ph-C ₆ H ₄	14	45
α -Naphthyl	12	62

The nucleophilic additions were carried out in MeCN/H₂O mixture at room temperature and the product obtained from cyclohexene oxide showed a total stereospecific *anti* addition.

Enantiopure 3-aryloxy-1,2-propanediols were prepared by nucleophilic ring-opening of (S)-glycidol (**20**) with phenols in water using catalytic amounts of NaOH (5 mol%).²⁴ The additions were totally regioselective and the optical purity of glycidol was preserved in the products (Table 5). The reactions occurred under heterogeneous conditions and the use of a phase-transfer catalyst did not affect the yield or the reaction times.

Metal Salt Catalyzed Reactions

Metal salt-promoted nucleophilic ring-opening of 1,2-epoxides were investigated in an attempt to improve the regioselectivity and the reaction rate attained by using classical conditions (NaN₃/NH₄Cl, in alcohol or alcohol/water at 70-80 °C). Most of the work was carried out in organic solvent using a variety of metal salts (LiClO₄, LiOTf, Mg(ClO₄)₂, Zn(OTf)₂, NaClO₄, KClO₄, Ti(O-*i*-Pr)₄),^{12,13,25} generally in large excess. Lewis acid-catalyzed reactions carried out in just water have been recently investigated focusing on the azidolysis,^{5d,7a,7b,26a} iodolysis,^{7a,26b} bromolysis,^{26b} and thiolysis^{7c} of 2,3-epoxycarboxylic acids and their esters to develop simple one-pot procedures in aqueous media for the synthesis of hydroxyamino acids, hydroxyacids, and hydroxysulfides.

Table 6. Azidolysis in water of α,β -epoxycyclohexanecarboxylic acid at pH 4.0 catalyzed by various metal salts


Catalyst	t (h)	C ^a (%)	C- β /C- α
none	1.5	10	>99
Yb(OTf) ₃	1.5	32	>99
Sc(OTf) ₃	1.5	30	>99
InCl ₃	1.5	>99	>99
Cu(NO ₃) ₂	1.5	93	>99
AlCl ₃	0.3	>99	>99
Al(NO ₃) ₃	0.3	>99 ^b	>99
AlF ₃	1	98	>99

^a. Reaction conversion; ^b. 4% of diol was present.

The azidolysis reaction of α,β -epoxycyclohexane carboxylic acid (**21**) (Table 6) was efficaciously catalyzed in water at pH 4.0 and 30°C by 1 mol% of $\text{Cu}(\text{NO}_3)_2$, InCl_3 , AlCl_3 , $\text{Al}(\text{NO}_3)_3$ and AlF_3 . These salts, with the exception of $\text{Cu}(\text{NO}_3)_2$, were not operative at pH 7.0. $\text{Yb}(\text{OTf})_3$, and $\text{Sc}(\text{OTf})_3$ were not efficient at either pH 4.0 or at pH 7.0. The reactions were always highly C- β -regioselective.

The catalytic effect of various Lewis acids on the azidolysis of *trans*- α,β -epoxyhexanoic acid (**22**) was also investigated. $\text{Cu}(\text{NO}_3)_2$, AlCl_3 and InCl_3 were again the best catalysts at pH 4.0, while SnCl_4 showed little activity and TiCl_4 slowed the reaction rate; both these salts gave mixtures of C- β and C- α adducts along with large amounts of diols (Table 7).

Table 7. Azidolysis in water of *trans*- α,β -epoxyhexanoic acid at pH 4.0 catalyzed by various salts




Catalyst	t (h)	C ^a (%)	C- β /C- α
none	1.5	28	1
$\text{Cu}(\text{NO}_3)_2$	4	90	16
AlCl_3	1.5	>99	>99
InCl_3	2	>99	>99
TiCl_4	70	75	1.6 ^b
SnCl_4	8	87	3 ^c

^a. Reaction conversion; ^b. 43 % of diol present; ^c. 57 % of diol present.


The stability of **22** at low pH values has allowed the investigation of the nucleophilic ring opening of the oxirane ring at pH 0-2.0. The iodolysis and bromolysis reactions were chosen because the very low concentration of azido ions at these pH values precludes the azidolysis reaction from being carried out under these conditions.

AlCl_3 and InCl_3 efficaciously catalyzed the iodolysis of **22** at pH 1.5 (Table 8) and the iodoalcohol derived from the *anti* attack at β -Carbon was practically the sole reaction product. Under these conditions even TiCl_4 and SnCl_4 were active catalysts but the process was less regioselective.

Table 8. Iodolysis in water of *trans*- α,β -epoxyhexanoic acid at pH 1.5 catalyzed by various metal salts


Catalyst	t (h)	C ^a (%)	C- β /C- α
none	0.5	18	0.6
AlCl ₃	0.5	93	49
TiCl ₄	2	92	2
InCl ₃	0.5	>99	49
SnCl ₄	0.5	92	8

^a. Reaction conversion.

Table 9. InCl₃ effect on the reactivity and selectivity of iodolysis in water of *trans*- α,β -epoxyhexanoic acid.


	W				With InCl ₃ (10 mol%)			
	pH 7.0	pH 4.0	pH 1.5	pH 0.0	pH 7.0	pH 4.0	pH 1.5	pH 0.0
t (h)	24	64	3	0.7	24	2.5	0.5	0.3
C ^a (%)	2	99	86	98	3	99	100	99
diol (%)	0	0	0	0	0	20	0	0
C- β (%)	0	1	36	43	0	78	98	65
C- α (%)	100	99	64	57	100	2	2	35

^a. Reaction conversion.

Table 10. Lewis acid-catalyzed azidolysis in water of α,β -epoxycarboxylic acids at pH 4.0

Epoxide	Catalyst (mol%)	T (°C)	t (h)	C ^a (%)	(C- β)/(C- α)
	none	30	16	25	1
	Cu(NO ₃) ₂ (10)	30	16	>99	>99
	AlCl ₃ (1)	30	3.5	>99	>99
	InCl ₃ (1)	30	10	>99	>99
	none	65	24	82	0.8
	Cu(NO ₃) ₂ (10)	30	18	>99	>99
	AlCl ₃ (1)	65	1.5	>99	>99
	InCl ₃ (1)	65	2	>99	>99
	none	30	3	13	>99
	Cu(NO ₃) ₂ (10)	30	3	>99	>99
	AlCl ₃ (1)	30	0.6	>99	>99
	InCl ₃ (1)	30	1.2	>99	>99
	none	30	8	20	>99
	Cu(NO ₃) ₂ (10)	30	12	>99	>99
	AlCl ₃ (1)	30	0.75	>99	>99
	InCl ₃ (1)	30	8	>99	>99
	none	30	1.5	12	>99
	Cu(NO ₃) ₂ (10)	30	0.25	>99	>99
	InCl ₃ (10)	30	0.02	>99	>99
		none	65	20	>99
AlCl ₃ (1)		65	0.25	99	99
InCl ₃ (1)		65	2	>99	>99
		none	30	0.25	2
	Cu(NO ₃) ₂ (10)	30	0.25	>99	>99
	AlCl ₃ (1)	30	0.25	98	>99
	InCl ₃ (1)	30	1.5	>99	>99

^a. Reaction conversion.

Table 11. InCl₃-catalyzed iodolysis in water of α,β -epoxycarboxylic acids at pH 1.5

Epoxide	Catalyst	T (°C)	t (h)	C ^a (%)	C- β /C- α
	none	0	2	18	0.45
	InCl ₃	0	2	100	49
	none	40	0.5	18	0.55
	InCl ₃	40	0.5	100	49
	none	40	0.5	20	99
	InCl ₃	40	0.5	100	99
	none	20	0.5	8	1.6
	InCl ₃	20	0.5	100	99
	none	0	0.5	12	6.7
	InCl ₃	0	0.5	100	99
	none	0	0.5	12	99
	InCl ₃	0	0.5	100	99
	none	0	9	5	99
	InCl ₃	0	9	100	99


^a. Reaction conversion.





The study of the pH dependence of iodolysis of **22** in the presence and in the absence of InCl₃ gave some interesting results from a synthetic point of view (Table 9).

At pH 4.0 the uncatalyzed reaction was complete in 64h and only the C- α adduct was isolated in quantitative yield. At pH 1.5 the InCl₃-catalyzed reaction afforded exclusively the C- β regioisomer in 0.5h.

All these results have been explained by the fact that the catalytic efficiency of Lewis acid salts in water depends on two factors: (i) the pH of the reaction medium must be below the pK_{1,1} hydrolysis constant of the aqua ion generated from the dissociation of the salt (presence of aqua ion) and (ii) the ability of the aqua ion to give an active complex with the reagents (affinity of the aqua ion). Cu(NO₃)₂, InCl₃, and AlCl₃ generate high concentrations of aqua ions at pH \leq 4.0 and have a great affinity for the α,β -epoxycarboxylic moiety of 1,2-epoxide and the azide and iodine nucleophiles. A revolutionary insight for the organic chemist is that Lewis acids, such as AlCl₃, TiCl₄ and SnCl₄ believed to be unusable as catalysts in organic reactions carried out in water, are efficacious catalysts in aqueous medium provided that they are used at suitable pH values.

Table 12. InBr₃-catalyzed bromolysis in water of α,β -epoxycarboxylic acids at pH 2.0



Epoxide	Catalyst	t (h)	C ^a (%)	C- β /C- α
	none	0.7	18	1.5
	InBr ₃	0.7	99	99
	none	1.3	15	1
	InBr ₃	1.3	99	99
	none	0.5	46	96
	InBr ₃	0.5	99	99
	none	0.3	47	19
	InBr ₃	0.3	99	99


a. Reaction conversion.

These results have lead to the investigation of the azidolysis and iodolysis of a variety of α,β -epoxycarboxylic acids in aqueous media catalyzed by Lewis acids (Tables 10 and 11).

The reactions were highly *regio*- and *diastereo*-selective and pure *anti* or *trans* C- β -adducts were isolated in high yields. Interestingly, the mother liquors from the work up of the reaction containing the metal ion were reused several times without adding fresh catalyst, and without loss of reaction yield and selectivity.


InBr₃ also efficiently catalyzed the bromolysis of α,β -epoxycarboxylic acids^{26b} (Table 12) producing C- β -adducts. However it was difficult to develop a protocol for a wide range of epoxides and sometimes it was difficult to isolate the reaction products so that the reaction yields (32-80%) were not always satisfactory. The reactivity and regioselectivity of thiolysis of alkyl- and aryl-1,2-epoxides with thiophenol in water, as the only solvent, was also strongly affected by

Table 13. Phenylthiolysis of the epoxide of methylenecyclohexane



pH	Catalyst (10 mol%)	t (min)	C- β /C- α	Yield (%)
9.0	none	20	99/1	95
4.0	none	1440	18/82	78
4.0	InCl ₃	5	18/82	78

Table 14. Phenylthiolysis of 2-methyl-2,3-epoxyheptane



pH	Catalyst (10 mol%)	t (min)	C- β /C- α	Yield (%)
9.0	none	2880	99/1	80
4.0	none	1440	37/63	50
4.0	InCl ₃	10	35/65	50

the pH and by the presence of Lewis acids, particularly the InCl_3 .^{7c} The uncatalyzed reaction at pH 9.0 proceeded *via* the $\text{S}_{\text{N}}2$ mechanism, and was generally slow (up to 48h) and occurred prevalently (82-99%) on the less substituted β -carbon of unsymmetrical epoxides. When the substituent was a phenyl ring the nucleophile was driven predominantly to the benzylic α -carbon by electronic effects. At pH 4.0, in the absence of InCl_3 , the reaction remained slow or was even slower, but the α -regioselectivity strongly increased and, in some cases, the C- α phenylsulfide adduct was the prevalent product. Finally the presence of 10% mol of InCl_3 strongly accelerated the reaction that was completed in 3-10 min. Two examples are reported in Table 13 and Table 14.

Coupling the protocol of catalyzed thiolysis of 1,2-epoxides with the preparation of sulfoxides by oxidation of sulfides in water under acidic conditions,²⁷ β -hydroxyphenyl sulfoxides were prepared from epoxides by a one-pot procedure in an aqueous medium. The procedure does not require organic solvent because the sulfoxides precipitate from the aqueous medium and are isolated by simple filtration. These results open the route to the synthesis of building blocks of synthetic interest such as allylic alcohols and ketones by environmentally benign procedures. Chemistry in aqueous media has seriously attracted the attention of organic chemists only during the last ten to fifteen years. Many unexpected results have been obtained already, but many more surprises are to be expected. Perhaps, it is not by chance that nature has chosen water as the reaction medium for its reactions.

Acknowledgments

The Ministero dell'Istruzione dell'Università e della Ricerca (MIUR) / Università degli Studi di Perugia: Progetto Nazionale ("Eterocicli non aromatici in processi stereocontrollati") and Fondazione Cassa di Risparmio di Perugia are thanked for financial support.

References

1. (a) *Green Chemistry: Challenging Perspectives* Eds.: P. Tundo, P.T. Anastas, Oxford Sc. Pub: 1999. (b) Lester T. *Chemistry in Britain* **1996**, 45.
2. (a) Bröll, D.; Kaul, C.; Krämmer, A.; Richter, A.; Jung, M.; Vogel, H.; Zehner, P. *Angew. Chem. Int. Ed.* **1999**, 38, 2999. (b) Siskin, M.; Katritzky, A.R. *Chem. Rev.* **2001**, 101, 825. (c) Katritzky, A. R.; Nichols, D.A.; Siskin, M.; Murugan, R.; Balasubramanian, M. *Chem. Rev.* **2001**, 101, 837.
3. (a) Weingärtner, H.; Franck, E.U.; Wiegand, G.; Dahmen, N.; Schwedt, G.; Frimmel, F.H.; Gordalla, B.C.; Joannsen, K.; Summers, R.S.; Höll, W.; Jekel, M.; Gimbel, R.; Rautenbach, R.; Glaze, W.H. In *Ullmann's Encyclopedia of Industrial Chemistry* Vol. A28, Elvers, B.;

- Hawkins, S. Eds. 5th Edn; VCH: Weinheim 1996. (b) Ohtaki, H.; Radnoi, T.; Yamaguchi, T. *Chem. Soc. Rev.* **1997**, 41.
4. Reichardt, C. *Solvents and Solvent Effects in Organic Chemistry* 2nd Ed; VCH: Weinheim, 1990.
 5. (a) Fringuelli, F.; Germani, R.; Pizzo, F.; Santinelli, F.; Savelli, G. *J. Org. Chem.* **1992**, 57, 1198. (b) Ye, D.; Fringuelli, F.; Piermatti, O.; Pizzo, F. *J. Org. Chem.* **1997**, 62, 3748. (c) Fringuelli, F.; Piermatti, O.; Pizzo, F.; Vaccaro, L. *J. Org. Chem.* **1999**, 64, 6094. (d) Fringuelli, F.; Pizzo, F.; Vaccaro, L. *Synlett* **2000**, 311. (e) Brufola, G.; Fringuelli, F.; Piermatti, O.; Pizzo, F. *Heterocycles* **1996**, 43, 1257; **1997**, 45, 1715.
 6. (a) Li, C.J.; Chang, T.H. *Organic Reactions in Aqueous Media*, Wiley: N.Y., 1997. (b) Grieco, P. A. Ed.; *Organic Synthesis in Water*, Blackie Acad. Professional: London, 1998. (c) Fringuelli, F.; Piermatti, O.; Pizzo, F.; Vaccaro, L. *Eur. J. Org. Chem.* **2001**, 439.
 7. (a) Fringuelli, F.; Pizzo, F.; Vaccaro, L. *J. Org. Chem.* **2001**, 66, 4719. (b) Fringuelli, F.; Pizzo, F.; Vaccaro, L. *Tetrahedron Lett.* **2001**, 42, 1131. (c) Fringuelli, F.; Pizzo, F.; Tortoioli, S.; Vaccaro, L. *Adv. Synth. Catal.* **2002**, 344, 379.
 8. Benson, S.W.; Siebert, E. D. *J. Am. Chem. Soc.* **1992**, 114, 4269.
 9. Fringuelli, F.; Piermatti, O.; Pizzo, F. *Trends in Organic Chemistry* 1997; Vol. 6 Res. Trends Pub. 181.
 10. Kolb, H.C.; Finn, M.G.; Sharpless, K.B. *Angew. Chem., Int. Ed.* **2001**, 40, 2005.
 11. (a) Coe, D.M.; Myers, P.L.; Parry, D.M.; Roberts, S.M.; Storer, R. *J. Chem. Soc., Chem. Comm.* **1990**, 151; (b) Jacobs, G.A.; Tino, J.A.; Zahler, R. *Tetrahedron Lett.* **1989**, 30, 6955.
 12. Chini, M.; Crotti, P.; Macchia, F. *Tetrahedron Lett.* **1990**, 31, 5641.
 13. Crotti, P.; Di Bussolo, V.; Favero, L.; Macchia, F.; Pineschi, M. *Eur. J. Org. Chem.* **1998**, 1675.
 14. Banwell, M.G.; Haddad, N.; Hudlicky, T.; Nugent, T.C.; Mackay, M.F.; Richards, S.L. *J. Chem. Soc., Perkin Trans. 1* **1997**, 1779.
 15. Kavadias, G.; Droghini, R. *Can. J. Chem.* **1979**, 57, 1870.
 16. Fringuelli, F.; Germani, R.; Pizzo, F.; Savelli, G. *Tetrahedron Lett.* **1989**, 30, 1427.
 17. Fringuelli, F.; Pizzo, F.; Vaccaro, L. *Synthesis* **2000**, 646.
 18. Adams, C.J.; Main, L. *Tetrahedron* **1991**, 47, 4979.
 19. Guy, A.; Doussot, J.; Garreau, R.; Godefroy-Falguieres, A. *Tetrahedron: Asymmetry* **1992**, 3, 247.
 20. (a) De Marinis, R.M.; Berchtold, G.A. *J. Am. Chem. Soc.* **1969**, 91, 6525. (b) Jeffrey, A.M.; Yeh, H.J.C.; Jerina, D.M.; De Marinis, R.M.; Foster, C.H.; Piccolo, D.E.; Berchtold, G.A. *J. Am. Chem. Soc.* **1974**, 96, 6929. (c) Mc Manus, M.J.; Berchtold, G.A. *J. Am. Chem. Soc.* **1985**, 107, 2977.
 21. (a) Whalen, D.L.; Montemarano, J.A.; Thakker, D.R.; Yagi, H.; Jerina, D.M. *J. Am. Chem. Soc.* **1997**, 99, 5522.
 22. Doan, L.; Lin, B.; Yagi, H.; Jerina, D.M.; Whalen, D.L. *J. Am. Chem. Soc.* **2001**, 123, 6785.
 23. Tamami, B.; Iranpoor, N.; Mahdavi, H. *Synth. Commun.* **2002**, 32, 1251.

24. Palermo, S.; Waykole, L.; Chen, K.-M.; Prashad, M.; Prasad, K.; Repic, O.; Blacklock, T.J. *Synth. Commun.* **1997**, *27*, 1757.
25. (a) Chong, J.M.; Sharpless, K.B. *J. Org. Chem.* **1985**, *50*, 1560. (b) Thijs, L.; Porskamp, J.J.M.; van Loon, A.A.W.M.; Derks, M.P.W.; Feenstra, R.W.; Legters J.; Zwanenburg, B. *Tetrahedron* **1990**, *46*, 2611. (c) Saito, S.; Takahashi, N.; Ishikawa, T.; Morikawa, T. *Tetrahedron Lett.* **1991**, *32*, 667. (d) Legters, J.; Thijs, L.; Zwanenburg, B. *Recl. Trav. Chim. Pays-Bas* **1992**, *111*, 1. (e) Azzena, F.; Crotti, P.; Favero, L.; Pineschi, M. *Tetrahedron* **1995**, *51*, 13409.
26. (a) Fringuelli, F.; Pizzo, F.; Vaccaro, L. *J. Org. Chem.* **2001**, *66*, 3554. (b) Amantini, D., Fringuelli, F.; Pizzo, F.; Vaccaro, L. *J. Org. Chem.* **2001**, *66*, 4463.
27. Fringuelli, F.; Pellegrino, R.; Pizzo, F. *Synth. Commun.* **1993**, *23*, 3157.