

Supplementary Material

Asymmetric aziridination of *N*-*tert*-butanesulfinyl imines with phenyldiazomethane via sulfur ylides


Zheng Xue, Veronica M. Dee, Louisa J. Hope-Weeks, Bruce R. Whittlesey,
and Michael F. Mayer*

*Department of Chemistry and Biochemistry, Texas Tech University, Lubbock, TX 79409-1061,
USA*



E-mail address: mf.mayer@ttu.edu

Table of Contents


1. ¹H NMR and ¹³C NMR spectra of (*R*_s,2*R*,3*R*)-**3a** and (*R*_s,2*S*,3*S*)-**3a**
at room temperature and –20 °C with expanded inserts S2-S4
2. ¹H NMR and ¹³C NMR spectra of compounds **3b-i** at room temperature
and –20 °C S5-S20





¹H NMR (top) and ¹³C NMR (bottom) of **(*R_S,2*R*,3*R*)*-3a** at 23 °C




¹H NMR (top) and ¹³C NMR (bottom) spectra of (*R_S*,2*S*,3*S*)-**3a** at 23 °C












^1H NMR (top) and ^{13}C NMR (bottom) of $(R_S,2S,3S)$ -**3d** at 23 °C





^1H NMR (top) and ^{13}C NMR (bottom) of $(R_S,2S,3S)$ -**3e** at 23 °C







^1H NMR (top) and ^{13}C NMR (bottom) of $(R_S,2R,3R)$ -**3f** at 23 °C










^1H NMR (top) and ^{13}C NMR (bottom) of $(R_S,2S,3R)$ -**3h** at 23 °C



^1H NMR (top) and ^{13}C NMR (bottom) of $(R_S,2R,3S)$ -**3h** at 23 °C







^1H NMR (top) and ^{13}C NMR (bottom) of 2,3-cis-3-2k at 23 °C