

Quinidine catalyzed reaction between 4-formyloxyazetidin-2-one and some thiophenols, thiols and alcohols

Anna Koziol,^a Elisa Altieri,^b Bartłomiej Furman,^a Jolanta Solecka,^c and Marek Chmielewski^{a*}

^a*Institute of Organic Chemistry, Polish Academy of Sciences, Kasprzaka 44/52, 01-224 Warsaw, Poland*

^b*Dip. di Chimica Organica e Biologica, Università, Vill. S. Agata, I 98166 Messina, Italy*

^c*National Institute of Public Health-National Institute of Hygiene, Chocimska 24, 00-791 Warsaw, Poland*

E-mail : chmiel@icho.edu.pl

Dedicated to Professor Siegfried Blechert in occasion of his 65th birthday

DOI: <http://dx.doi.org/10.3998/ark.5550190.0012.405>

Abstract

Readily available 4-formyloxyazetidinone was enantioselectively transformed into 4-substituted azetidinones upon treatment with 0.1 equiv. of the cinchona alkaloid (quinidine) in toluene *via* intermolecular nucleophilic trapping of the *N*-acylimine intermediate by mercapto-, or hydroxy-moieties of thiophenols, thiols, phenols and alcohols. Additionally, biological activity tests were performed on the newly synthesized β -lactams.

Keywords: Chiral Lewis base, chiral Lewis acid, enantioselective synthesis, β -lactams

Introduction


Compounds containing a β -lactam core are synthetic targets of great importance for academic and industrial laboratories owing to their various biological activities such as antibacterial,¹ inhibition of enzymes like HLE,² thrombin,³ human cytomegalovirus protease,⁴ or highly specific cholesterol inhibitors.⁵ The synthesis of β -lactam compounds can be achieved in two ways: by transformation of readily available natural products, or by total synthesis.⁶ In the latter case, the method of formation of the β -lactam ring together with control of the stereogenic centers at C-3 and C-4 of the azetidin-2-one ring is crucial for the success of the synthesis.

Recently, we have demonstrated that a 4-acyloxy substituent in the azetidin-2-one ring could be replaced by phenols in both chiral base and chiral acid catalyzed reactions leading to enantiomerically enhanced products. In particular, quinidine effectively catalysed the reaction between **1** and salicyl aldehydes **2** providing the corresponding 3,4-benzo-2-hydroxy-5-oxacephams **5** (Scheme 1).⁷ The key step of this cascade reaction is based on the chiral Lewis base (quinidine) mediated, enantioselective intermolecular alkylation of the phenol hydroxy group with neutral 1,4-dehydroazetidin-2-one **3** which is followed by formation of the cyclic hemiacetal **5**.⁸ Other phenols react with **1** with similar enantioselectivity providing the corresponding 4-aryloxyazetidinones.

This methodology enables the formation of a new stereogenic centre at C-4 of the azetidin-2-one ring with an enantioselectivity of up to 50% *ee* and a good yield 70% - 90%. The absolute configuration of **5** was assigned on the basis of the CD spectra and depends on the cinchona alkaloid used.⁷

We have also described a chiral Lewis acid mediated intramolecular alkylation of the phenol hydroxy group by the acyliminium ion **8** generated from the corresponding 4-formyloxy-*N*-substituted-azetidinones **6**.⁹ This reaction, however, always requires an equimolar amount of the acid promoter **7** and proceeds in a low yield but high asymmetric induction as a kinetic resolution of the primary formed racemic 3,4-benzo-5-oxacepham **9-rac**. Contrary to the base catalyzed reactions, Lewis acid promoted intermolecular reactions between **1** and phenols proceed in low yield and low asymmetric induction.

So far, we have reported asymmetric reactions concerned with the formation of 3,4-benzo-5-oxacepham skeletons or 4-aryloxyazetidinones. Examination of whether other nucleophiles e.g thiophenols, thiols, or alcohols, can be applied in similar asymmetric processes is a logical consequence of our previous investigations. Formation of enantiomerically enriched 4-substituted azetidinones would open an access to interesting monocyclic β -lactams, or after proper transformations, to a variety of clavams, penams and cephams.




Scheme 1. Synthetic strategy.

Results and Discussion

It was of interest to check whether cinchona alkaloids would catalyze the nucleophilic substitution at C-4 of the azetidin-2-one ring involving, more nucleophilic than oxygen congeners, thiophenols. 4-Formyloxyazetidin-2-one **1** was prepared by the ozonolysis of 4-vinyloxyazetidin-2-one at $-78\text{ }^{\circ}\text{C}$ in CH_2Cl_2 , followed by reductive workup with dimethyl sulfide. High yields, and reproducibility of ozonolysis was achieved by the inclusion of a small amount of ozonizable dye (Sudan red 7B) as an internal standard, which indicated the reaction end point.¹⁰ The results of the experiments performed under conditions found optimal for the corresponding phenols are shown in (Table 1).

Table 1. Quinidine-catalyzed nucleophilic substitution with thiols **10**



Entry	Thiol	R	Product	Yield [%] ^a	<i>ee</i> [%] ^b
1	10a	H	11a	84	42
2	10b	2-Me	11b	89	49
3	10c	4-OH	11c	95	11
4	10d	4-OMe	11d	75	34
5	10e	4-Br	11e	95	34

^aIsolated yield determined after flash chromatography on SiO_2 . Reaction was carried out until disappearance of substrate **1** (TLC monitored).


^bThe enantiomeric excess was determined by chiral HPLC.

After a few initial attempts we found that the reaction of 4-formyloxyazetidin-2-one **1** with the thiophenols **10a-e**, was most effective in the presence of a catalytic amount (0.1 mol. equiv.) of quinidine **3**, in toluene at room temperature. As shown in Table 1, the enantiomeric excess, as well as the product's yield depended on the nature of the substituent in the thiophenol ring. In all cases, however, good chemical yields and moderate enantioselectivities of 4-aryltioazetidin-2-ones **11a-e** were observed.

The scope of nucleophiles was then extended to thiosalicylaldehyde **12**. Under the same conditions, azetidinone **1** underwent condensation with **12** to provide 3,4-benzo-2-


hydroxycepham **13** in 85% yield, with 38% *ee* (Scheme 2). The selectivity in the formation of **13** was similar to that found for reaction of **2** with salicylaldehyde.⁷

The experiments on cinchonine catalyzed substitutions testify that the switch from the oxygen nucleophile to the sulfur analog does not affect the enantioselectivity of the process. This might suggest that the elimination-addition mechanism of the first step is the same for both nucleophiles. In the case of thiosalicylaldehyde **12**, the second step leading to the ring closure is under thermodynamic control and consequently leads to the *exo* location of the hemiacetal hydroxyl group.



Scheme 2. Quinidine-catalyzed nucleophilic substitution with *o*-thiosalicylaldehyde **12**.

Subsequently we decided to examine whether aliphatic alcohols and thiols would be proper substrates in the same nucleophilic substitution processes. First experiments showed that the products obtained **16/17** were volatile and therefore the yield of the reaction and the proportions of the enantiomers by HPLC were hard to be correctly assign. We found that in most cases direct benzylation of the nitrogen atom in **16/17** leading to **18/19** allowed us to determine the yield and enantiomeric excess of the reaction (Scheme 3). Independent experiments performed on 4-benzyloxyazetidin-2-one **16c** showed that *N*-benzylation is almost a quantitative process which does not affect the enantioselectivity of the nucleophilic substitution.




Scheme 3. Synthesis of *N*-benzylated 4-substituted azetidin-2-ones **18** and **19**.

As has been demonstrated recently, the reaction between phenols and 4-formyloxyazetidin-2-one **1** in the presence of a catalytic amount of quinidine proceeds in good yield and moderate enantioselectivity to afford the corresponding 4-aryloxyazetidin-2-ones.⁷ Different results were noticed when aliphatic alcohols or mercaptans were applied in the same reaction (Table 2, 3). In the case of aliphatic alcohols **14** an equimolar amount of quinidine was required to obtain full

conversion of **1**. An excess of the nucleophile **14** should also be raised from 1.5 to 3 equivalents (Table 2). The results obtained are modest and show a good yield only for benzyl alcohol, whereas the enantioselectivities are low for all nucleophiles.

Table 2. Quinidine promoted nucleophilic substitution with alcohols **14a-f**




Entry	Substrate	R	Product	Yield[%] ^a	ee [%] ^b
1	14a	CH ₂ =CHCH	18a	15	10
2	14b	CH≡CCH ₂	18b	30	12
3	14c	Ph CH ₂	18c	77	4
4	14d	Et	18d	12	nd
5	14e	<i>i</i> -Pr	18e	15	11
6	14f	<i>t</i> -Bu	18f	0	-

^aIsolated yield determined after flash chromatography on SiO₂. Reaction was carried out until the disappearance of the substrate **1** (TLC monitored).

^bThe enantiomeric excess was determined by chiral HPLC.

In contrast to aliphatic alcohols **14**, analogous thiols **15** undergo reaction with **1** to provide the corresponding sulfides **19** under standard catalytic reaction conditions in good chemical yields and enantioselectivities similar to that observed for phenols (Table 3).

Comparison of the corresponding results of alcohols and thiols shows much better results for the latter. When *t*-butyl alcohol was used no reaction was observed (Table 2, entry 6), but its thio- congener gave an expected product in 65% yield and 12% *ee* (Table 3, entry 3). The best result was obtained for triphenylmethyl thiol (Table 3, entry 4). It gave an excellent yield of **20** and 51% *ee* (Table 3, entry 4). In this case, owing to high molecular weight, further benzylation of the nitrogen atom in order to assign enantioselectivity was not necessary.

Table 3. Quinidine-catalyzed nucleophilic substitution with thioalcohols **15 a-d**


Entry	Substrate	R	Product	Yield[%] ^a	ee [%] ^b
1	15a	CH ₂ =CHCH	19a	40	38
2	15b	PhCH ₂	19b	86	34
3	15c	<i>t</i> -Bu	19c	65	12
4	15d	Ph ₃ C	20	89	51

^aIsolated yield determined after flash chromatography on SiO₂. Reaction was carried out until the disappearance of the substrate **1** (TLC monitored).

^bThe enantiomeric excess was determined by chiral HPLC.

In contrast to chiral base catalyzed nucleophilic substitution at C-4 of azetidinone **1**, the intermolecular reaction promoted by chiral Lewis acid – equimolar mixture of SnCl₄ and (*S*)- α -di-naphthyl-BINOL **7** – proceeded in a different way for phenols and thiophenols.⁹ We found that the direct reaction between **1** and thiophenol **10a** in the presence of the BINOL/SnCl₄ complex **7** provided the racemic compound **11a** in low yield. Asymmetric degradation of the racemic 4-phenylthioazetidin-2-ones **11** gave better enantiomer selectivities. After 6h **11a** underwent destruction in 41% yield and 42% *ee* (Table 4, entry 1). In comparison to phenols, nucleophilic substitution involving thiophenols is faster, whereas asymmetric degradation of the primary formed racemic compound is slower due to a more stable *N,S*-acetal than *N,O*-congener. The moderate asymmetric induction for the Lewis acid promoted kinetic resolution process may be also caused by a reversible process abstraction-addition of the thiophenoxy group which should lead to partial racemization of the product.

Chiral Lewis acid **7** catalyzed, asymmetric degradation of the racemic 3,4-benzocepham **22** after 6 h at 0 °C provided an enantiomerically enhanced compound in 48% yield and 46% *ee* (Scheme 4). The lower result of the kinetic resolution of **22** in comparison with **9** is caused by the same reasons as suggested for **11** and **21**.


Table 4. Chiral Lewis acid **7** promoted deracemization of 4-thioaryloxyazetid-2-ones

Entry	Lewis acid 7	R	R ¹	Product	Yield [%] ^a	ee [%] ^b
1	<i>R</i>	H	H	11a	41	42 (<i>S</i>)
2	<i>R</i>	H	4-OMe	11d	45	34 (<i>S</i>)
3	<i>S</i>	Bn	H	21a	40	37 (<i>R</i>)
4	<i>S</i>	Bn	4-OMe	21b	43	21 (<i>R</i>)

^aIsolated yield determined after flash chromatography on SiO₂.

^bThe enantiomeric excess was determined by chiral HPLC.

Similar asymmetric degradation was performed for 5-oxacepham **23** obtained by us a few years ago.¹¹ Under the same conditions ee was found to be only 22% (Scheme 4).

**Scheme 4.** Chiral Lewis acid **7** promoted deracemization of cepham's congeners.

Biological activity

4-Thiophenoxy compounds **11a-e** were tested for their biological activity. An inhibition of the DD-carboxypeptidase activity and, separately, an inhibition of β -lactamase were measured.¹²⁻¹³

Within the studied series, all of the tested azetidin-2-ones **11a-e** showed a low activity towards DD-peptidase. Except **11b** which showed the inhibition of β -lactamase class A activity, expressed by the value of 2.5×10^{-2} , other compounds do not showed any activity as inhibitors of the β -lactamase.

Conclusions

In conclusion, we report a simple approach to the asymmetric synthesis of 3,4-benzo-2-hydroxycephams, 4-thiophenoxyazetidin-2-ones and 4-alkoxyazetidin-2-ones. The key step of this method is based on the chiral Lewis base mediated, enantioselective, intermolecular alkylation of mercapto or hydroxyl groups by intermediately formed dehydro-azetidinone. It is important to note that the chiral Lewis base is used in catalytic amounts without effecting both the yield and enantioselectivity. The reactions proceed in similar yield and with similar selectivity for thiols and alcohols. This is in contrast with that previously described for the chiral Lewis acid mediated enantioselective alkylation of the phenol hydroxy group by the *N*-acyliminium ion generated from the 4-formyloxyazetidinone **1**, which requires always an equimolar amount of the promoter and proceeds in low yield but high asymmetric induction as the intramolecular process, or in good yield but low asymmetric induction as the intermolecular process. Due to the higher stability of the *N,S*-acetal versus the *N,O*-one asymmetric degradations of 4-alkoxyazetidinones or 5-oxacephams proceed more readily.

Experimental Section

General. All reactions involving air- and moisture-sensitive materials were performed under an atmosphere of dry argon using dry solvents. Reagents were purchased from commercial suppliers and used without further purification, unless noted. Tetrahydrofuran was distilled from Na and bezophenon ketyl and dichloromethane were distilled from CaH_2 . All processes were performed until the disappearance of the starting material (TLC monitoring). Melting points were determined using a hot-stage apparatus with a microscope and are uncorrected. The column chromatography was performed on Kiesel gel (230 – 400 mesh). Thin layer chromatography (TLC) was performed on aluminum sheets Silica Gel 60 F₂₅₄ (20 x 20 x 0.2). TLC spots were visualized in UV (254 nm) and by treatment by the solution of ceric acid. Routine NMR spectra were obtained at 500 MHz for ^1H NMR and 128 MHz for ^{13}C NMR, using CDCl_3 as a solvent and TMS as an integral reference (δ 0 for ^1H and δ 0 ^{13}C). IR data were obtained with a FT-IR-1600-Perkin – Elmer spectrophotometer. High resolution mass spectra were recorded on an AMD 604 mass spectrometer (EI, 70 eV) and ESI-TOF Mariner spectrometer (Perspective Biosystem). The optical rotations were measured using JASCO J-1020 digital polarimeters. The high pressure liquid chromatography (HPLC) was performed using a

Merck – Hitachi chromatograph with a L-2130 pump, diode array detector L-2450, analytical columns : Chiralpark® AD-H, Chiralcel® OD-H and OJ-H and hexane / *i*-propanole as solvents.

Quinidine-catalyzed nucleophilic substitution with thiols (10) and (12)

To a solution of 4-formyloxazetididin-2-one **1** (30 mg, 0.26 mmol, 1.0 equiv.) and thiol **10**, **12** (0.29 mmol, 1.1 equiv.) in toluene (5 mL) at room temperature was added quinidine (9 mg, 0.1 equiv.) under an argon atmosphere. The reaction mixture was stirred for 48 h till the disappearance of the starting material and diluted with water (5 mL). The aqueous phase was extracted with ether (3x10 mL). The organic extracts were combined and dried over Na₂SO₄. The solution was filtered and the filtrate evaporated. The crude product was purified by column chromatography to yield **11**, **13** respectively.

(4S)-4-Phenylthioazetididin-2-one (11a). White solid, yield 84%, 42% *ee*, 39 mg (0.22 mmol), m.p. 54-56°C, IR (ν_{\max} , cm⁻¹): 1778 (C=O), $[\alpha]_{\text{D}}^{26} = -110.7$ (c 0.73, CHCl₃). ¹H NMR (500 MHz, CDCl₃) δ_{H} 2.89 (ddd, $J = 15.2, 2.3, 1.9$ Hz, 1H, CH₂), 3.36 (ddd, $J = 15.2, 4.9, 1.9$ Hz, 1H, CH₂), 5.00 (dd, $J = 4.9, 2.3$ Hz, 1H, CH-S), 6.38 (s, 1H, NH), 7.35 (m, 3H, Ar), 7.45 (m, 2H, Ar). ¹³C NMR (125 MHz, CDCl₃) δ_{C} 45.4, 54.2, 128.6, 129.4, 131.4, 133.5, 165.8. HR MS (TOF MS EI+) Calcd for M⁺ C₉H₉NOS 179.0405, Found: 179.0396. HPLC [OD-H, hexane : *i*-propanole = 8 : 2, 0.5 mL/min, t_{R} [R]= 19.9 (minor), t_{R} [S]= 23.8 (major)].

(4S)-4-(*o*-Tolylthio)-azetididin-2-one (11b). White solid, yield 89%, 49% *ee*, 44 mg (0.23 mmol), m.p. 62-64°C, IR (ν_{\max} , cm⁻¹): 1776 (C=O), $[\alpha]_{\text{D}}^{26} = -15.8$ (c 1.0, CHCl₃). ¹H NMR (500 MHz, CDCl₃) δ_{H} 2.44 (s, 3H, CH₃), 2.93 (ddd, $J = 15.2, 2.3, 1.9$ Hz, 1H, CH₂), 3.45 (ddd, $J = 15.2, 4.9, 1.9$ Hz, 1H, CH₂), 4.99 (dd, $J = 4.9, 2.3$ Hz, 1H, CH-S), 6.25 (s, 1H, NH), 7.19 (m, 1H, Ar), 7.25 (m $J = 2\text{H}$, Ar), 7.37 (m, 1H, Ar). ¹³C NMR (125 MHz, CDCl₃) δ_{C} 20.9, 45.6, 53.7, 126.8, 128.5, 130.8, 131.4, 132.9, 140.5, 165.8. HR MS (TOF MS E+) Calcd for M⁺ C₁₀H₁₁NOS 193.0561, Found: 193.0569. HPLC [OD-H, hexane : *i*-propanole = 8 : 2, 0.5 mL/min, t_{R} [R]= 17.0 (minor), t_{R} [S]= 22.4 (major)].

(4S)-4-(4'-Hydroxyphenylthio)-azetididin-2-one (11c). White solid, yield 95%, 11% *ee*, 48 mg (0.25 mmol), m.p. 110-112°C, IR (ν_{\max} , cm⁻¹): 1788 (C=O), $[\alpha]_{\text{D}}^{26} = -1.8$ (c 0.50, MeOH). ¹H NMR (500 MHz, CD₃OD) δ_{H} 2.69 (d, $J = 15.2, 2.1$ Hz, 1H, CH₂), 3.22 (dd, $J = 15.2, 4.8$ Hz, 1H, CH₂), 4.87 (dd, $J = 4.8, 2.1$ Hz, 1H, CH-S), 6.80 (d, $J = 8.8$ Hz, 2H, Ar), 7.37 (d, $J = 8.8$ Hz, 2H, Ar). ¹³C NMR (CD₃OD) δ_{C} 44.8, 55.5, 117.2, 120.3, 138.3, 159.9, 169.2. HR MS (ESI) Calcd for [M + Na]⁺ C₉H₈NO₂NaS 218.0246, Found: 218.0242. HPLC [OD-H, hexane : *i*-propanole = 8 : 2, 0.5 mL/min, t_{R} [R]= 21.3 (minor), t_{R} [S]= 23.6 (major)].

(4S)-4-(*p*-Methoxyphenylthio)-azetididin-2-one (11d). White solid, yield 75%, 34% *ee*, 40 mg (0.19 mmol), m.p. 71-73°C, IR (ν_{\max} , cm⁻¹): 1776 (C=O), $[\alpha]_{\text{D}}^{26} = -24.8$ (c 0.755, CHCl₃). ¹H NMR (500 MHz, CDCl₃) δ_{H} 2.82 (ddd, $J = 15.2, 2.2, 1.6$ Hz, 1H, CH₂), 3.29 (ddd, $J = 15.2, 4.9, 1.6$ Hz, 1H, CH₂), 3.81 (s, 3H, CH₃), 4.88 (dd, $J = 4.9, 2.2$ Hz, 1H, CH-S), 6.11 (s, 1H, NH), 6.89 (d, $J = 8.8$ Hz, 2H, Ar), 7.43 (d, $J = 8.8$ Hz, 2H, Ar). ¹³C NMR (CDCl₃) δ_{C} 44.9, 54.7, 55.4, 114.9, 120.7, 136.7, 160.7, 165.7. HR MS (ESI) Calcd for [M + Na]⁺ C₁₀H₁₁NO₂NaS 232.0402,

Found: 232.0398. HPLC [OD-H, hexane : *i*-propanole = 8 : 2, 0.5 mL/min, t_R [R]= 21.2 (minor), t_R [S]= 25.4 (major)].

(4S)-4-(4'-Bromophenylthio)-azetid-2-one (11e). White solid, yield 95%, 34% *ee*, 66 mg (0.25 mmol), m.p. 107-109°C, IR (ν_{\max} , cm^{-1}): 1779 (C=O), $[\alpha]_{\text{D}}^{26} = -32.9$ (c 0.96, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.88 (dm, $J = 15.12$, 1H, CH_2), 3.39 (ddd, $J = 15.2$, 4.9, 2.0 Hz, 1H, CH_2), 4.98 (dd, $J = 4.9$, 2.3 Hz, 1H, CH-S), 6.29 (s, 1H, NH), 7.32 (d, $J = 8.5$ Hz, 2H, Ar), 7.49 (d, $J = 8.5$ Hz, 2H, Ar). ^{13}C NMR (125 MHz, CDCl_3) δ_{C} 45.5, 54.2, 123.3, 132.6, 135.0, 165.5. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+$ $\text{C}_9\text{H}_8\text{NONaSBr}$ 279.9402, Found: 279.9396. HPLC [OD-H hexane : *i*-propanole = 8 : 2, 0.5 mL/min, t_R [R]= 20.3 (minor), t_R [S]= 21.9 (major)].

(2R, 6S)-3,4-Benzocepham (13). White solid, yield 85%, 38% *ee*, 46 mg (0.22 mmol), m.p. 130-132°C, IR (ν_{\max} , cm^{-1}): 1778 (C=O), $[\alpha]_{\text{D}}^{26} = -25.8$ (c 1.0, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 3.00 (d, $J = 15.1$ Hz, 1H, CH_2), 3.351 (dd, $J = 15.1$, 4.3 Hz, 1H, CH_2), 4.58 (s, 1H, OH), 4.93 (d, $J = 4.3$ Hz, 1H, CH-S), 6.04 (s, 1H, CH), 7.26 (m, 3H, Ar), 7.64 (d, $J = 5.4$ Hz, 1H, Ar). ^{13}C NMR (125 MHz, CDCl_3) δ_{C} 45.8, 45.9, 69.7, 126.2, 128.6, 129.3, 129.5, 129.9, 130.1, 165.7. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+$ $\text{C}_{10}\text{H}_9\text{NO}_2\text{NaS}$ 230.0246, Found: 230.0257. HPLC [OD-H, hexane : *i*-propanole = 8 : 2, 0.5 mL/min, t_R [6R]= 16.8 (minor), t_R [6S]= 21.3 (major)].

Quinidine-catalyzed nucleophilic substitution with alcohols (14)

To the solution of 4-formyloxyazetid-2-one **1** (30 mg, 0.26 mmol, 1.0 equiv.) and alcohol **14** (0.78 mmol, 3.0 equiv.) in toluene (5 mL) at room temperature was added quinidine (90 mg, 1.0 equiv.) under argon atmosphere. The reaction mixture was stirred for 48 h till disappearance of starting material and filtered through small layer of silica gel. The silica layer was washed with toluene (2 mL). Subsequently the filtrate containing 4-alkoxyazetid-2-one **16** was slowly added to solution of 50% aqueous NaOH (5 mL), and Bu_4NBr (84 mg, 0.26 mmol, 1.0 equiv.) and benzyl chloride (135 μL , 1.17 mmol, 4.5 equiv.) in 1 mL of toluene. The reaction mixture was stirred for 30 min till disappearance of starting material and diluted with water (10 mL). The aqueous phase was extracted with ethyl acetate (3x15 mL). The organic extracts were combined and dried over MgSO_4 . The solution was filtered and filtrate evaporated. The crude product was purified by column chromatography to yield **18**.

(4S)-N-Benzyl-4-allyloxyazetid-2-one (18a). Yellow oil, yield 15%, 9 mg (0.04 mmol), 10% *ee*, IR (ν_{\max} , cm^{-1}): 1759 (C=O), $[\alpha]_{\text{D}}^{26} = -8.3$ (c 0.5, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.89 (dm, $J = 14.9$ Hz, 1H, CH_2), 3.05 (dd, $J = 14.9$, 3.8 Hz, 1H, CH_2), 3.89 (ddt, $J = 12.7$, 5.7, 1.5 Hz, 1H, $\text{CH}_2\text{-O}$), 3.95 (ddt, $J = 12.7$, 5.7, 1.5 Hz, 1H, $\text{CH}_2\text{-O}$), 4.21 (d, $J = 15.2$ Hz, 1H, CH-N), 4.60 (d, $J = 15.2$ Hz, 1H, $\text{CH}_2\text{-N}$), 4.91 (dd, $J = 6.1$, 1.5 Hz, 1H, $\text{CH}_2=$), 5.15 (dm, $J = 9.7$ Hz, 1H, $\text{CH}_2=$), 5.20 (dd, $J = 3.8$, 1.7 Hz, 1H, CH-O), 5.80 (m, 1H, CH=), 7.30 (m, 3H, Ar), 7.35 (m, 2H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 44.4, 44.7, 68.5, 80.7, 117.6, 127.8, 128.3, 128.8, 133.5, 135.7, 165.9. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+$ $\text{C}_{13}\text{H}_{15}\text{NO}_2\text{Na}$ 240.0995, Found: 240.0987. HPLC [AD-H, hexane : *i*-propanole = 95:5, 0.5 mL/min, t_R [R]= 27.9 (minor), t_R [S]= 30.1 (major)].

(4S)-N-Benzyl-4-propargyloxyazetididin-2-one (18b). Yellow oil, yield 30%, 12% *ee*, 17 mg (0.08 mmol), IR (ν_{\max} , cm^{-1}): 1762 (C=O), $[\alpha]_{\text{D}}^{26} = -11.5$ (c 0.7, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.44 (t, $J = 2.4$ Hz, 1H, CH), 2.97 (dd, $J = 14.9, 1.2$ Hz, 1H, CH_2), 3.09 (dd, $J = 14.9, 3.7$ Hz, 1H, CH_2), 4.05 (dd, $J = 16.0, 2.4$ Hz, 1H, $\text{CH}_2\text{-C}$), 4.11 (dd, $J = 16.0, 2.4$ Hz, 1H, $\text{CH}_2\text{-C}$), 4.23 (d, $J = 15.2$ Hz, 1H, $\text{CH}_2\text{-N}$), 4.59 (d, $J = 15.2$ Hz, 1H, $\text{CH}_2\text{-N}$), 5.02 (dd, $J = 3.7, 1.2$ Hz, 1H, CH-O), 7.30 (m, 3H, Ar), 7.36 (m, 2H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 44.6, 44.7, 55.2, 75.3, 78.8, 80.7, 127.8, 128.3, 128.8, 135.6, 165.8. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{13}\text{H}_{13}\text{NO}_2\text{Na}$ 238.0838, Found: 238.0845. HPLC [AD-H, hexane : *i*-propanole = 95 : 5, 0.7 mL/min, t_{R} [R]= 34.2 (minor), t_{R} [S]= 35.8 (major)].

(4S)-N-Benzyl-4-benzyloxyazetididin-2-one (18c). Yellow oil, yield 77%, 4% *ee*, 53 mg (0.20 mmol), IR (ν_{\max} , cm^{-1}): 1758 (C=O), $[\alpha]_{\text{D}}^{26} = -4.0$ (c 0.5, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.90 (dm, $J = 14.8$ Hz, 1H, CH_2), 3.04 (dd, $J = 14.8, 3.8$ Hz, 1H, CH_2), 4.18 (d, $J = 15.1$ Hz, 1H, $\text{CH}_2\text{-N}$), 4.42 (d, $J = 11.7$ Hz, 1H, $\text{CH}_2\text{-O}$), 4.47 (d, $J = 11.7$ Hz, 1H, $\text{CH}_2\text{-O}$), 4.59 (d, $J = 15.1$ Hz, 1H, $\text{CH}_2\text{-N}$), 4.98 (dd, $J = 3.8, 1.3$ Hz, 1H, CH-O), 7.19 (m, 2H, Ar), 7.30 (m, 8H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 43.8, 45.5, 70.0, 77.8, 127.5, 127.7, 128.5, 128.7, 128.9, 128.6, 135.3, 136.5, 166.3. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{17}\text{H}_{17}\text{NO}_2\text{Na}$ 290.1259, Found: 290.1262. HPLC [AD-H, hexane : *i*-propanole = 95 : 5, 0.7 mL/min, t_{R} [S]= 33.6 (major), t_{R} [R]= 35.9 (minor)].

(4S)-N-Benzyl-4-ethoxyazetididin-2-one (18d). Yellow oil, Yield 12%, *ee* not determined, 6 mg (0.03 mmol), IR (ν_{\max} , cm^{-1}): 1758 (C=O), $[\alpha]_{\text{D}}^{26} = -5.2$ (c 0.6, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 1.13 (t, $J = 6.9$ Hz, 3H, CH_3), 2.87 (dm, $J = 14.7$ Hz, 1H, CH_2), 3.03 (dd, $J = 14.7, 3.8$ Hz, 1H, CH_2), 3.44 (m, 2H, CH_2), 4.20 (d, $J = 15.2$ Hz, 1H, $\text{CH}_2\text{-N}$), 4.59 (d, $J = 15.2$ Hz, 1H, CH-N), 4.86 (dd, $J = 3.8, 1.5$ Hz, 1H, $\text{CH}_2\text{-O}$), 7.29 (m, 3H, Ar), 7.34 (m, 2H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 15.1, 44.3, 44.7, 63.1, 80.8, 127.7, 128.2, 127.8, 135.8, 166.1. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{12}\text{H}_{15}\text{NO}_2\text{Na}$ 228.0995, Found: 228.1000.

(4S)-N-Benzyl-4-isopropoxyazetididin-2-one (18e). Yellow oil, yield 15%, 11% *ee*, 9 mg (0.04 mmol), IR (ν_{\max} , cm^{-1}): 1756 (C=O), $[\alpha]_{\text{D}}^{26} = -3.1$ (c 0.71, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 1.09 (d, $J = 6.1$ Hz, 6H, $2 \times \text{CH}_3$), 2.82 (dm, $J = 14.7$ Hz, 1H, CH_2), 3.07 (dd, $J = 14.7, 3.7$ Hz, 1H, CH_2), 3.60 (septet, $J = 6.1$ Hz, 1H, $\text{CH}(\text{CH}_3)_2$), 4.13 (d, $J = 15.2$ Hz, 1H, CH-N), 4.85 (dd, $J = 3.7, 1.2$ Hz, 1H, CH-O), 7.29 (m, 3H, Ar), 7.34 (m, 2H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 22.5, 22.7, 44.2, 45.9, 71.0, 79.5, 127.7, 128.3, 128.8, 135.8, 166.2. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{13}\text{H}_{17}\text{NO}_2\text{Na}$ 242.1152, Found: 242.1153. HPLC [AD-H, hexane : *i*-propanole = 95 : 5, 0.5 mL/min, t_{R} [R]= 27.9 (minor), t_{R} [S]= 30.1 (major)].

Quinidine-catalyzed nucleophilic substitution with thioalcohols (15)

To the solution of 4-formyloxyazetididin-2-one (1) (30 mg, 0.26 mmol, 1.0 equiv.) and thioalcohol 15 (0.39 mmol, 1.5 equiv.) in toluene (5 mL) at room temperature was added quinidine (90 mg, 1.0 equiv.) under argon atmosphere. The reaction mixture was stirred for 48 h till disappearance of starting material and filtered through small layer of silica gel. The silica layer was washed with toluene (2 mL). Subsequently the filtrate containing 4-thio-azetididin-2-one 17 was slowly

added to solution of 50% aqueous NaOH (5 mL), and Bu₄NBr (84 mg, 0.26 mmol, 1.0 equiv.) and benzyl chloride (90 μ L, 0.78 mmol, 3.0 equiv.) in 1 mL of toluene. The reaction mixture was stirred for 30 min till disappearance of starting material and diluted with water (10 mL). The aqueous phase was extracted with ethyl acetate (3x15 mL). The organic extracts were combined and dried over MgSO₄. The solution was filtered and filtrate evaporated. The crude product was purified by column chromatography to yield 19.

(4S)-4-Allylthio-N-benzylzetidin-2-one (19a). Yellow oil, yield 40%, 38% *ee*, 23 mg (0.10 mmol), IR (ν_{\max} , cm⁻¹): 1755 (C=O), [α]_D²⁶ = -20.3 (c 1.0, CHCl₃). ¹H NMR (500 MHz, CDCl₃) δ_{H} 3.10 (m, 3H, CH₂, CH₂S), 3.34 (dd, *J* = 15.1, 4.9 Hz, 1H, CH₂), 4.07 (d, *J* = 15.3 Hz, 1H, CH₂-N), 4.53 (dd, *J* = 4.9, 2.2 Hz, 1H, CH-S), 4.70 (d, *J* = 15.3 Hz, 1H, CH₂-N), 4.90 (dd, *J* = 16.9, 1.1 Hz, 1H, CH₂=), 5.01 (d, *J* = 9.9 Hz, 1H, CH₂=), 5.70 (m, 1H, CH=), 7.30 (m, 3H, Ar), 7.35 (m, 2H, Ar). ¹³C NMR (CDCl₃) δ_{C} 32.5, 44.1, 45.1, 55.6, 117.8, 127.8, 128.3, 128.8, 133.7, 135.5, 165.4. HR MS (ESI) Calcd for [M + Na]⁺ C₁₃H₁₅NONaS 256.0767, Found: 256.0769. HPLC [AD-H, hexane : *i*-propanole = 95 : 5, 0.5 mL/min, *t*_R [R]= 34.2 (minor), *t*_R [S]= 35.8 (major)].

(4S)-N-Benzyl-4-benzylthioazetid-2-one (19b). Yellow oil, yield 86%, 34% *ee*, 62 mg (0.22 mmol), IR (ν_{\max} , cm⁻¹): 1758 (C=O), [α]_D²⁶ = -9.3 (c 1.0, CHCl₃). ¹H NMR (500 MHz, CDCl₃) δ_{H} 2.93 (dd, *J* = 15.1, 2.2 Hz, 1H, CH₂), 3.26 (dd, *J* = 15.1, 4.9 Hz, 1H, CH₂), 3.58 (d, *J* = 13.5 Hz, 1H, CH₂-S), 3.67 (d, *J* = 13.5 Hz, 1H, CH₂-S), 3.94 (d, *J* = 15.3 Hz, 1H, CH₂-N), 4.50 (dd, *J* = 4.9, 2.2 Hz, 1H, CH-S), 4.59 (d, *J* = 15.3 Hz, 1H, CH₂-N), 7.13 (m, 2H, Ar), 7.30 (m, 8H, Ar). ¹³C NMR (CDCl₃) δ_{C} 33.5, 44.1, 44.9, 55.9, 127.3, 127.8, 128.3, 128.6, 128.7, 128.8, 135.4, 137.5, 165.3. HR MS (ESI) Calcd for [M + Na]⁺ C₁₇H₁₇NONaS 306.0923, Found: 306.0921. HPLC [OD-H, hexane : *i*-propanole = 9 : 1, 0.5 mL/min, *t*_R [R]= 21.2 (minor), *t*_R [S]= 25.4 (major)].

(4S)-N-Benzyl-4-tert-butylthioazetid-2-one (19c). Yellow oil, yield 65%, 12% *ee*, 37 mg (0.15 mmol), IR (ν_{\max} , cm⁻¹): 1754 (C=O), [α]_D²⁶ = -3.4 (c 0.41, CHCl₃). ¹H NMR (500 MHz, CDCl₃) δ_{H} 1.27 (s, 9H, 3xCH₃), 2.99 (dm, *J* = 14.8 Hz, 1H, CH₂), 3.46 (dd, *J* = 14.8, 4.7 Hz, 1H, CH₂), 4.00 (d, *J* = 15.4 Hz, 1H, CH-N), 4.59 (dd, *J* = 4.7, 1.9 Hz, 1H, CH-O), 4.75 (d, *J* = 15.4 Hz, 1H, CH-N), 7.28 (m, 3H, Ar), 7.35 (m, 2H, Ar). ¹³C NMR (CDCl₃) δ_{C} 29.7, 31.1, 43.7, 48.9, 53.8, 127.6, 128.2, 128.7, 135.9, 165.8. HR MS (ESI) Calcd for [M + Na]⁺ C₁₄H₁₉NONaS 272.1080, Found: 272.1085. HPLC [AD-H, hexane : *i*-propanole = 95 : 5, 1.0 mL/min, *t*_R [R]= 15.1 (major), *t*_R [S]= 19.8 (minor)].

Quinidine-catalyzed nucleophilic substitution with thioalcohols (15)

To the solution of 4-formyloxyazetid-2-one 1 (30 mg, 0.26 mmol, 1.0 equiv.) and thioalcohol 15 (0.39 mmol, 1.5 equiv.) in toluene (5 mL) at room temperature was added quinidine (90 mg, 1.0 equiv.) under argon atmosphere. The reaction mixture was stirred for 48 h till disappearance of starting material and diluted with water (5 mL). Aqueous phase was extracted with ether (3x10 mL). The organic extracts were combined and dried over Na₂SO₄. The solution was

filtered and filtrate evaporated. The crude product was purified by column chromatography (silica gel, 2:3 hexane/diethyl ether).

(4S)-4-Tritylthioazetid-2-one (20). White solid, yield 89%, 51% *ee*, 80 mg (0.23 mmol), m.p. 140-142°C, IR (ν_{\max} , cm^{-1}): 1772 (C=O), $[\alpha]_{\text{D}}^{26} = 12.5$ (c 1.0, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.88 (ddd, $J = 15.2, 2.7, 1.6$ Hz, 1H, CH_2), 3.27 (ddd, $J = 15.2, 5.3, 1.6$ Hz, 1H, CH_2), 4.44 (dd, $J = 5.3, 2.7$ Hz, 1H, CH-O), 7.26 (m, 3H, Ar), 7.32 (m, 6H, Ar), 7.46 (m, 6H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 45.0, 52.4, 67.3, 127.1, 127.9, 128.3, 129.3, 143.9, 165.7. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{22}\text{H}_{19}\text{NONaS}$ 368.1080, Found: 368.1088. HPLC [OD-H, hexane : *i*-propanole = 8 : 2, 0.5 mL/min, $t_{\text{R}}[\text{S}] = 15.0$ (major), $t_{\text{R}}[\text{R}] = 21.2$ (minor)].

Chiral Lewis Acids 7 promoted deracemization

The chiral Lewis acid promoter **7** was generated *in situ* by the addition of 1M solution of SnCl_4 in CH_2Cl_2 (420 μL , 1.0 equiv.) to the solution of (*S*) or (*R*) α -di-naphthyl-BINOL (225 mg, 0.42 mmol, 1.0 equiv.) in CH_2Cl_2 (10 mL) at 0°C under argon atmosphere. After 15 minutes racemic azetidione (0.42 mmol, 1.0 equiv.) in appropriate temperature was added. The reaction mixture was stirred for 6h to 36h and diluted with saturated NaHCO_3 (10 mL). The aqueous phase was extracted with CH_2Cl_2 (3x20 mL). The organic extracts were combined and dried over MgSO_4 . The solution was filtered and filtrate evaporated. The crude product was purified by column chromatography to yield **11a**, **11d**, **21a-b**, **22** or **23**.

(4R)-N-Benzyl-4-phenylthioazetid-2-one (21a). White solid, yield 40%, 37% *ee*, 46 mg (0.17 mmol), m.p. 129-130°C, IR (ν_{\max} , cm^{-1}): 1779 (C=O), $[\alpha]_{\text{D}}^{26} = +50.7$ (c 0.6, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.99 (ddd, $J = 14.9, 2.2, 1.0$ Hz, 1H, CH_2), 3.29 (ddm, $J = 14.9, 4.9$ Hz, 1H, CH_2), 4.58 (d, $J = 15.1$ Hz, 1H, CH-N), 4.73 (d, $J = 15.1$ Hz, 1H, CH-N), 4.79 (dd, $J = 4.9, 2.2$ Hz, 1H, CH-S), 7.14 (m, 2H, Ar), 7.32 (m, 8H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 44.3, 44.5, 58.2, 127.8, 128.4, 128.7, 128.8, 129.3, 130.4, 134.1, 135.4, 165.1. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{16}\text{H}_{15}\text{NONaS}$ 292.0767, Found: 292.0771. HPLC [OD-H, hexane : *i*-propanole = 9 : 1, 0.5 mL/min, $t_{\text{R}}[\text{S}] = 19.9$ (minor), $t_{\text{R}}[\text{R}] = 23.8$ (major)].

(4R)-N-Benzyl-4-(*p*-methoxyphenylthio)-azetid-2-one (21b). White solid, yield 43%, 21% *ee*, 54 mg (0.18 mmol), m.p. 131-132°C, IR (ν_{\max} , cm^{-1}): 1776 (C=O), $[\alpha]_{\text{D}}^{26} = +16.7$ (c 0.6, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.89 (ddd, $J = 15.1, 2.2, 1.0$ Hz, 1H, CH_2), 3.29 (ddd, $J = 15.1, 4.9, 1.6$ Hz, 1H, CH_2), 3.81 (s, 3H, CH_3), 4.58 (d, $J = 15.1$ Hz, 1H, CH-N), 4.73 (d, $J = 15.1$ Hz, 1H, CH-N), 4.87 (dd, $J = 4.9, 2.2$ Hz, 1H, CH-S), 6.89 (d, $J = 8.7$ Hz, 2H, Ar), 7.14 (m, 2H, Ph), 7.32 (m, 3H, Ph), 7.43 (d, $J = 8.7$ Hz, 2H, Ar). ^{13}C NMR (CDCl_3) δ_{C} 44.8, 46.5, 54.5, 56.0, 114.9, 118.2, 121.3, 126.2, 127.6, 128.1, 128.7, 130.4, 134.1, 158.4, 166.1. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{17}\text{H}_{17}\text{NO}_2\text{NaS}$ 322.0980, Found: 322.0976. HPLC [OD-H, hexane : *i*-propanole = 9 : 1, 0.5 mL/min, $t_{\text{R}}[\text{S}] = 21.6$ (minor), $t_{\text{R}}[\text{R}] = 25.8$ (major)].

(6R)-3,4-Benzocepham (22). Yellow oil, yield 48%, 46% *ee*, 38 mg (0.2 mmol), IR (ν_{\max} , cm^{-1}): 1764 (C=O), $[\alpha]_{\text{D}}^{26} = +20.2$ (c 0.41, CHCl_3). ^1H NMR (500 MHz, CDCl_3) δ_{H} 2.97 (dm, $J = 14.8$ Hz, 1H, CH_2), 3.51 (ddm, $J = 14.8, 4.1$ Hz, 1H, CH_2), 4.25 (d, $J = 16.6$ Hz, 1H, CH-N), 4.79 (d, $J = 16.6$ Hz, 1H, CH-N), 4.81 (dm, $J = 4.1$ Hz, 1H, CH-S), 7.17 (m, 3H, Ar), 7.24 (m, 1H, Ar).

^{13}C NMR (CDCl_3) δ_{C} 42.1, 45.9, 49.1, 126.2, 127.7, 128.3, 128.4, 129.5, 130.0, 166.6. HR MS (ESI) Calcd for $[\text{M} + \text{Na}]^+ \text{C}_{10}\text{H}_9\text{NONaS}$ 214.0297, Found: 214.0288. HPLC [OD-H, hexane : *i*-propanole = 9 : 1, 1.0 mL/min, $t_{\text{R}}[\text{R}]$ = 20.9 (major), $t_{\text{R}}[\text{S}]$ = 25.5 (minor)].

(6R)-3-Methylene-5-oxacepham (23). Yield 48%, 22% *ee*, 28 mg (0.2 mmol), $[\alpha]_{\text{D}}^{26} = +9.2$ (c 0.8, CHCl_3). HPLC [AD-H, hexane : *i*-propanole = 95 : 5, 1.0 mL/min, $t_{\text{R}}[\text{R}]$ = 14.9 (major), $t_{\text{R}}[\text{S}]$ = 20.7 (minor)].

Assay of DD-carboxypeptidase activity

The enzyme activity was measured as described previously.^{154,153} Samples for assay of the DD-carboxypeptidase activity consisted of 10 μL of DD-carboxypeptidase from *Saccharopolyspora erytraea* PZH TZ 64-575 (40 units/mg), 20 μL of substrate solution containing 4.52 mg/mL *N*, *N* ϵ -diacetyl-L-lysyl-D-alanyl-D-alanine in 0.1 M phosphate buffer, pH 8.0 and 10 μL of 0.1 M phosphate buffer, pH 8.0. A standard sample containing 20 μL of D-alanine in distilled water. The reaction mixture for assay of the DD-carboxypeptidase activity consisted of 60 μL of 0.05 mg/mL flavin adenine dinucleotide in 0.1 M phosphate buffer, pH 8.0, 10 μL of 0.05 mg/mL horseradish peroxidase (1230 units/mg) in distilled water, 5.0 μL of 5.0 mg/mL *o*-dianisidine in methanol, and 2.0 μL of 11.77 mg/mL D-amino acid oxidase from porcine kidney (6.7 units/mg) in 0.1 M phosphate buffer, pH 8.0. Samples were incubated for 30 minutes at 37 $^{\circ}\text{C}$ and then boiled for 2 minutes. After cooling, 77 μL of the reaction mixture was added, and all samples were incubated for 10 minutes at 37 $^{\circ}\text{C}$. Next, was added 350 μL to each sample of a mixture consisting of MeOH, distilled water and sulfuric acid (5 : 5 : 6 by volume). Extinction of the resulting solution was measured at 540 nm. The inhibition of DD-peptidase 64-575 by the discussed compounds was evaluated. Mixtures of 10 μL of DD-peptidase 64-575 (40 units/mg), 5.0 μL of 0.1 M phosphate buffer, pH 8.0 were incubated for 45 minutes at 37 $^{\circ}\text{C}$. The concentration of tested compounds in the mixture was from 0.04 to 0.0001 M. following the incubation, 20 μL of substrate solution was added to 20 μL of each sample and the resulting mixtures were incubated again.

Assay of β -lactamase activity

The inhibition of penicillinase was evaluated following a literature method. The sample for assay of inhibition of β -lactamase consisted of 10 μL of penicillinase (Penase, 5×10^6 IU/mL, Bacto), 20 μL 0.1 M phosphate buffer, pH 7.0, 10 μL solution of tested compound in methanol. The samples were incubated for 30 minutes at 37 $^{\circ}\text{C}$, then 30 μL of nitrocephin, 430 μL 0.1 M phosphate buffer pH 7.0 were added and all samples were incubated for 10 minutes at 37 $^{\circ}\text{C}$. Absorption was measured at 428 nm.

Acknowledgements

This work was supported by the Ministry of Education and Science, grant N N204 192738. Anna Koziol thanks the President of Polish Academy of Sciences for a doctoral fellowship.

References

1. (a) Morin, R. B.; Gorman, M. *Chemistry and Biology of β -Lactam Antibiotics*; Academic: New York, 1982; Vol. 1-3. (b) Bruggink, A. *Synthesis of β -Lactam Antibiotics*; Kluwer: Dordrecht, Netherlands, 2001.
2. Shah, S. K.; Dorn, C. P.; Finke, P. E.; Hale, J. J.; Hagmann, W. K.; Brause, K. A.; Chandler, G. O.; Kissinger, A. L.; Ashe, B. M.; Weston, H.; Knight, W. B.; Maycock, A. L.; Dellea, P. S.; Fletcher, D. S.; Hand, K. M.; Mumford, R. A.; Underwood, D. J.; Doherty, J. B. *J. Med. Chem.* **1992**, *35*, 3745.
3. Han, W. T.; Trehan, A. K.; Wright, J. J. K.; Federici, M. E.; Seiler, S. M.; Meanwell, N. A. *Bioorg. Med. Chem.* **1995**, *3*, 1123.
4. (a) Borthwick, A. D.; Weingarten, G.; Haley, T. M.; Tomaszewski, M.; Wang, W.; Hu, Z.; Bedard, J.; Jin, H.; Yuen, L.; Mansour, T. S. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 365. (b) Deziel, R.; Malenfant, B. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 143. (c) Yoakim, C.; Ogilvie, W.W.; Cameron, D.R.; Chabot, C.; Guse, I.; Hache, B.; Naud, J.; O'Meara, J.A.; Plante, R.; Deziel, R. *J. Med. Chem.* **1998**, *41*, 2882. (d) Ogilvie, W. W.; Yoakim, C.; Do, F.; Hache, B.; Lagace, L.; Naud, J.; O'Meara, J.A.; Deziel, R. *Bioorg. Med. Chem.* **1999**, *7*, 1521. (e) Bonneau, P. R.; Hasani, F.; Plouffe, C.; Malenfant, E.; LaPlane, S. R.; Guse, I.; Ogilvie, W.W.; Plante, R.; Davidson, W. C.; Hopkins, J. L.; Morelock, M. M.; Cordingley, M. G.; Deziel, R. *J. Am. Chem. Soc.* **1999**, *121*, 2965.
5. Rosenblum, S. B. In *The Art of Drug Synthesis*; Johnson, D. S.; Li, J. J. Eds.; Wiley: Hoboken, 2007, Chapter 13, p 183.
6. (a) Wild, H. In *The Organic Chemistry of β -Lactams*; Georg, G. I. Eds.; VCH Publishers: Weinheim, 1993; p 49. (b) Ojima, I. *Acc. Chem. Res.*, **1995**, *28*, 383. (c) Ojima, I.; Delalogue, F. *Chem. Soc. Rev.* **1997**, *26*, 377; (d) Singh, G. S. *Tetrahedron* **2003**, *59*, 7631. (e) Palomo, C.; Aizpurua, J. M.; Ganboa, I.; Oiarbide, M. *Curr. Med. Chem.* **2004**, *11*, 1837. (f) Fisher, J. F.; Meroueh, S. O.; Mobashery, S. *Chem. Rev.* **2005**, *105*, 395. (g) Alcaide, B.; Almendros, P.; Aragoncillo, C. *Chem. Rev.* **2007**, *107*, 4437. (h) D'hooghe, M.; Dekeukeleire, S.; Leemans, E.; De Kimpe, N. *Pure Appl. Chem.*, **2010**, *82*, 1749.
7. Kozioł, A.; Furman, B.; Frelek, J.; Woźnica, M.; Altieri, E.; Chmielewski, M. *J. Org. Chem.* **2009**, *74*, 5687.
8. Nagaraja, S.R.; More O'Ferral, R. A. *J. Am. Chem. Soc.* **1990**, *112*, 2729.
9. Kozioł, A.; Frelek, J.; Woźnica, M.; Furman, B.; Chmielewski, M. *Eur. J. Org. Chem.* **2009**, 338.
10. Veysoglu T.; Mitscher L. A.; Swayze J. K. *Synthesis*, **1980**, 807.
11. Kałuża, Z.; Kazimierski, A.; Lewandowski, K.; Suwińska, K.; Szczęsna, B.; Chmielewski, M. *Tetrahedron* **2003**, *59*, 5893.
12. Knox, J. R.; Moews, P. C., *J. Mol. Biol.* **1991**, *220*, 435.

13. a) Kurzatkowski, W.; Solecka, J.; Filipek, J.; Kurzatkowski, J. D.; Kurylowicz, W. *Appl. Microbiol. Biotechnol.* **1990**, *33*, 452; b) Rohl, F.; Rabenhorst, J.; Zahner, H. *Arch. Microbiol.* **1987**, *147*, 315.