

Supplementary Material

Efficient synthesis of 2,3-dihydroquinazolin-4(*1H*)-ones using heterogeneous solid acid catalysts: unexpected formation of 2,3-dihydro-2-(4-(tetrahydro-2*H*-pyran-2-yloxy)butyl)quinazolin-4(*1H*)-one

Sandip B. Bharate,* Nagaraju Mupparapu, Sudhakar Manda, Jaideep B. Bharate, Ramesh Mudududdla, Rammohan R. Yadav, and Ram A. Vishwakarma*

Medicinal Chemistry Division, Indian Institute of Integrative Medicine (Council of Scientific and Industrial Research), Canal Road, Jammu-180001, India


E-mail: sbharate@iiim.ac.in ; ram@iiim.ac.in

Table of Contents


1.	¹ H NMR spectrum of compound 6	S2
2.	¹³ C NMR spectrum of compound 6	S2
3.	DEPT135 NMR spectrum of compound 6	S3
4.	MS NMR spectrum of compound 6	S3
5.	Key HMBC correlations of compound 6	S4

1. ^1H NMR spectrum of 6


J.B.21 (A)

2. ^{13}C NMR spectrum of compound 6


JB-21A




3. DEPT135 NMR spectrum of compound 6



4. MS spectrum of compound 6



5. Key HMBC correlations of compound 6

HMBC correlations of compound 6