

Aluminium(III) chloride hexahydrate: an efficient and versatile reagent in organic synthesis

Saikat Das Sarma, Pallab Pahari, Swapnali Hazarika, Parasa Hazarika, Manash Jyoti Borah, and Dilip Konwar*

*Synthetic Organic Chemistry Division, CSIR - North East Institute of Science & Technology
Jorhat-785006, Assam, India
E-mail: dkonwar@yahoo.co.uk*

DOI: <http://dx.doi.org/10.3998/ark.5550190.p007.863>

Abstract

A short review on the application of aluminium chloride hexahydrate [AlCl₃.6H₂O], a mild, comparatively less toxic, and easily available reagent has been presented. The article is covering those areas of organic synthesis where the AlCl₃.6H₂O was used either as a single reagent or in combination with a co-catalyst.

Keywords: Aluminium chloride hexahydrate, isoxazolidine, chemoselective reduction, deoxygenation, dehydration

Table of Contents

1. Introduction
2. Reduction reactions
 - 2.1 Reductive cleavage of 2,1-benzisoxazoles
 - 2.2 Reductive cleavage of isoxazolidines
 - 2.3 Chemoselective reduction of ketones
 - 2.4 Chemoselective reduction of C=C bonds of α , β -unsaturated carbonyl compounds
 - 2.5 Deoxygenation of organic N-oxides
 - 2.6 Reduction of azides
3. Dehydration of oximes and amides in hydrated media
4. Protection and deprotection
 - 4.1 Tetrahydropyranylation and depyranylation of alcohols and phenols
 - 4.2 Chemoselective deprotection of esters, ethers, and ketals
5. Synthesis of 1,4-dihydropyridines and their aromatization
6. Oxidation of alcohols to carbonyls
7. Miscellaneous reactions

- 7.1 Esterification of glycerine with lauric acid
- 7.2 Preparation of ionic liquids bearing tetrachloroaluminate anions
- 7.3 Iodination of benzyl alcohols
- 8. Conclusions
- 9. Acknowledgement
- 10. References

1. Introduction

Development of non-hazardous synthetic methodologies for organic reactions is one of the latest challenges to the organic chemists. In this connection, the use of efficient, non-toxic and inexpensive reagents for various organic transformations is an important research area. Aluminium, a group 13 element, is the most abundant metal in the earth's crust. Anhydrous AlCl_3 has been extensively used as a strong Lewis acid in many organic reactions such as Friedel-Crafts reactions, Fries rearrangement, Scholl reaction etc.¹ But, the utility of the hydrated counterpart i.e. $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ is not fully explored yet.

Aluminium(III) chloride hexahydrate is a non-flammable; yellowish-white to colorless deliquescent granular crystals with slight HCl odor. It is highly soluble in water and also soluble in alcohol, ether, glycerol, propylene glycol and acetonitrile. It finds wide application in deodorants and antiperspirant preparations, refining crude oil, dyeing fabrics, manufacturing parchment paper, textile finishing, preserving wood and disinfecting stables. It is also clinically used to control severe excessive sweating called hyperhidrosis. Generally the reagent is prepared either by treating conc. hydrochloric acid to the solution of anhydrous AlCl_3 in water followed by saturating with dry hydrogen chloride or heating aluminium hydroxide with conc. hydrochloric acid in a sealed tube. It has been observed that $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ exists in reality as $[\text{Al}(\text{H}_2\text{O})_6]\text{Cl}_3$ and its aqueous solution is acidic due to the ability of the hydrated cation $[\text{Al}(\text{H}_2\text{O})_6]^{+3}$ to act as proton donor giving $[\text{Al}(\text{H}_2\text{O})_5(\text{OH})]^{2+}$, $[\text{Al}(\text{H}_2\text{O})_4(\text{OH})_2]^+$, etc.^{1c}


The cost effectiveness, ease of handling, good reactivity, experimental simplicity, and excellent solubility in water and organic solvents render $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ as a promising green reagent and catalyst in organic synthesis. In this review, we will outline over a decade of research work on the development of $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ as a reagent and catalyst in various organic reactions by different group of scientists including us.

2. Reduction reactions

2.1. Reductive cleavage of 2,1-benzisoxazoles

$\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ in combination with metallic zinc in moist THF can cleave the 2,1-benzisoxazoles **1** and their ammonium salts **3** to yield *o*-amino and *N*-alkylaminobenzophenones **2** & **4**


respectively (Scheme 1).² *o*-Amino and *N*-alkylaminobenzophenones³ are the key intermediates for the synthesis of several clinically used drugs of 1,4-benzodiazepine class, e.g., oxazepam, diazepam, flurazepam, alprazolam etc.⁴ In addition, appropriately substituted 2-aminobenzophenone derivatives act as antiproliferative agents and antimitotic agents.⁵ A wide range of derivatives of 2,1-benzisoxazoles and its salts was successfully investigated under this protocol. It was found that the cleavage of 2,1-benzisoxazoles to furnish corresponding *o*-aminobenzophenones occurs in better yield (88-92%) compared to its salts which produced corresponding *N*-alkylaminobenzophenones in moderate yield (60-65%).



Scheme 1

2.2. Reductive cleavage of isoxazolidines

Zn-AlCl₃·6H₂O system can also perform the reductive cleavage of the N-O bond of isoxazolidines **5** to produce 1,3-amino alcohols **6**.⁶ A variety of isoxazolidine derivatives have been examined and the conversion found to proceed successfully under ambient conditions within a period of short reaction time (Scheme 2). Naturally occurring 1,3-amino alcohols as well as synthetic derivatives thereof exhibit wide ranges of biological activities.⁷ They are also widely used as ligands for asymmetric catalysts and synthetic intermediates.⁸



Scheme 2

2.3. Chemoselective reduction of ketones

Chowdhury and Borah⁹ developed a mild and chemoselective method for the reduction of aliphatic ketones to the corresponding methylene derivatives using Zn-AlCl₃.6H₂O-THF-H₂O system at room temperature. A broad range of structurally diverse ketones **7a-j** have been tested using the catalytic system (Table 1). It was observed that the reaction was more successful for the cyclic ketones (entry 1-5, Table 1) compared to the acyclic ketones (entry 6, Table 1). The reagent could chemoselectively reduce a six membered ring ketone in the preference of a five membered one (entry 3, Table 1). The free ester group and isolated double bonds were unaffected under the reaction condition. It was noteworthy that the ketone **7e** (entry 5, Table 1) was deoxygenated efficiently by this method which under Clemmensen¹⁰ or Wolf-Kishner reduction¹¹ condition produced a mixture of products. Interestingly, aliphatic and aromatic aldehydes (**7g** & **7h**), aromatic ketones **7i** and α,β -unsaturated ketones **7j** were reduced to the corresponding alcohols (entry 7-10, Table 1).

Table 1. Chemoselective reduction of carbonyl compounds by Zn-AlCl₃.6H₂O-THF-H₂O system



Entry	Substrates	Products	Yield (%)
1	 7a	 8a	50

Table 1. Continued
















Entry	Substrates	Products	Yield (%)
2	 7b	 8b	90
3	 7c	 8c	80
4	 7d	 8d	60
5	 7e	 8e	70
6	 7f	 8f	20
7	$\text{CH}_3(\text{CH}_2)_{14}\text{CHO}$ 7g	$\text{CH}_3(\text{CH}_2)_{14}\text{CH}_2\text{OH}$ 8g	70
8	PhCHO 7h	PhCH_2OH 8h	90

Table 1. Continued

Entry	Substrates	Products	Yield (%)
9	 7i	 8i	85
10	 7j	 8j	90

2.4. Chemoselective reduction of C=C bonds of α , β -unsaturated carbonyl compounds



Scheme 3

Konwar et al. showed that the Zn-AlCl₃·6H₂O combination in moist THF can act as an excellent reducing system for the chemo-selective reduction of carbon-carbon double bond of the α,β -unsaturated carbonyl compounds **9** (Scheme 3).¹² The methodology was generalized using different α,β -unsaturated ketones (entry 1-4, table 2), esters (entry 5-6, table 2) and acids (entry 7, table 2). It was observed that the process was also successful for the reduction of ene-diacid (entry 7, Table 2) and ene-diester (entry 6, Table 2). The quinones were reduced to the corresponding dihydroquinones in excellent yields (entries 8 and 9, Table 2). Notably, aromatic and heteroaromatic rings were not affected under the reaction condition.

Table 2. Selective reduction of carbon-carbon double bonds by Zn-AlCl₃·6H₂O system



















Entry	Substrates	Products	Yield (%)
1	 9a	 10a	90
2	 9b	 10b	88

Table 2. Continued

Entry	Substrates	Products	Yield (%)
3	 9c	 10c	60
4	 9d	 10d	82
5	 9e	 10e	88
6	 9f	 10f	98
7	 9g	 10g	95
8	 9h	 10h	100
9	 9i	 10i	98

2.5. Deoxygenation of organic N-oxides

Dutta and Konwar used $Zn-AlCl_3 \cdot 6H_2O$ system for the N-deoxygenation of nitrones **11a-g**, heteroarene N-oxides **11h** and azoxy benzenes **11i** to their corresponding imines.¹³ The sensitive carbon-carbon double bonds, furan and thiophene remained unaffected under the reaction conditions. Later on Boruah and Konwar replaced zinc with KI and developed even milder reagent system $AlCl_3 \cdot 6H_2O/KI/CH_3CN/H_2O$ which was effective for the similar N-deoxygenations in hydrated medium at room temperature (Table 3).¹⁴ The best result was obtained when acetonitrile and water was used in 2:1 ratio in the reaction mixtures. Similar to the earlier reagent system the $AlCl_3 \cdot 6H_2O/KI$ also did not affect the aromatic or heteroaromatic rings.

Table 3. Reduction of organic N-oxides using AlCl₃.6H₂O/KI






















Entry	Substrates	Products	Yield (%)
1	 11a	 12a	90
2	 11b	 12b	92
3	 11c	 12c	85
4	 11d	 12d	80
5	 11e	 12e	72
6	 11f	 12f	75
7	 11g	 12g	70
8	 11h	 12h	80

Table 3. Continued


Entry	Substrates	Products	Yield (%)
9	 11i	 12i	72
10	 11j	 12j	75

But the expected mechanism for the N-deoxygenation using KI/ $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ is different from the Zn/ $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ system. It is proposed that the $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ which in reality exist as $[\text{Al}(\text{H}_2\text{O})_6]\text{Cl}_3$ in solution^{1c} reacts with KI to form $[\text{Al}(\text{H}_2\text{O})_6]\text{I}_3$. The iodide ion thus produced adds to the double bond of **13** and with the help of aluminium performs the deoxygenation to imines **15** through the formation of the intermediate **14** (Scheme 4).¹⁴

**Scheme 4.** Proposed mechanism for the deoxygenation of nitrones

2.6. Reduction of azides

The reduction of azides is a most common strategy for the synthesis of primary amines.¹⁵ Li et al.¹⁶ found that Zn/ $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ could be used as an effective reagent for the reduction of azides to the corresponding primary amines in water or aq. ethanol at ambient condition. In another method, Zheng et al.¹⁷ also efficiently reduced azides **16** to their corresponding primary amines **17** employing $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ in combination with iron (Scheme 5). The mild reaction condition did not affect sensitive functional groups like nitro, halides, carbonyl and ethers.



Scheme 5. Reduction of azides using Fe-AlCl₃·6H₂O system

3. Dehydration of oximes and amides in hydrated media

Boruah and Konwar¹⁸ showed that the AlCl₃·6H₂O/KI reagent system could perform the dehydration of aldoximes **18a-f** and amides **18g** to their corresponding nitriles. The interesting fact was that the dehydration actually was taking place in aqueous medium. The reaction was performed in CH₃CN/H₂O solvent system at 80 °C and there was no report of formation of side products like triazines. The authors showed flexibility of the reagent system using both aliphatic and aromatic aldoximes and amides (entry 1-7, table 4). The mild reaction condition allowed the sensitive functional groups like methyl ether, furan, thiophene etc. to remain intact. It was observed that cyclodehydration of appropriately substituted amides produced isoquinoline derivatives in a similar manner to the Bischler–Napieralski reaction (entry 8, table 4). Interestingly, when ketoximes were allowed to react with the reagent system, the Beckmann rearrangement took place. So open chain ketoximes produced anilides or amides (entry 9, table 4) and cyclic ketoximes produced corresponding lactams (entry 10, table 4).

Table 4. Dehydration of oximes and amides using AlCl₃·6H₂O/KI





















Entry	Substrates	Products	Yield (%)
1	 18a	 19a	96
2	 18b	 19b	75


Table 4. Continued

Entry	Substrates	Products	Yield (%)
3	 18c	 19c	69
4	 18d	 19d	60
5	 18e	 19e	67
6	 18f	 19f	75
7	 18g	 19g	92
8	 18h	 19h	87
9	 18i	 19i	82
10	 18j	 19j	72

4. Protection and deprotection

4.1. Tetrahydropyranylation and depyranylation of alcohols and phenols

Varma and Namboodiri¹⁹ reported a $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ catalyzed solvent free tetrahydropyranylation of alcohols and phenols by the reaction of aliphatic and aromatic alcohols **20** with 3,4-dihydro-2H-pyran (DHP) (**21**) (Scheme 6). A wide variety of alcohol and phenolic compounds (e.g., 3-hexanol, 4-butanediol, *t*-butanol, cyclohexanol, allyl alcohol, geraniol, 4-chlorobenzyl alcohol, phenol, 4-chlorophenol, 2-naphthol, benzhydrol etc.) were tested and in each case the corresponding THP ethers produced in good to excellent yield (74-96%). No dehydration reaction occurred when acid sensitive alcohols like *t*-butanol were pyranylated under the reaction condition. It was reported that the rate of the pyranylation reaction decreased with increase in substrate to catalyst ratio. In that case increase in the reaction temperature compensated the effect. Also substituted benzyl alcohols and solid alcohols needed comparatively higher temperature (60-80 °C) for the pyranylation. This method can be used for the mono tetrahydropyranylation of diols which is difficult to achieve with conventional methods. The method is also suitable for the moist substrates although the rate of the reaction decreases with increase in the water content. Interestingly, the same reagent can be used for the deprotection of the generated THP ethers. The deprotection is carried out at high substrate to catalyst ratio (500 mmol: 1 mmol) in excess methanol.




















Scheme 6

4.2. Chemoselective deprotection of esters, ethers, and ketals

The $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}/\text{KI}$ system is also useful for the cleavage of the C–O bonds of esters **24a-f**, ethers **24g**, and 1,3-oxathiolanes **24h-j**.²⁰ The reactions are performed in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ medium at 80 °C. Under the reaction conditions, esters (entries 1-3, Table 5) and acetates (entries 4-6, Table 5) are converted to the corresponding acids and phenols, respectively. The reaction can be used for the chemoselective cleavage of the acetate functionalities in the presence of esters (entry 6, Table 5). Selective cleavage of either esters or acetates can also be achieved successfully without affecting the ether functionalities (entries 3 and 4, Table 5). But ethers also can be cleaved under the reaction condition to provide corresponding alcohols and iodides when only ether functionality is present in the molecule (entry 7, Table 5). The reagent system effectively can deprotect 1,3-oxathiolanes to the corresponding carbonyls (entries 8-10, Table 5). In these cases the deprotection can effectively be performed without disturbing either the ethers (entries 8 and 10, Table 5) or the esters (entry 9, Table 5).


Table 5. Deprotection of esters, ethers, and ketals by $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}/\text{KI}/\text{CH}_3\text{CN}/\text{H}_2\text{O}$ system

Entry	Substrate	Product	Yield (%)
1	$\text{CH}_2(\text{COOEt})_2$ 24a	$\text{CH}_2(\text{COOH})_2$ 25a	65
2	 24b	Ph-COOH 25b	78
3	 24c	 25c	76
4	 24d	 25d	83
5	 24e	 25e	90
6	 24f	 25f	80
7	 24g	 25g	89
8	 24h	 25h	90
9	 24i	 25i	84
10	 24j	 25j	77

5. Synthesis of 1,4-dihydropyridines and their aromatization

The versatility of the $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ as reagent was proved when it was found to catalyze the one pot three component Hantzsch dihydropyridine synthesis. So a 10 mol% $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ was able to catalyze the condensation of an aldehyde **26**, a beta keto ester **27** and NH_4OAc to produce dihydropyridines **28** at 60 °C in high yields under solvent free conditions (Scheme 7).²¹ A broad range of structurally diverse aldehydes (both aromatic and aliphatic) and 1,3-dicarbonyl compounds were used to produce the corresponding dihydropyridines. The sensitive functional groups like furans were unaffected under the reaction condition. Interestingly, it was found that $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ also could catalyze the oxidation of dihydropyridines **28** to the corresponding pyridines **29** in the presence of H_2O_2 under microwave irradiation. Most probably the liberated HCl from $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ reacts with H_2O_2 to produce HOCl which performs the oxidation through the formation of a chlorine free radical.

As both the two steps in the synthesis of pyridines **29** were catalyzed by $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$, it was also possible to combine them in one pot following the same protocol. So, substituted pyridines were synthesized by the condensation of aldehyde **26**, beta keto ester **27** and NH_4OAc in the presence of 10 mol% $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ and subsequent oxidation with 30% H_2O_2 in the same pot under microwave irradiation. The product was not separated or purified after the first step and no additional amount of $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ was added in the oxidation step.




Scheme 7. $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ catalyzed synthesis of 1,4-dihydropyridines and pyridines.

Entry	R ¹	R ²	Yield of dihydropyridines 28	Yield of pyridines 29
1	C ₆ H ₅	Et	80	99
2	4-MeOC ₆ H ₄	Et	77	97
3	4-O ₂ NC ₆ H ₄	Et	75	96
4	4-ClC ₆ H ₄	Et	76	97
5	CH ₃	Et	74	96
6	CH ₃ CH ₂	Et	76	95
7	(CH ₃) ₂ CH	Et	78	100
8	2-Furyl	Et	73	94
9	C ₆ H ₅	Me	78	98
10	4-MeOC ₆ H ₄	Me	75	97

Yield refers to isolated yield (%).

6. Oxidation of alcohols to carbonyls

Chang et al.²² developed an efficient and environmentally friendly method for the oxidation of benzylic alcohols **30** (both primary and secondary) to aldehydes and ketones **31** using $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ and NCS in ionic liquid $[\text{bmim}]\text{BF}_4$ (Scheme 8). A broad range of benzylic alcohols bearing both electron-donating and electron-withdrawing substituents were oxidized in this oxidation condition. Primary benzylic alcohols reacted at a faster rate compared to secondary alcohols. It was proposed that the oxidation reactions were accomplished by the reaction of alcohols with chlorium ion which was formed by the interaction of NCS with $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$. Gogoi et al.²⁰ also reported a similar oxidation of both primary and secondary alcohols using $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}/\text{KI}$ along with DMSO under refluxing condition.



$\text{R}^1 = \text{H, Me, Et, Ph, PhCO}$

$\text{R}^2 = \text{H, MeO}$

$\text{R}^3 = \text{Me, F, Cl, Br, NO}_2$


$\text{R}^4 = \text{H, MeO}$

Scheme 8. Oxidation of benzylic alcohols using NCS/ $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$

7. Miscellaneous reactions

7.1. Esterification of glycerine with lauric acid

Sugi and coworkers²³ described an efficient method for the selective esterification of glycerine (**32**) with lauric acid using $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ as the catalyst in solvent free condition. The reaction selectively produced lauric acid monoglyceride (monolaurin) **33** in a moderate yield (Scheme 9). It was proved that the product was formed by a direct mono esterification of glycerine, not via the transesterification of lauric acid triglyceride (trilaurin). When the reaction was performed in excess lauric acid the process also produced di and triglycerides, albeit in a lower proportion. The mono- and di-glycerides of long chain fatty acids are widely used in the food, cosmetic and pharmaceutical industry.



Scheme 9. Monoesterification of glycerine with lauric acid

7.2. Preparation of ionic liquids bearing tetrachloroaluminate anions


Namboodiri and Varma²⁴ reported an efficient method for the preparation of ionic liquid dialkylimidazolium tetrachloroaluminates $[\text{C}_n\text{MIM}][\text{AlCl}_4]$ **35** by the reaction of N,N' -dialkylimidazolium chloride **34** and $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ in quantitative yield using microwave irradiation (Scheme 10). The authors have prepared three different ionic liquids namely $[\text{C}_4\text{MIM}][\text{AlCl}_4]$, $[\text{C}_6\text{MIM}][\text{AlCl}_4]$, and $[\text{C}_8\text{MIM}][\text{AlCl}_4]$ from respective N,N' -dialkylimidazolium chlorides using the procedure. It is to be noted that the ionic liquids with tetrachloroaluminate anion are very useful for several organic reactions. The utility of the synthesized ionic liquids was established by using them as recyclable catalyst for the tetrahydropyranlation and depyranlation of different alcohols.



Scheme 10. Preparation of $[\text{C}_n\text{MIM}][\text{AlCl}_4]$ using microwave irradiation

7.3. Iodination of benzyl alcohols

The $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}/\text{KI}$ system can also act as an excellent iodinating agent for the iodination of benzyl alcohols **36** to benzyl iodides **37** in hydrated media (Scheme 11).²⁰ The reaction is equally applicable for different type of benzyl alcohols carrying both electron-donating and withdrawing groups.



Scheme 11. Iodination of benzyl alcohols

8. Conclusions

The exploration of the chemistry of $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ by different group of scientists has resulted in understanding of the applicability of such an efficient and environment friendly reagent and catalyst in diverse range of organic reactions. By virtue of its low toxicity and versatility, it holds considerable promise as a reagent and catalyst. It offers the added advantage of being inexpensive and easy to handle. The results described in this review lead us to hope that there will be new and novel applications of the reagent in the near future.

Acknowledgements

The authors thank CSIR, New Delhi for financial help and the Director, CSIR-NEIST, Jorhat, Assam, India, for his support.

References

1. (a) Thomas, C. A.; Moshier, M. B.; Morris, H. E.; Moshier, R. W. *Anhydrous Aluminum Chloride in Organic Chemistry*, Reinhold: New York, **1941**. (b) Roberts, R. M.; Khalaf, A. A. *Friedel-Crafts Alkylation Chemistry, A Century of Discovery*; Marcel Dekker: New York, **1984**. (c) Greenwood, N. N.; Earnshaw, A. *Chemistry of the elements*; Pergamon Press: Oxford, **1984**; p 254.
2. Dutta, D. K.; Konwar, D. *Indian J. Chem.* **1994**, *33B*, 690-691.
3. For review see: Welsh, D. A. *Synthesis* **1980**, 677-688.
<http://dx.doi.org/10.1055/s-1980-29169>
4. (a) Trimble, M. R.; Hindmarch, I. *Benzodiazepines*, Taylor & Francis, **2000**. (b) Sternbach, L. H. *Angew. Chem. Int. Ed.* **1971**, *10*, 34-43.
<http://dx.doi.org/10.1002/anie.197100341>
5. Liou, J. -P.; Chang, C. -W.; Song, J. -S.; Yang, Y. -N.; Yeh, C. -F.; Tseng, H. -Y.; Lo, Y. -K.; Chang, Y. -L.; Chang, C. -M.; Hsieh, H. -P. *J. Med. Chem.* **2002**, *45*, 2556-2562.
<http://dx.doi.org/10.1021/jm010365+>
6. Boruah, M.; Konwar, D. *J. Chem. Research (S)* **2000**, 232-233.
7. (a) Shibahara, S.; Kondo, S.; Maeda, K.; Umezawa, H.; Ohno, M. *J. Am. Chem. Soc.* **1972**, *94*, 4353-4354.
<http://dx.doi.org/10.1021/ja00767a059>
(b) Hashiguchi, S.; Kawada, A.; Natsugari, H. *J. Chem. Soc., Perkin Trans. 1*, **1991**, 2435-2444.

- (c) Sham, H. L.; Zhao, C.; Li, L.; Betebenner, D. A.; Saldivar, A.; Vasavanonda, S.; Kempf, D. J.; Plattner, J. J.; Norbeck, D. W. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 3101-3103.
[http://dx.doi.org/10.1016/S0960-894X\(02\)00643-1](http://dx.doi.org/10.1016/S0960-894X(02)00643-1)
8. (a) Hulst, R.; Heres, H.; Peper, N. C. M. W.; Kellogg, R. M. *Tetrahedron: Asymm.* **1996**, *7*, 1373-1384.
[http://dx.doi.org/10.1016/0957-4166\(96\)00154-1](http://dx.doi.org/10.1016/0957-4166(96)00154-1)
(b) Li, X.; Yeung, C.-H.; Chan, A. S. C.; Yang, T.-K. *Tetrahedron: Asymm.* **1999**, *10*, 759-763.
[http://dx.doi.org/10.1016/S0957-4166\(99\)00043-9](http://dx.doi.org/10.1016/S0957-4166(99)00043-9)
(c) Evans, P. A.; Holmes, A. B.; McGeary, R. P.; Nadin, A.; Russell, K.; O'Hanlon, P. J.; Pearson, N. D. *J. Chem. Soc., Perkin Trans. 1* **1996**, 123-138.
9. Chowdhury, P. K.; Borah, P. *J. Chem. Research (S)* **1994**, 230-231.
10. Martin, E. L. *Org. React.* **1942**, *1*, 155-209. (b) Vedejs, E. *Org. React.* **1975**, *22*, 401-422.
11. Todd, D. *Org. React.* **1948**, *4*, 378-422.
12. Konwar, D.; Dutta, D. K.; Sandhu, J. S. *Indian J. Chem.* **1995**, *34B*, 725-727.
13. Dutta, D. K.; Konwar, D. *J. Chem. Research (S)* **1998**, 266-267.
14. Boruah, M.; Konwar, D. *Synlett* **2001**, 795-796.
<http://dx.doi.org/10.1055/s-2001-14617>
15. Bartoli, G.; Antonio, G. D.; Giovannini, R.; Giuli, S.; Lanari, S.; Paoletti, M.; Marcantoni, E. *J. Org. Chem.* **2008**, *73*, 1919-1924, and references cited therein.
<http://dx.doi.org/10.1021/jo7024288>
16. Li, C. B.; Zheng, P. W.; Zhao, Z. X.; Zhang, W. Q.; Li, M. B.; Yang, Q. C.; Cui, Y.; Xu, Y. L. *Chin. Chem. Lett.* **2003**, *14*, 773-775.
17. Zheng, P. -W.; Duan, X. -M.; Wang, W.; Wang, Z. -J.; Guo, Y. -P.; Tu, Q. -D. *Chin. J. Chem.* **2006**, *24*, 825-826.
<http://dx.doi.org/10.1002/cjoc.200690157>
18. Boruah, M.; Konwar, D. *J. Org. Chem.* **2002**, *67*, 7138-7139.
<http://dx.doi.org/10.1021/jo020005+>
19. Namboodiri, V. V.; Varma, R. S. *Tetrahedron Lett.* **2002**, *43*, 1143-1146.
[http://dx.doi.org/10.1016/S0040-4039\(01\)02372-3](http://dx.doi.org/10.1016/S0040-4039(01)02372-3)
20. Gogoi, P.; Konwar, D.; Das Sharma, S.; Gogoi, P. K. *Synth. Commun.* **2006**, *36*, 1259-1264.
<http://dx.doi.org/10.1080/00397910500516316>
21. Das Sharma, S.; Hazarika, P.; Konwar, D. *Catal. Commun.* **2008**, *9*, 709-714.
<http://dx.doi.org/10.1016/j.catcom.2007.08.008>
22. Chang, S. U.; Cho, J. H.; Lee, J. C. *Bull. Korean Chem. Soc.* **2008**, *29*, 27-28.
<http://dx.doi.org/10.5012/bkcs.2008.29.1.027>
23. Nakamura, R.; Komura, K.; Sugi, Y. *Catal. Commun.* **2008**, *9*, 511-515.
<http://dx.doi.org/10.1016/j.catcom.2007.03.024>
24. Namboodiri, V. V.; Varma, R. S. *Chem. Commun.* **2002**, 342-343.
<http://dx.doi.org/10.1039/b110565e>

Authors' Biographies



Dr. Saikat Das Sharma was born in Agartala, Tripura, India. He received his B.Sc. (Hons) in Chemistry from M. B. B. College, Agartala and M.Sc. in Organic Chemistry from Tripura University, Agartala. In 2004, he joined the group of Dr. Dilip Konwar where he obtained his PhD working on the development of green methods for the synthesis of bioactive molecules. Presently, he is an Assistant Professor in Dasaratha Deb Memorial College, Khowai, Tripura, India. His research interests include organic reactions in aqueous media, transition metal catalyzed reactions and synthesis of bio-active heterocycles.



Dr. Pallab Pahari was born in Midnapur, West Bengal, India. He obtained his PhD in 2008 from the Indian Institute of Technology Kharagpur, India, under the supervision of Prof. Dipakranjan Mal. During the year 2007 to 2011, he was a post-doctoral fellow in the lab of Prof. Jürgen Rohr, University of Kentucky, USA. Currently he is a scientist at CSIR-North East Institute of Science and Technology, Assam, India. His research interests include development of green chemical methodologies, synthesis of bioactive heterocycles, and isolation of bioactive natural products.



Dr. Swapnali Hazarika was born in 1973 at Jorhat, Assam, India. She obtained her PhD from Dibrugarh University under the guidance of Dr N N Dutta, Chief Scientist, CSIR-NEIST at 2002. In 2005, she joined CSIR-North East Institute of Science and Technology and currently holding the position of Scientist. She has been conducting research in the development of Separation and Purification Technology as well as Reaction Engineering. She has published one book and more than 30 research papers in National and International journal of repute.



Dr. Parasa Hazarika is a native of Jorhat, Assam, India. She received her master's from Gauhati University, Assam, India, and Ph.D. from Dibrugarh University, Assam, India, working in the lab of Dr. Dilip Konwar. Currently, she is an Assistant Professor in the Department of Chemistry, Jorhat Institute of Science & Technology, Assam. Her research is centered on the study of the green synthetic reagent systems for the synthesis of biologically active compounds and application of nanomaterials as green catalyst for the synthesis of important drug intermediates.



Manash Jyoti Borah was born in 1987 at Jorhat, Assam, India. He completed his B.Sc from Dibrugarh University in 2009. In 2010, he joined CSIR-NEIST and currently working in the lab of Dr. Dilip Konwar as a Research Assistant. His research is centered on the area of development of green chemical and enzymatic methodologies for the synthesis of bioactive heterocyclic molecules.



Dr. Dilip Konwar was born in 1956 at Sibsagar, Assam, India. He completed his PhD in 1987 from Dibrugarh University, Assam. He worked as post-doctoral fellow in the lab of Prof. K. T. Potts, Rensselaer Polytechnic Institute, Troy, NY, USA and Prof. J. D. Woolins, University of St. Andrews, UK. From 1982 he is working as scientist at CSIR-North East Institute of Science and Technology and currently holding the position of Chief Scientist. His research interests include green chemistry, synthesis of heterocyclic compounds and organometallic chemistry. He has published more than 60 research papers and holding 15 patents.