

One-step synthesis of furan-2,5-dicarboxylic acid from furan-2-carboxylic acid using carbon dioxide

Róbert Fischer* and Mária Fišerová

Pulp and Paper Research Institute, Lamačská cesta 3, 841 04 Bratislava, Slovak Republic

E-mail: robofis@centrum.sk

DOI: <http://dx.doi.org/10.3998/ark.5550190.p008.285>

Abstract

A short and effective method for preparation and isolation of furan-2,5-dicarboxylic acid has been developed. Furan-2,5-dicarboxylic acid was prepared directly from furan-2-carboxylic acid in one step under strong basic conditions using carbon dioxide. This method allows to obtain the product with high added value, furan-2,5-dicarboxylic acid, from natural sources such as hemicelluloses containing pentoses.

Keywords: Furan-2,5-dicarboxylic acid, furan-2-carboxylic acids, high added value chemicals, hemicelluloses

Introduction

Interest in renewable sources of energy and chemicals is increasing because of expected worldwide shortages of easy-to-access oil and gas and growing concern regarding accumulation of greenhouse gases in the atmosphere. The concept of an Integrated Forest Products Biorefinery has been developed by a number of investigators. The idea is to convert biomass into bio-energy and chemicals.¹ Integrated biorefinery is a process for extracting carbohydrates, oils, lignin, and other materials from biomass and converting them into fuels, high added value chemicals and other materials with almost no waste.² Pulp and paper mills that produce multiple products from biomass can be categorized as biorefineries. Approximately 20% of the dry wood weight are hemicelluloses which are dissolved in the black liquor and are afterwards burned in the chemical recovery systems. Pre-extraction of hemicelluloses prior to kraft pulping and further conversion to high added value chemicals offer an interesting economic opportunity for pulp and paper industry.³

Furan-2,5-dicarboxylic acid (**1**, FDCA, Figure 1) is a very important chemical substance due to its application in polymerization. It compensates the building blocks in the production of

polymers which contain an aromatic moiety. It is also valuable in pharmacology.⁴⁻⁶ Moreover, furan-2,5-dicarboxylic framework is a current versatile building block for the synthesis of interesting organic compounds and biologically active natural products.^{7,8} The production of FDCA **1** is usually based on the conversion of hexoses under strong acid conditions at high temperatures^{9,10} or on various oxidative processes of 2,5-disubstituted furans, mainly 5-(hydroxymethyl)furfural and 5-(hydroxymethyl)furoic acid as C₆-substances, compounds containing six carbon atoms mostly derived from biomasses.¹¹⁻¹³ The mostly investigated method in laboratory involves conversion of furan-2-carboxylic acid methyl ester *via* chloromethylation - exchange of chloro-substituent with hydroxyl group - oxidation process⁷ along with earlier works dealing with the synthesis of furan-2,5-dicarboxylic species.^{14,15}

The last named procedure represents multistep synthesis providing FDCA **1** and its derivatives in low total yield. On the other hand, this allows the possibility to obtain high added value product FDCA **1** from C₅-compounds that can be manufactured from hemicelluloses containing mostly various pentoses¹⁶ using furan-2-carboxylic acid (**2**) as intermediate (Figure 1). This strategy has been published for the first time by Fu, Y. and co-workers, who reported a new pathway of converting furfural, a bulk biomass-based chemical, into FDCA including catalytic disproportionation of furoate to furan and 2,5-furandicarboxylate catalyzed by zinc chloride.¹⁷





Figure 1. Retrosynthetic analysis of FDCA **1** from C₅-compounds.

Results and Discussion

The addition of the organo-metallic compounds or carbanions generated directly from substituted furans on carbon dioxide is considered to be universal synthetic route for the preparation of a wide range of furan carboxylic acids.¹⁸⁻²¹ Furan ring deprotonisation of furan-2-carboxylic acid derivatives with strong bases such as *n*-butyllithium or LDA (lithiumdiisopropylamide) has been also published.^{22,23} The first work concerning the generation of dianions directly from furan-2-carboxylic acid (**2**) was published by Knight and Nott.²⁴ The authors demonstrated the formation of dianion species by regioselective deprotonisation of acid **2** in the position H-5 with LDA and its reactivity with D₂O and other more or less reactive electrophiles, except CO₂. On the other hand, Carpenter and Chadwick investigated the course of lithiation of acid **2**, using *n*-butyllithium and found that the regioselectivity is strongly dependent on the type of the lithium base.²⁵ In comparison with LDA, the use of *n*-butyllithium led to the mixture of 3- and 5-

dicarboxylic acid (**3**) in ratio 50:50. Furthermore, we observed the inversion of regioselectivity in favour of 2,3-regioisomer in comparison with the results reported by Knight and Nott²⁴ by addition of methyl iodide (instead CO₂) retaining the mixture one hour at -78 °C. The reaction also provided the mixture of two regioisomers 5-methylfuran-2-carboxylic acid (**4**) and 3-methylfuran-2-carboxylic acid (**5**) in ratio 35:65 in the yield of 83%. All obtained results are summarized in Table 1.

Because it is difficult to isolate the carboxylic acids **1** and **3** we decided to convert them into their methyl esters **6**, **7** (Scheme 2). Conventional esterification catalyzed by toluene-4-sulfonic acid failed due to low solubility of initial acids. Furthermore, earlier work provided dimethyl ester **6** in lower yields of 68% and 59%, respectively.¹⁰ Therefore, we carried out the reaction in the presence of trimethylsilyl chloride (TMSCl, 2 eq.) in neat methanol under reflux (Scheme 2). This method was already applied in the esterification of furan-2,3-dicarboxylic acid **3**.²⁶ Now we have used it in the case of furan-2,5-dicarboxylic acid **1** for the first time. All obtained results are summarized in Table 2.



Scheme 2. Esterification of furandicarboxylic acids using of TMSCl.

Table 2. Preparation of furandicarboxylic acid dimethyl esters

Dicarboxylic acid	Methyl ester	Yield	mp
		%	°C
1 (purchased from Aldrich)	6	82	111-112 ^e
1 (solid, Table 1, Entry 1)	6	55 ^c	110-111 ^e
1, 3 (crude product, Table 1, Entry 2) ^a	6, 7 ^b	52 ^{c,d}	---

^aRatio **1:3** = 50:50.

^bRatio **6:7** = 50:50.

^cYield calculated after two steps from furan-2-carboxylic acid.

^dEsters were isolated as the mixture of both regioisomers.

^emp 109-110 °C.²⁷

Unfortunately, we were not able to separate the dimethyl esters **6** and **7** from each other by column chromatography on silica gel.

Conclusions

We have successfully prepared furan-2,5-dicarboxylic acid (**1**, FDCA) from furan-2-carboxylic acid (**2**) by reaction with LDA in THF at -78 °C using carbon dioxide as electrophile. This reaction proceeded regioselectively and provided FDCA **1** in 73% yield and in satisfactory purity. FDCA **1** was prepared from furan-2-carboxylic acid (**2**) as representative C₅-building block readily available from D-xylose. In addition, we have converted FDCA **1** into furan-2,5-dicarboxylic acid dimethyl ester (**6**) by reaction in neat methanol in the presence of TMSCl with the yield of 82%. Though FDCA is commercially available, the presented method could be an alternative way of its preparation or relative derivatives with high added value from hemicelluloses rich in pentoses. Also, treatment of lithium 5-lithiofuran-2-carboxylate with other reactive electrophiles might be examined in order to prepare interesting building blocks with different carbonyl species in the C-2 and C-5 positions of furan ring. Further studies and experiments are currently in progress.

Experimental Section

General. All melting points were measured on a Melting Point B-540 apparatus (Büchi, Switzerland) and are uncorrected. NMR spectra were recorded on a Varian VRX-300 spectrometer (¹H, 300 MHz Varian, USA) in CDCl₃ and DMSO-*d*₆ using tetramethylsilane (TMS) as internal standard. TLC analyses were carried out using TLC Silica gel 60 F₂₅₄ (aluminium sheets, Merck, Germany) and visualized by UV light or by permanganate solution. Column chromatographies were performed on Büchi system (Pump Manager C-615 and Fraction Collector C-660, Büchi, Switzerland) using Normasil 60 silica gel (0.040-0.063 mm) (VWR, Germany). All solvents were dried and distilled according to conventional methods. All reagents were purchased from Aldrich (Germany), Acros Organics (Germany), Alfa-Aesar (Germany), Merck (Germany) and Mikrochem Trade (Slovakia) and were used without further purification.

Furan-2,5-dicarboxylic acid (1). A well-dried reaction flask sealed with rubber septum was flushed with argon and charged with diisopropylamine (1.06 mL, 7.5 mmol, stored over molecular sieves). *n*-Butyllithium (4.70 mL, 7.5 mmol, 1.6 M solution in hexane) was added dropwise at -10 °C over 10 min. under argon balloon. The resulting viscous oil stayed at this temperature for 25 min. Afterwards, the reaction flask was cooled to -78 °C and *in situ* prepared LDA was dissolved in dry THF (12 mL). A solution of furan-2-carboxylic acid (**2**, 400 mg, 3.57 mmol) in dry THF (6 mL) was added dropwise over a period of 20 min. When the solution was added the reaction mixture was stirred for one hour. Subsequently, dry CO₂ was bubbled into the solution for 15 min at -78 °C and next 15 min at 0 °C. Then the resulting slurry was diluted with saturated aq. ammonium chloride (25 mL). Water was added (10 mL) and the final liquid phase

was extracted with ether (25 mL). Water layer was separated, cooled with ice-bath and acidified with 2N HCl (until pH 1). The precipitated solid was collected and stirred at 80 °C in 2 N HCl for two hours. The final furan-2,5-dicarboxylic acid (**1**) was collected by filtration and well dried. Colourless crystals, yield 73%, 407 mg; ¹H NMR (300 MHz, DMSO-d₆): δ_H 7.28 (2H, s, H-3, H-4). The NMR spectra correspond to the literature data.²⁸

Furan-2,5-dicarboxylic acid dimethyl ester (6). A reaction flask was charged with furan-2,5-dicarboxylic acid (**1**, 250 mg, 1.55 mmol, 97%, purchased from Aldrich), equipped with condenser and flushed with argon. Methanol (0.8 mL) was added, followed by trimethylsilyl chloride (0.4 mL, 3.10 mmol). The reaction mixture was stirred for 2 hours under reflux. After removal of the solvent *in vacuo*, product was isolated by flash column chromatography on silica gel (CH₂Cl₂). Colourless crystals, yield 82%, 234 mg, mp 111-112 °C; ¹H NMR (300 MHz, CDCl₃): δ_H 3.94 (6H, s, COOCH₃), 7.23 (2H, s, H-3, H-4). The NMR spectra correspond to the literature data.²⁷

Additional ¹H NMR data (300 MHz, DMSO-d₆) for furan-2,3-dicarboxylic acid (**3**), 5-methylfuran-2-carboxylic acid (**4**), 3-methylfuran-2-carboxylic acid (**5**) and furan-2,3-dicarboxylic acid dimethyl ester (**7**) were obtained from the corresponding mixtures of both regioisomers:

Furan-2,3-dicarboxylic acid (3): δ_H 6.87 (1H, d, H-4, *J* 1.8 Hz), 7.90 (1H, d, H-5, *J* 1.8 Hz). The NMR spectra correspond to the literature data.³¹

5-Methylfuran-2-carboxylic acid (4): δ_H 2.33 (3H, s, CH₃), 6.27-6.28 (1H, m, H-4), 7.10 (1H, d, H-3, *J* 3.5 Hz), 12.86 (1H, bs, COOH). The NMR spectra correspond to the literature data.²⁹

3-Methylfuran-2-carboxylic acid (5): δ_H 2.27 (3H, s, CH₃), 6.53 (1H, d, H-4, *J* 1.8 Hz), 7.75 (1H, d, H-5, *J* 1.8 Hz), 12.86 (1H, bs, COOH). The NMR spectra correspond to the literature data.³⁰

Furan-2,3-dicarboxylic acid dimethyl ester (7): δ_H 3.91 (3H, s, COOCH₃), 3.95 (3H, s, COOCH₃), 6.79 (1H, d, H-4, *J* 1.8 Hz), 7.53 (1H, d, H-5, *J* 1.8 Hz). The NMR spectra correspond to the literature data.²⁶

Acknowledgements

This work was supported by the Slovak Research and Development Agency under contract No. APVV-0367-10.

References

1. Mao, H.; Genco, J. M.; Yoon, S.-H.; van Heiningen, A.; Pendse, H. *J. Biobased Mater. Bio.* **2008**, *2*, 177.
<http://dx.doi.org/10.1166/jbmb.2008.309>

2. Carvalheiro, F.; Duarte, L. C.; Gírio, F. M. *J. Sci. Ind. Res.* **2008**, *67*, 849.
3. van Heiningen, A. *Pulp and Paper Canada* **2006**, *107*, 38.
4. Gandini, A.; Belgacem, M. N. *Prog. Polym. Sci.* **1997**, *22*, 1203.
[http://dx.doi.org/10.1016/S0079-6700\(97\)00004-X](http://dx.doi.org/10.1016/S0079-6700(97)00004-X)
5. Lewkowski, J. *Arkivoc* **2001**, (i), 17.
6. Bozell, J. J.; Petersen, G. R. *Green Chem.* **2010**, *12*, 539.
<http://dx.doi.org/10.1039/b922014c>
7. Schmuck, C.; Machon, U. *Eur. J. Org. Chem.* **2006**, 4385.
<http://dx.doi.org/10.1002/ejoc.200600324>
8. Yin, B.-L.; Yang, Z.-M.; Hu, T.-S.; Wu, Y.-L. *Synthesis* **2003**, 1995.
<http://dx.doi.org/10.1055/s-2003-41042>
9. Taguchi, Y.; Oishi, A.; Iida, H. *Chem. Lett.* **2008**, *37*, 50. DOI: 10.1246/cl.2008.50.
<http://dx.doi.org/10.1246/cl.2008.50>
10. Lewkowski, J. *Pol. J. Chem.* **2001**, *75*, 1943.
11. Koopman, F.; Wierckx, N.; de Winde, J. H.; Ruijssenaars, H. J. *Bioresour. Technol.* **2010**, *107*, 6291.
<http://dx.doi.org/10.1016/j.biortech.2010.03.050>
12. Partenheimer, W.; Grushin, V. V. *Adv. Synth. Catal.* **2001**, *343*, 102.
[http://dx.doi.org/10.1002/1615-4169\(20010129\)343:1<102::AID-ADSC102>3.0.CO;2-Q](http://dx.doi.org/10.1002/1615-4169(20010129)343:1<102::AID-ADSC102>3.0.CO;2-Q)
13. Brasholz, M.; von Känel, K.; Hornung, C. H.; Saubern, S.; Tsanaktsidis, J. *Green Chem.* **2011**, *13*, 1114.
<http://dx.doi.org/10.1039/c1gc15107j>
14. Moldenhauer, O.; Trautmann, G.; Irion, W.; Pfluger, R.; Döser, H.; Mastaglio, D.; Marwitz, H. *Liebigs Ann. Chem.* **1953**, *580*, 169.
<http://dx.doi.org/10.1002/jlac.19535800302>
15. Nielek, S.; Lesiak, T. *J. Prakt. Chem.* **1988**, *330*, 825.
<http://dx.doi.org/10.1002/prac.19883300523>
16. Feather, M. S.; Harris, D. W.; Nichols, S. B. *J. Org. Chem.* **1972**, *37*, 1606.
<http://dx.doi.org/10.1021/jo00975a032>
17. Pan, T.; Deng, J.; Xu, Q.; Zuo, Y.; Guo, Q.-X.; Fu, Y. *ChemSusChem.* **2013**, *6*, 47.
<http://dx.doi.org/10.1002/cssc.201200652>
18. Dawson, M. I.; Chan, R.; Hobbs, P. D.; Chao, W.; Schiff, L. J. *J. Med. Chem.* **1983**, *26*, 1282.
<http://dx.doi.org/10.1021/jm00363a013>
19. Carpenter, A. J.; Chadwick, D. J. *Tetrahedron* **1985**, *41*, 3803.
[http://dx.doi.org/10.1016/S0040-4020\(01\)91401-2](http://dx.doi.org/10.1016/S0040-4020(01)91401-2)
20. Magnus, P.; Diorazio, L.; Donohoe, T. J.; Giles, M.; Pye, P.; Tarrant, J.; Thom, S. *Tetrahedron* **1996**, *52*, 14147.
[http://dx.doi.org/10.1016/0040-4020\(96\)00865-4](http://dx.doi.org/10.1016/0040-4020(96)00865-4)

21. Takahashi, M.; Kuroda, T.; Ogiku, T.; Ohmizu, H.; Kondo, K.; Iwasaki, T. *Heterocycles* **1993**, *36*, 1867.
<http://dx.doi.org/10.3987/COM-93-6394>
22. Chadwick, D. J.; McKnight, M. V.; Ngochindo, R. *J. Chem. Soc., Perkin Trans. 1* **1982**, 1343.
<http://dx.doi.org/10.1039/p19820001343>
23. Carpenter, A. J.; Chadwick, D. J. *Tetrahedron Lett.* **1985**, *26*, 5335.
[http://dx.doi.org/10.1016/S0040-4039\(00\)95032-9](http://dx.doi.org/10.1016/S0040-4039(00)95032-9)
24. Knight, D. W.; Nott, A. P. *J. Chem. Soc., Perkin Trans. 1* **1981**, 1125.
<http://dx.doi.org/10.1039/p19810001125>
25. Carpenter, A. J.; Chadwick, D. J. *Tetrahedron Lett.* **1985**, *26*, 1777.
[http://dx.doi.org/10.1016/S0040-4039\(00\)98336-9](http://dx.doi.org/10.1016/S0040-4039(00)98336-9)
26. Klein, M.; Sandner, P.; Frey, R.; Riedl, B.; Christensen, O. WO 2007/118602 (A1), Bayer Healthcare AG.
27. Barton, D. H. R.; Brown, B. D.; Ridley, D. D.; Widdowson, D. A.; Keys, A. J.; Leaver, C. J. *J. Chem. Soc., Perkin Trans. 1* **1975**, 2069.
<http://dx.doi.org/10.1039/p19750002069>
28. Yoon, D.-W.; Gross, D. E.; Lynch, V. M.; Sessler, J. L.; Hay, B. P.; Lee, C.-H. *Angew. Chem. Int. Ed.* **2008**, *47*, 5038.
<http://dx.doi.org/10.1002/anie.200801426>
29. Collins, M. J.; Hatton, P. M.; Sternhell, S. *Aust. J. Chem.* **1992**, *45*, 1119.
<http://dx.doi.org/10.1071/CH9921119>
30. Cahiez, G.; Chavant, P.-Y.; Metais, E. *Tetrahedron Lett.* **1992**, *33*, 5245.
<http://dx.doi.org/10.1016/j.bbr.2011.03.031>
31. Dumas, J. P.; Boyer, S. J.; Dixon, J. A.; Joe, T. K.; Kluender, H. C. E.; Lee, W.; Nagarathnam, D.; Sibley, R. N.; Su, N. US 6689883(B1), 2004, Bayer Pharmaceuticals Corporation.