

Supplementary Material



Reduction of quinolines to 1,2,3,4-tetrahydroquinolines with hydrosilane/ethanol catalyzed by TiO₂-supported gold nanoparticles under solvent free conditions



Anastasia Louka, Charis Gryparis, and Manolis Stratakis*



Department of Chemistry, University of Crete, Voutes, 71003 Iraklion, Greece



E-mail: stratakis@chemistry.uoc.gr



Dedicated to Professor Michael Orfanopoulos on the occasion of his 67th birthday and retirement, and for his remarkable contribution in physical organic chemistry





^1H and ^{13}C NMR spectra of synthesized quinolines















^1H and ^{13}C NMR spectra of isolated 1,2,3,4-tetrahydroquinolines


































^1H NMR spectrum of co-isolated 2a and intermediate product 2b during the reduction of parent quinoline 2




**^1H NMR spectrum of reduction product 1a in the presence of
 $\text{PhMe}_2\text{SiD}/\text{EtOH}$**



**^1H NMR spectrum of reduction product 1a in the presence of
 $\text{PhMe}_2\text{SiH}/\text{CD}_3\text{OD}$**



Region of the ^1H NMR spectrum of reduction product **8a** in the presence of
PhMe₂SiD/EtOH



^1H NMR and MS spectra of isolated side-product $\text{PhMe}_2\text{SiOCH}_2\text{CH}_3$ 