

Solvent-free one-pot efficient and highly regioselective access to functionalized thiazolopyridones from α -enolic dithioesters

**Anugula Nagaraju, B. Janaki Ramulu, Gaurav Shukla, Abhijeet Srivastava,
Girijesh Kumar Verma, and Maya Shankar Singh***

Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi-221005, India


E-mail: mayashankarbhu@gmail.com


Dedicated to Professor (Dr.) J. S. Yadav on the occasion of his 65th birthday


Contents


¹H & ¹³C NMR Spectra


S2-S27


¹H NMR of 4a


^{13}C NMR of **4a**


^1H NMR of **4b**


^{13}C NMR of **4b**


^1H NMR of **4c**


^{13}C NMR of **4c**


^1H NMR of **4d**


^{13}C NMR of **4d**


^1H NMR of 4e


^{13}C NMR of **4e**


^1H NMR of **4f**


^{13}C NMR of **4f**


^1H NMR of **4g**


^{13}C NMR of **4g**


^1H NMR of **4h**


^{13}C NMR of **4h**


^1H NMR of **4i**


^{13}C NMR of **4i**


^1H NMR of **4j**


^{13}C NMR of **4j**

^1H NMR of **4k**

^{13}C NMR of **4k**

^1H NMR of **41**

^{13}C NMR of **4l**

¹H NMR of **4m**

^{13}C NMR of **4m**