

Supplementary Material

Reaction of 3-aminopyrrole with chloropyrimidines to give pyrroloaminopyrimidines


Michael De Rosa*, David Arnold, Austin Cameron Wright and Young Son


*Department of Chemistry, Penn State Brandywine, 25 Yearsley Mill Road,
Media, Pa 19063, USA*

E-mail: mxdl19@psu.edu


Table of Contents


1. ¹ H NMR (400 MHz, CDCl ₃) (10)	2
2. ¹³ C NMR (100 MHz, CDCl ₃) (10)	3
3. ¹ H NMR (400 MHz, acetone- <i>d</i> ₆) (11)	4
4. ¹³ C NMR (100 MHz, acetone- <i>d</i> ₆) (11)	5
5. ¹ H NMR (400 MHz, acetone- <i>d</i> ₆) (12)	6
6. ¹³ C NMR (100 MHz, acetone- <i>d</i> ₆) (12)	7
7. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) (13)	8
8. ¹³ C NMR (100 MHz, DMSO- <i>d</i> ₆) (13)	9
9. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) (14)	10
10. ¹³ C NMR (100 MHz, DMSO- <i>d</i> ₆) (14)	11
11. ¹ H NMR (400 MHz, acetone- <i>d</i> ₆) (15)	12
12. ¹³ C NMR (100 MHz, acetone- <i>d</i> ₆) (15)	13
13. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) (16)	14
14. ¹³ C NMR (100 MHz, DMSO- <i>d</i> ₆) (16)	15
15. ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) (17)	16
16. ¹³ C NMR (100 MHz, DMSO- <i>d</i> ₆) (17)	17
17. X-ray Structure of 4,6-Dichloro- <i>N</i> -(1 <i>H</i> -pyrrol-3-yl)pyrimidin-2-amine (11)	18





(10) ^{13}C NMR (100 MHz, CDCl_3)


(11) ¹H NMR (400 MHz, acetone-d₆)










(12) ^{13}C NMR (100 MHz, acetone- d_6)




(13) ^{13}C NMR (100 MHz, DMSO-d_6)


(14) ¹H NMR (400 MHz, DMSO-d₆)




(14) ^{13}C NMR (100 MHz, DMSO-d_6)


(15) ¹H NMR (400 MHz, acetone-d₆)




(15) ^{13}C NMR (100 MHz, acetone- d_6)


(16) ^1H NMR (400 MHz, DMSO- d_6)



(16) ^{13}C NMR (100 MHz, DMSO-d_6)

(17) ¹H NMR (400 MHz, DMSO-d₆)



(17) ^{13}C NMR (100 MHz, DMSO- d_6)

Crystall Structure of 4,6-Dichloro-*N*-(1*H*-pyrrol-3-yl)pyrimidin-2-amine (11):

