

Synthesis and cytotoxic activity of new pyrazolo[1,5-*a*]pyrimidines and determination of pyrimidine regio-specific ring formation with 2D NMR

Eman K. A. Abdelall* and John N. Philoppes

Department of Pharmaceutical Organic Chemistry, Faculty of Pharmacy, Beni-Suef University, Beni-Suef 62514, Egypt
E-mail: eman.ahmed@pharm.bsu.edu.eg

DOI: <http://dx.doi.org/10.3998/ark.5550190.p009.743>

Abstract

Novel pyrazolo[1,5-*a*]pyrimidines (**9a**, **9b**, and **10a-c**) were synthesized in high and efficient yields. Their pathway involves the formation of *N,S*-ketene derivatives (**7a** and **7b**) that reacted with hydrazine hydrate to get the key intermediate aminopyrazolo derivatives (**8a** and **8b**). These aminopyrazoles were further reacted with either acetylacetone or β -ketoesters resulting in the targeted pyrazolopyrimidines (**9a**, **9b**, and **10a-c**). All prepared compounds were fully characterized by spectral methods and the cyclization of **10a-d** was proved by 2D NMR such as HMBC, HSQC and NOESY. The targeted pyrazolopyrimidines were subjected to *in vitro* anticancer screening and all of them showed promising cytotoxic activity when compared to doxorubicin. Compound **10d** was the most active with IC_{50} = 1.98, 2.20 and 2.61 μ M against MCF-7, BT474 and A549 cancer cell lines.

Keywords: Benzothiazole; 2D NMR; pyrazolopyrimidines; anticancer; MCF-7

Introduction

The development of new anticancer agents was a major area of research during the last decade. Instead of that complete over control of cancer has not been achieved yet.¹ Beside to surgery, chemotherapy is still one of the most important medical options. Reasonably, many researches are directed to develop a new anti-cancer agent that targeting cancer cells and also with a low toxicity to normal cells.²⁻⁴ In addition, pyrazolo-pyrimidine base scaffolds are belonging to a biologically active class as they are structurally related to intrinsic purine bases. Obviously, several pyrazolopyrimidine models as compounds **1-3** (Fig.1) were reported to possess anti-cancer and cytotoxic activity.⁵⁻⁷ Compound **2** was reported to be an effective anti-tumour agent and a scaffold to adenosine (ATP) binding receptor in several kinases.⁶ Moreover, various

synthetic methods of pyrazolopyrimidines have been described,⁷⁻¹¹ and one of these methods involves the formation of ketene-*S,S* or *N,S*- acetals. Such ketenes reacted with a bi-nucleophilic reagent such as hydrazine or substituted hydrazine to get the key intermediates 5-amino-pyrazoles.⁷




Figure1. Showing different potent anti-cancers: substituted pyrazolopyrimidines (**1-3**), and 2-(3,4-dimethoxyphenyl)-5-fluorobenzothiazole (**PMX610**) (**4**) [4-(1,3-benzothiazol-2-yl)phenyl]amine (**5b**) as anticancer agent and the design of new pyrazolopyrimidine - benzothiazole or oxazoles drug hybrid drugs.


Such intermediates are versatile reagents for building more substituted pyrazolopyrimidines by further condensation with either diketones or β -ketoesters. In addition, several benzothiazole derivatives as [4-(1,3-benzothiazol-2-yl)phenyl]amine (**CJM126**)(**5b**) and 2-

(3,4-dimethoxyphenyl)-5-fluorobenzothiazole (**PMX610**)¹²⁻¹⁵ are famous with their potency that enable them to be effective anti-cancer agents. Based on these findings, and as a continuation of previous work,¹⁶⁻²⁰ we synthesized new expected drug hybrid of two active moieties pyrazolopyrimidine and benzothiazole or oxazole. The main objective of the research is to synthesize two series (**9a,b**) and (**10a-d**) in a convenient method as a new combination of pyrazolopyrimidine based scaffold substituted with benzothiazole or oxazole. This method starts by converting of either 4-aminophenyl-benzothiazole or oxazole to their corresponding 5-aminopyrazole derivatives as a common precursor to pyrazolopyrimidines when reacted with diketones or β -ketoesters. These series of hybrids might be of more synergistic activity against **MCF-7**, **BT474** and **A549** cancer cell lines than the initial entities. Also, one of research objectives is directed toward exploring the mechanism of pyrazolopyrimidine cyclization through reaction of 5-aminopyrazole and β -ketoesters.

Results and Discussion

The reaction of amines **5a** or **5b**¹⁴ with ethyl cyanoacetate in DMF gave 2-cyano-*N*-substituted acetamido derivatives **6a** and **6b**. The structure of **6a** was confirmed by its ¹H NMR that showed (CH₂C≡N) at δ 3.97. **6a** and **6b** were further reacted with phenylisothiocyanate and methyl iodide in the presence of KOH giving the *N*, *S*-ketene acetal derivatives **7a** and **7b**.


Structure **7a** was elucidated by its IR spectrum that showed two NH peaks at 3347 and 3291 cm⁻¹ as well as a C≡N band at 2179 cm⁻¹ and also its ¹H NMR spectrum that exhibited a singlet at δ 2.28 ppm (SCH₃) and two D₂O exchangeable peaks at δ 9.97 and 11.55 corresponding to (NHCO) and (NH phenyl) respectively. Also, ¹³C NMR of **7a** confirm the appearance of signals for (S-CH₃), β -ketene and α -ketene at δ 16.89, 74.35, 167.91 in sequence. These keteneacetals **7a** and **7b** were cyclized upon reaction with hydrazine in ethanol resulting in **8a** and **8b**. The structure **8** is postulated through its spectral data and subsequent reaction. The ¹H NMR of **8a** showed a singlet D₂O exchangeable peak at δ 6.11 corresponding to (NH₂) group (Scheme1).



Scheme 1. Synthesis of 5-amino-pyrazoles **8a** and **8b**, Reagents and conditions: (a) DMF, 6 h., (b) 2M KOH, r.t, DMF, PhNCS, MeI, 6h., (c) NH₂NH₂, EtOH, 3h.

Then, these 5-aminopyrazoles were cyclized to the target compounds **9a** and **9b** through their reaction with 2,4-pentanedione in boiling acetic acid. With a symmetric diketone, only one compound was formed, the reaction proceeded that either NH or NH₂ reacted with any carbonyl followed by condensation and removal of 2 moles of water giving **9a** and **9b** (Scheme 2).

The structure **9** was elucidated by elemental analysis and spectral data. Its IR revealed the disappearance of the cyano group and ¹H NMR spectrum of **9b** showed singlets at δ2.65, 2.69 for two (CH₃) while pyrimidine H-5 appeared at 6.80-7.00 as a multiplet with H-4 of NH-phenyl (Scheme 2).




Scheme 2. preparation of of dimethylpyrazolo[1,5-*a*]pyrimidine **9a** and **9b**, Reagent and conditions: (a) AcOH, Reflux, 3 h.

Additionally, the 5-aminopyrazoles **8a,b** were reacted with β-ketoesters such as ethyl acetoacetate or ethyl 3-oxo-3-phenylpropanoate resulting in pyrazolopyrimidine compounds **10a-d** (scheme3). The structure **10a-d** was confirmed by elemental analysis, spectral data and MO calculation (Table1).

Theoretical MO calculations of heat of formation and bend energy (energy required to bend all bonds in a molecule that could be a measure of its stability) was done by using Chemdraw ultra and MM2 property revealed that **10a-d** is more stable in form **A** than **B**. The MO calculation of (**10a,A**) showed a heat of formation = -339.40086 Kcal/mol and bend energy = 19.8561 Kcal/mol, while in form **B** a heat of formation and bend energy = -369.39 Kcal/mol and 20.508 Kcal/mol were found respectively. The previous result indicates that form **A** is more favorable than **B**. Practically, for **10a** and due to lack of formation of a single crystal analysis, the structure of **10a** was illustrated using HSQC, HMBC and NOESY techniques (Fig. 2). Upon studying 3D model of **10a** in either form **A** or **B**, form **A** showed a correlation between the

protons of methyl group and both of pyrimidine H, and C-4H of NH-phenyl in a distance within the range that appeared in the NOESY experiment (Fig. 2), while form **B** does not show these correlations. NOESY scan confirmed that **10a** is in form **A** and not **B**.



Scheme 3. Synthesis of substituted pyrazol[1,5-*a*]pyrimidines **10a-d**. Reagent and condition: ethyl acetoacetate or ethyl 3-oxo-3-phenylpropanoate, AcOH, Reflux, 5 h.




Figure 2. 3D representation of 10a in form A and B, NOESY scan of 10a.

Table 1: Molecular orbital (M.O) calculation of heat of formation (HF) and bend energy (BE) for expected **10a-d** (structure A or structure B)

	HF*\BE* of structure A	HF*\BE*of structure B
10a	HF: -339.40086 Kcal/mol BE: 19.8561 Kcal/mol	HF: -369.39 Kcal/mol BE: 20.508 Kcal/mol
10b	H F: - 359.7533 kcal/mol BE: 22.9076 Kcal/mol	H F: -546.65 Kcal/mol BE: 23.1722 Kcal/mol
10c	HF: - 395.6209 kcal/mol BE: 19.5831 Kcal/mol	HF: - 426.6263 kcal/mol BE: 20.107 Kcal/mol
10d	H F: -411.8006 kcal/mol BE: 23.5574 Kcal/mol	H F: -679.98 Kcal/mol BE: 23.5864 Kcal/mol

HF*: heat of formation calculated after energy minimization of structures at 25° using Chemdraw 3D ultra version8 and MM2 property BE*: Bend energy calculated at 25°

Anticancer screening:

Cytotoxic study was operated using MTT(3,4-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium Bromide assay method.²¹ All tested compounds showed moderate to high cytotoxic activity against **MCF-7**, **BT474** and **A549** cell lines. The obtained data are listed Table 2. Most synthesized compounds (**7a**, **8a**, **8b**, **9a**, **9b**, **10a** and **10c**) showed moderate cytotoxic activity when compared with the reference. **7b** is the least potent one that exhibited a cytotoxic activity with IC₅₀ 9.02, 9.87, 8.42 μM against **MCF-7**, **BT474** and **A549** cell lines. Regarding compounds with an oxazole moiety only (**6a** and **8a**) they showed nearly the same moderate activity against all cell lines and they are more potent than (**6b** and **8b**) with a benzothiazole moiety. The sight was directed to the compound of benzoythiazole or oxazole /pyrazolopyrimidine hybrids (**9a**, **9b** and **10a-d**). Their potency was ordered **10d** > **10b** > **9b** > **9a** > **10a** > **10c**. The most potent is **10d** with IC₅₀ 1.98, 2.2, 2.61 μM and its result is close to the reference doxorubicin with IC₅₀ 1.72, 1.81, 1.21 μM against **MCF-7**, **BT474** and **A549** cell lines. **10d** is a pyrazolopyrimidine-benzothiazole hybrid and this reactivity could be related to, 1- high potency of benzothiazole moiety, 2- presence of NH tautomeric OH that increase incidence of hydrogen bond (HB) with proposed receptor active site and 3- high lipophilicity due to phenyl moiety when compared to **10b** which is also pyrazolopyrimidine-benzothiazole hybrid with a CH₃ moiety. In addition **10b** also showed higher potency activity with IC₅₀ = 2.01, 2.36, 2.47 μM against **MCF-7**, **BT474** and **A549** cell lines respectively.

Table 2: IC₅₀ of testing compounds and standard doxorubicin against MCF-7, BT474 and A549

Compound no	IC ₅₀ (μM)		
	MCF-7	BT474	A549
6a	5.87±1.07	6.74±1.03	6.41±1.07
6b	9.02±0.98	9.87±1.05	8.42±1.09
8a	5.21±0.77	6.32±1.09	6.27±1.02
8b	6.54±0.88	6.98±1.09	6.96±0.99
9a	3.42±0.89	4.26±0.77	3.65±0.93
9b	2.25±0.12	2.11±0.87	2.74±0.89
10a	2.73±0.70	3.07±0.14	2.91±0.78
10b	2.01±0.40	2.36±0.44	2.47±0.38
10c	5.47±0.80	5.96±1.05	6.36±0.94
10d	1.98± 0.70	2.20±0.50	2.61±0.45
Doxorubicin	1.72± 0.7	1.81±0.19	1.21±0.33

Several benzothiazole and oxazole compounds (**6a**, **6b**, **7a,7b**, **8a** and **8b**) were prepared and also pyrazolopyrimidine with a benzoxazole (**9a**, **10a** and **10c**) those showed considerable cytotoxic activity while their corresponding pyrazolopyrimidines with a benzothiazole (**9b**, **10b** and **10d**) showed the highest activity. Such pyrazolopyrimidine-benzothiazole combination is the successful one that resulted in **10b** and **10d** as the most active compounds. Structure elucidation of **10a** with 2D and NOE excluded their structural form.

Experimental Section

General. Melting points were determined on an Electrothermal digital melting point apparatus and are uncorrected. IR spectra were recorded on an R 435 spectrophotometer (Middlton, Madison West, WI, USA) and values were reported in cm⁻¹. ¹H-NMR and ¹³C-NMR were carried out on Bruker Advance III 400 MHz spectrophotometer (Bruker BioSpin AG, Fällanden, Switzerland) for ¹H and 100 MHz for ¹³C with BBFO Smart Probe and Bruker 400 AEON Nitrogen-Free Magnet, using TMS as an internal standard and chemical shifts were recorded in ppm on δ scale, Faculty of Pharmacy, Beni Suef University, Egypt. The electron impact (EI) mass spectra were recorded on a Hewlett Packard 5988 spectrometer (Palo Alto, CA, USA), Microanalyses for C, H and N were carried out on Perkin-Elmer 2400 analyzer (Perkin-Elmer, Norwalk, CT, USA) at the Micro analytical unit of Cairo University, Egypt, and all compounds were within ±0.4% of the theoretical values. Thin-layer chromatography (TLC) was performed on Merck (Darmstadt, Germany) TLC aluminium sheets silica gel 60 F254 with detection by UV

quenching at 254 nm to follow the course of reactions and to check the purity of products. All reagents and solvents were purified and dried by standard techniques.

General procedure for synthesis of 6a,b. A solution of equivalent amount of ethyl cyanoacetate (0.56 g, 5 mmol) with the corresponding amines **5a** and **5b**¹⁴ in dimethylformamide (20 mL) was heated under reflux for 6 h. The reaction mixture was cooled and concentrated under vacuum to get the solid product which was crystallized from ethanol/dimethylformamide mixture (1:1).

N-(4-(Benzoxazol-2-ylphenyl)-2-cyanoacetamide (6a). Buff solid; Yield 87%; mp 265-267 °C; IR(KBr, cm⁻¹): 3439 (2NH), 2199 (C≡N), 1656 (C=O); ¹H NMR (DMSO-*d*₆) δ 3.97 (s, 2H, CH₂), 7.39-7.42 (m, 2H, H-5 & H-6 of benzoxazole), 7.75-7.80 (m, 4H, H-4 & H-7 of benzoxazole and H-2&H-6 of amidophenyl), 8.18 (d, *J* 8.4 Hz, 2H, H-3&H-5 of amidophenyl), 10.64 (s, 1H, NHCO, D₂O exchangeable); MS (*m/z*, %): 278 (M+1, 13.60%), 277 (M⁺, 77.30%), 68 (C₃H₂NO⁺, 100.00%); Anal. Calcd for C₁₆H₁₁N₃O₂ (277.28): C, 69.31; H, 4.00; N, 15.15. Found: C, 69.57; H, 3.85; N, 15.28.

N-(4-(Benzothiazol-2-ylphenyl)-2-cyanoacetamide (6b). Buff solid; Yield 82%; mp 235-237 °C; IR (KBr, cm⁻¹): 3406 (2NH), 2292 (C≡N), 1640 (C=O); ¹H NMR (DMSO-*d*₆) δ 3.96 (s, 2H, CH₂), 7.42 (t *J* 7.6 Hz, 1H, H-6-of benzothiazole), 7.53 (t *J* 7.6 Hz, 1H, H-5 benzothiazole), 7.75 (d, *J* 8.4 Hz, 2H, H-2&H-6 of amidophenyl), 8.01-8.13 (m, 4H, H-4&H-7 of benzothiazole, H-3 & H-5 of amidophenyl), 10.60 (s, 1H, NHCO, D₂O exchangeable); Anal. Calcd for C₁₆H₁₁N₃OS (293.34): C, 65.51; H, 3.78; N, 14.32. Found: C, 65.37; H, 3.95; N, 14.29.

General procedure for synthesis of 7a, 7b. A solution of **6a** or **6b** (0.1 mmol), potassium hydroxide (0.11 g, 2 mmol), phenyl isothiocyanate (0.27 g, 2 mmol) in dimethylformamide (20 mL) was stirred for 6 h. To the reaction mixture methyl iodide was added (0.28 g, 2 mmol) and the mixture stirred for 3 h, the reaction mixture was concentrated under vacuum to get the product **7a** or **7b** that crystallized from ethanol/ dimethylformamide mixture (1:1)

N-(4-Benzoxazol-2-ylphenyl)-2-cyano-3-methylsulfanyl-3-phenylaminoacetamide (7a). Yellow solid; Yield 85%; mp 201-203 °C; IR (KBr, cm⁻¹): 3447, 3291 (2NH), 2197 (C≡N), 1629 (C=O); ¹H NMR (DMSO-*d*₆) δ 2.28 (s, 3H, SCH₃), 7.21-7.37 (m, 1H, H-4 of NH-phenyl), 7.39-7.55 (m, 6H, H-2, H-6, H-3, H-5-of NH-phenyl and H-5 & H-6 of benzoxazole), 7.74-7.78 (m, 4H, H-2&H-6 of amidophenyl and H-4&H-7 of benzoxazole), 8.01 (d *J* 7.2, 2H, H-3&H-5 of amidophenyl), 9.97 (s, 1H, NH (D₂O exchangeable), 11.55 (s, 1H, NH (D₂O exchangeable), ¹³C NMR (DMSO-*d*₆) δ 16.89, 74.35, 111.23, 118.90, 120.02, 121.11, 121.56, 123.96, 125.26, 125.67, 126.47, 128.31, 129.72, 139.14, 142.10, 142.33, 150.59, 162.66, 164.5, 167.91; Anal. Calcd for C₂₄H₁₈N₄O₂S (426.49): C, 67.59; H, 4.25; N, 13.14. Found: C, 67.37; H, 4.55; N, 13.38.

N-(4-Benzothiazol-2-ylphenyl)-2-cyano-3-methylsulfanyl-3-phenylaminoacetamide (7b). Red solid; Yield 82%; mp 203-205 °C; IR(KBr, cm⁻¹): 3440, 3290 (2NH), 2195 (C≡N), 1625 (C=O); ¹H NMR (DMSO-*d*₆): δ 2.44 (s, 3H, SCH₃), 7.21-7.23 (m, 1H, H-4 of NH-phenyl), 7.31-

7.48 (m, 6H, H-2, H-6, H-3, H-5 of NH-phenyl and H-5 & H-6 of benzothiazole), 7.68 (d J 8.4, 2H, H-2&H-6 of amidophenyl), 7.96 (d J 8.4, 2H, H-3 & H-5 of amidophenyl), 8.02 (m, 2H, H-4&H-7 of benzothiazole), 9.88 (s, 1H, NH, NHCO (D₂O exchangeable)), 11.57 (s, 1H, NH, NH-phenyl (D₂O exchangeable)), ¹³C NMR (DMSO-*d*₆) δ 16.90, 71.01, 111.25, 121.29, 122.75, 123.06, 124.00, 125.76, 126.50, 127.07, 128.12, 128.31, 129.72, 134.76, 141.66, 146.80, 154.09, 167.42, 168.25, 172.66; MS(*m/z*, %): 443 (M+1, 15.41%), 442 (M⁺, 42.51%), 226 (C₁₃H₁₀N₂S⁺, 100.00 %); Anal. Calcd for C₂₄H₁₈N₄OS₂ (442.56): C, 65.13; H, 4.10; N, 12.66. Found: C, 64.98; H, 4.27; N, 12.45.

General procedure for synthesis of 8a, 8b. A suspension of **7a** or **7b** (5 mmol) in ethanol (20 ml) and hydrazine hydrate (0.16 gm, 5 mmol) was heated under reflux for 3 h, the solid formed on hot was separated and crystallized from ethanol/dimethylformamide mixture (1:1).

5-Amino-3-phenylamino-1H-pyrazol-4-carboxylic acid (4-benzoxazol-2-ylphenyl)-amide (8a). Buff solid; Yield 73%; mp 259-261 °C; IR (KBr, cm⁻¹): 3438-3295 (3NH and NH₂), 1661 (C=O); 1604 (C=N); ¹H NMR (DMSO-*d*₆): δ 6.11 (s, 2H, NH₂ 5-amino-pyrazole (D₂O exchangeable)), 6.76-7.92(m, 1H, H-4 of NH-phenyl), 7.18-7.43 (m, 6H, H-2, H-6, H-3, H-5 of NH-phenyl and H-5 & H-6 of benzoxazole), 7.74-7.77 (m, 4H, H-4 & H-7 of benzoxazole and H-2&H-6 of amidophenyl), 8.13 (d J 8.4, 2H, H-3&H-5 of amidophenyl), 8.53 (s, 1H, NH-phenyl (D₂O exchangeable)), 9.09 (s, NH, NHCO (D₂O exchangeable)), 11.36 (s, 1H, NH, NH-pyrazole (D₂O exchangeable)). MS (*m/z*, %): 411 (M+1, 6.34%), 410 (M⁺, 22.54%), 210 (C₁₃H₁₀N₂O⁺, 100.00%); Anal. Calcd for C₂₃H₁₈N₆O₂ (410.43): C, 67.31; H, 4.42; N, 20.48. Found: C, 67.38; H, 4.65; N, 20.39.

5-Amino-3-phenylamino-1H-pyrazol-4-carboxylic acid (4-benzothiazol-2-ylphenyl)amide (8b). Buff solid; Yield 68%; mp 254-256 °C; IR(KBr, cm⁻¹): 3458-3261 (3NH and NH₂), 1644 (C=O); 1592 (C=N); ¹H NMR (DMSO-*d*₆): δ 6.14 (s, 2H, NH₂, 5-aminopyrazole (D₂O exchangeable)), 6.80 (t J 7.2 Hz, 1H, H-4 of NH-phenyl), 7.22 (t J 7.2 Hz, 2H, H-2&H-6 of NH-phenyl), 7.32 (d J 7.2, 2H, H-3 & H-5 of NH-phenyl), 7.51 (t J 7.2 Hz, 1H, H-6 benzothiazole), 7.55 (t J 7.2 Hz, 1H, H-5 benzothiazole), 7.74 (d J 8.4, 2H, H-2&H-6 of amidophenyl), 8.01-8.08 (m, 3H, H-3 & H-5 of amidophenyl and H-7 of benzothiazole), 8.11(d, J 7.2, 1H, H-4 of benzothiazole), 8.57 (s, 1H, NH-phenyl (D₂O exchangeable), 9.07 (s, 1H, NHCO), 11.39 (s, 1H, NH-pyrazole (D₂O exchangeable)). Anal. Calcd for C₂₃H₁₈N₆OS (426.49): C, 64.77; H, 4.25; N, 19.70. Found: C, 64.98; H, 4.46; N, 19.89.

General procedure for synthesis of (9a and 9b). A mixture of the appropriate aminopyrazole **8a** or **8b** (2 mmol) and acetylacetone (0.2 g, 2 mmol) in glacial acetic acid (15 mL) was boiled under reflux for 3 h. The resulting solid was collected and recrystallized from ethanol/DMF mixture (3:1) to achieve **9a** or **9b**.

N-(4-(Benzo[*d*]oxazol-2-yl)phenyl)-5,7-dimethyl-2-(phenylamino)pyrazolo[1,5-*a*]pyrimidine-3-carboxamide (9a). Yellowish white; Yield 62%; mp 283-285 °C; IR (KBr, cm⁻¹): 3320, 3222 (2NH), 2919 (CH aliphatic), 1669 (C=O); 1595 (C=N); ¹H NMR (DMSO-*d*₆): δ

2.70 (s, 3H, CH₃), 2.77 (s, 3H, CH₃), 7.12 (s, 1H, CH pyrimidine), 7.12-7.16 (m, 1H, H-4 of NH-phenyl), 7.36-7.57 (m, 2H, H-4, H-2&H-6 of NH-phenyl), 7.81 (d *J* 7.6, 2H, H-3&H-5 of NH-phenyl), 8.03-8.14 (m, 8H, H-5 & H-6, H-4 & H-7 of benzoxazole and H-2&H-6, H-3&H-5 of amidophenyl), 9.37 (s, 1H, NHCO (D₂O exchangeable)), 10.49 (s, NH, NH-phenyl (D₂O exchangeable)), Anal. Calcd for C₂₈H₂₂N₆O₂ (474.51): C, 70.87; H, 4.67; N, 17.71. Found: C, 71.11; H, 4.82; N, 17.76.

***N*-(4-(Benzo[*d*]thiazol-2-yl)phenyl)-5,7-dimethyl-2-(phenylamino)pyrazolo[1,5-*a*]pyrimidine-3-carboxamide (9b).** Yellow solid; Yield 63%; mp 251-253 °C; IR (KBr, cm⁻¹): 3312, 3229 (2NH), 2921 (CH aliphatic), 1666 (C=O); 1599 (C=N); ¹H NMR (DMSO-*d*₆): δ 2.65 (s, 3H, CH₃), 2.69 (s, 3H, CH₃), 6.80-7.00 (m, 2H, H-4 of NH-phenyl and CH pyrimidine), 7.37 (t *J* 8.0 Hz, 2H, H-2&H-6 of NH-phenyl), 7.43 (t *J* 7.2 Hz, 1H, H-6 benzothiazole), 7.51-7.53 (t *J* 8.0 Hz, 1H, H-5 benzothiazole), 7.76 (d *J* 7.2 Hz, 2H, H-3 & H-5 of NH-phenyl), 7.99 (d *J* 8.8 Hz, 2H, H-2&H-6 of amidophenyl), 8.01 (d *J* 8.0 Hz, 1H, H-7 of benzothiazole), 8.06-8.10 (m, 3H, H-3&H-5 of amidophenyl and H-4 of benzothiazole), 9.31 (s, 1H, NHCO (D₂O exchangeable)), 10.28 (s, NH, NH-phenyl (D₂O exchangeable)), MS (*m/z*, %) 491 (M+1, 10.39%), 490 (M⁺, 29.97%), 265 (C₁₄H₁₁N₅O⁺, 100.00%); Anal. Calcd for C₂₈H₂₂N₆OS (490.58): C, 68.55; H, 4.52; N, 17.13. Found: C, 68.33; H, 4.38; N, 17.16.

General procedure for synthesis of 10a-d. A mixture of the appropriate aminopyrazole **8a** or **8b** (2 mmol) and ethylacetoacetate or ethyl 3-oxo-3-phenylpropanoate (2 mmol) in glacial acetic acid (15 mL) was boiled under reflux for 5 h. The resulting solid was collected and recrystallized from ethanol/DMF mixture to afford **10a-d**.

***N*-(4-(Benzo[*d*]oxazol-2-yl)phenyl)-4,7-dihydro-5-methyl-7-oxo-2-(phenylamino)pyrazolo[1,5-*a*]pyrimidine-3-carboxamide (10a).** Brown solid; Yield 65%; mp 298-300 °C; IR (KBr, cm⁻¹): 3471, 3287, 3182 (3NH), 2935 (CH aliphatic), 1657, 1622 (2C=O); ¹H NMR (DMSO-*d*₆): δ 2.37 (s, 3H, CH₃), 5.80 (s, 1H, CH Pyrimidine), 7.29 (t *J* 8.2 Hz, 1H, H-4), 7.40-7.42 (m, *J* 4.0 Hz, 2H, H-5&H-6 of benzoxazole and H-2&H-6 of NH-phenyl), 7.71 (d *J* 8.0 Hz, 2H, H-3&H-5 of NH-phenyl), 7.78 (t *J* 4.0 Hz, 2H, H-4 & H-7 of benzoxazole), 7.92 (d *J* 8.2 Hz, 2H, H-2&H-6 of amidophenyl), 8.22 (d *J* 8.2 Hz, 2H, H-3&H-5 of amidophenyl), 8.77 (s, 1H, NH, NH pyrazole (D₂O exchangeable)), (s, 1H, NHCO (D₂O exchangeable)), 12.05 (s, 1H, NH of NH-phenyl (D₂O exchangeable)), ¹³C NMR (DMSO-*d*₆) δ 21.53, 89.00, 98.84, 110.73, 111.25, 117.68, 120.01, 121.03, 121.38, 125.30, 125.70, 128.43, 129.28, 135.40, 141.38, 142.12, 142.84, 150.59, 153.24, 155.31, 162.10, 162.74, 172.50, MS (*m/z*, %) 477 (M+1, 9.94%), 476 (M⁺, 29.04%), 210 (C₁₃H₁₀N₂O⁺, 100.00%); Anal. Calcd for C₂₇H₂₀N₆O₃ (476.49): C, 68.06; H, 4.23; N, 17.74. Found: C, 67.88; H, 3.95; N, 17.88.; HMBC, HSQC and NOESY scan are in supplementary materials.

***N*-(4-(Benzo[*d*]thiazol-2-yl)phenyl)-4,7-dihydro-5-methyl-7-oxo-2-(phenylamino)pyrazolo[1,5-*a*]pyrimidine-3-carboxamide (10b).** Buff solid; Yield 69%; mp 284-286 °C; IR (KBr, cm⁻¹): 3438, 3280, 3120 (3NH), 1674, 1642 (2C=O); ¹H NMR (DMSO-*d*₆): δ 2.37 (s, 3H, CH₃), 5.81 (s, 1H, pyrimidine H), 6.92 (t *J* 7.60 Hz, 1H, H-4 of NH-phenyl), 7.31 (t *J* 7.60 Hz,

2H, H-3& H-5 of NH-phenyl), 7.45 (t J 7.2 Hz, 1H, H-6 benzothiazole), 7.55 (t J 7.2 Hz, 1H, H-5 benzothiazole), 7.70 (d J 7.6 Hz, 2H, H-2& H-6 of NH-phenyl), 7.89(d J 8.40 Hz, 2H, H-2&H-6 of amidophenyl), 8.04 (d J 7.2 Hz, 1H, H-7 benzothiazole), 8.11-8.15 (m, 3H, H-3&H-5 of amidophenyl and H-4 of benzothiazole), 8.77 (s, 1H, NH pyrazole (D₂O exchangeable)), 10.10 (s, 1H, NHCO (D₂O exchangeable)), 12.04 (s, 1H, NH, NH-phenyl (D₂O exchangeable)); MS (m/z ,%) 539 (M+1, 0.10%), 538 (M⁺, 0.12%), 80 (C₃H₂N₃⁺, 100.00%); Anal.Calcd for C₃₂H₂₂N₆O₃ (538.56): C, 71.37; H, 4.12; N, 15.60. Found: C, 71.53; H, 3.90; N, 15.78.

***N*-(4-(Benzo[d]oxazol-2-yl)phenyl)-4,7-dihydro-5-phenyl-7-oxo-2-(phenylamino)pyrazolo [1,5-*a*] pyrimidine-3-carboxamide (10c).** Buff solid; Yield 62%; mp 287-289 °C; IR (KBr, cm⁻¹): 3439, 3288, 3185 (3NH), 2925 (CH aliphatic), 1657, 1624 (2C=O); ¹H NMR (DMSO-*d*₆): δ 6.30 (s, 1H, CH pyrimidine), 6.95 (m, 1H, H-4 of NH-phenyl), 7.33-7.42 (m, 4H, H-3& H-5 of NH-phenyl and H-5&H-6 of benzoxazole), 7.60-7.61 (m, 3H, H-3, H-4 and H-5 pyrimidine phenyl), 7.78-7.79 (d J 6.0 Hz, 4H, H-2&H-6 of NH-phenyl and H-2&H-6 of pyrimidine phenyl), 7.97-8.00 (m, 4H, H-4 & H-7 of benzoxazole and H-2&H-6 of amidophenyl), 8.23 (d J 8.2, 2H, H-3&H-5 of amidophenyl), 8.99 (s, 1H, NH, NHCO(D₂O exchangeable)), 10.55 (s, 1H, NH phenyl (D₂O exchangeable)); ¹³C NMR (DMSO-*d*₆) δ 19.40, 88.75, 98.84, 117.40, 117.70, 121.07, 121.22, 122.74, 123.04, 125.78, 127.11, 128.21, 129.30, 134.75, 136.66, 141.40, 142.31, 153.24, 154.10, 155.31, 162.05, 164.70, 167.50; Anal.Calcd for C₂₇H₂₀N₆O₂S (492.55): C, 65.84; H, 4.09; N, 17.06. Found: C, 66.05; H, 3.85; N, 16.96.

***N*-(4-(benzo[d]thiazol-2-yl)phenyl)-4,7-dihydro-5-phenyl-7-oxo-2-(phenylamino)pyrazolo [1,5-*a*]pyrimidine-3-carboxamide (10d).** Buff solid; Yield 64%; mp 287-289 °C; IR (KBr, cm⁻¹): 3414, 3281, 3180 (3NH), 1666, 1632 (2C=O); ¹H NMR (DMSO-*d*₆): δ 6.31 (s, 1H, pyrimidine H), 6.92 (t J 8.00 Hz, 1H, H-4 of NH-phenyl), 7.34 (t J 7.60 Hz, 2H, H-2& H-6 of NH-phenyl), 7.45 (t J 7.2 Hz, 1H, H-6 benzothiazole), 7.53-7.67 (m, 4H, H-5 benzothiazole and H-3&H-5 of pyrimidine phenyl), 7.62 (d J 7.6 Hz, 2H, H-3& H-5 of NH-phenyl), 7.92 (d J 8.40 Hz, 2H, H-2&H-6 of pyrimidine phenyl), 8.03-8.05 (m, 4H, H-7&H-4 benzothiazole and H-2&H-6 of amidophenyl), 8.15 (d J 8.2 Hz, 2H, H-3&H-5 of amidophenyl) 8.98 (s, 1H, NHCO (D₂O exchangeable)), 10.60 (s, 1H, NH, NH-phenyl (D₂O exchangeable)), pyrazole NH not appeared. Anal.Calcd for C₃₂H₂₂N₆O₂S (554.62): C, 69.30; H, 4.00; N, 15.15. Found: C, 69.43; H, 3.87; N, 14.94.

Anti-tumour activity. Three human cancer cell lines [Human Breast Adenocarcinoma (MCF-7) and BT474 and Non-Small Cell Lung Cancer (A549)] were obtained from the American Type Culture Collection. Cells were maintained in **Dulbecco's Modified Eagle's Medium (DMEM: Gibco, USA) supplemented with 10% fetal bovine serum (Gibco), penicillin/streptomycin (Gibco).** Cells were incubated at 37 °C in a humidified incubator containing 5% CO₂. Cell viability count was assessed using MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) (Sigma-Aldrich, St. Louis, MO, USA) in 6

Acknowledgements

We are grateful to Pharmaceutical Organic Chemistry for kind support during progressing of this research.

References

1. Lesyk, R.; Validzimirska, O.; Holota, S.; Zaprutko, L.; Gazell, A. *Eur. J. Med. Chem.* **2007**, *42*, 641.
<http://dx.doi.org/10.1016/j.ejmech.2006.12.006>
2. Raj, T.; Bhatia, R. K.; Kapur, A.; Sharma, M.; Saxena, A. K.; Ishar, M. P. S. *Eur. J. Med. Chem.* **2010**, *45*, 790.
<http://dx.doi.org/10.1016/j.ejmech.2009.11.001>
3. Drews, J. *Drug discovery* **2000**, *287*, 1960.
4. Staben, S.T.; Heffron, T. P.; Sutherlin, D. P.; Bhat, S. R.; Castanedo, G. M.; Chuckowree, I. S.; Dotson, J.; Folkes, A. J.; Friedman, L. S.; Lee, L.; Lesnick, J. *Bioorg. Med. Chem. Lett.* **2010**, *20*, 6048.
<http://dx.doi.org/10.1016/j.bmcl.2010.08.067>
5. Manetti, D.; Brullo, C.; Magnani, M.; Mosci, F.; Chelli, B.; Crespan, E.; Schenone, S.; Naldini, A.; Bruno, O.; Trincavelli, M. L. *J. Med. Chem.* **2008**, *51*, 1252.
<http://dx.doi.org/10.1021/jm701240c>
6. Radi, M.; Dreassi, E.; Brullo, C.; Crespan, E.; Tintori, C.; Brenardo, V.; Valoti, M.; Zamperini, C.; Daigi, H.; Musumeci, F.; Carraro, F. *J. Med. Chem.* **2011**, *54*, 2610.
<http://dx.doi.org/10.1021/jm1012819>
7. Elgemie, G. H.; Elghandour, A. H.; Elzanti, S. M.; Ahmed, S. A. *J. Chem. Res. S*, 1998, 163-165.
8. Catalano, J. G.; Gaitonde, V.; Beesu, M.; Leivers, A. L.; Shotwell, J. B. *Tetrahedron Lett.* **2015**, *56*, 6077.
<http://dx.doi.org/10.1016/j.tetlet.2015.09.068>
9. Zhao, Y. H.; Li, Y.; Guo, T.; Tang, Z.; Xie, W.; Zhao, G. *Tetrahedron Lett.* **2016**, *57*, 2257.
<http://dx.doi.org/10.1016/j.tetlet.2016.04.037>
10. Shekarrao, K.; Kaishap, P. P.; Gogoi, S.; Gogoi, S.; Boruah, R. C. *Tetrahedron Lett.* **2014**, *55*, 5251.
<http://dx.doi.org/10.1016/j.tetlet.2014.07.119>
11. Saikia, P.; Kaishap, P. P.; Prakash, R.; Shekarrao, K.; Gogoi, S.; Boruah, R. C. *Tetrahedron Lett.* **2014**, *55*, 3896.
<http://dx.doi.org/10.1016/j.tetlet.2014.05.021>
12. Kandeel, M. M.; Abdelall, E. K. A.; Abdelhamid, M. K.; Abdelgwad, M. A.; Philoppes, J. N. *Journal of chemical and pharmaceutical research* **2013**, *5*, 16.

13. Hutchinson, I.; Chua, M.-S.; Browne, H. L.; Trapani, V.; Bradshaw, T. D.; Westwell, A. D.; Stevens, M.F.G. *J. Med. Chem.* **2001**, *44*, 1446.
<http://dx.doi.org/10.1021/jm001104n>
14. Shi, D.-F.; Bradshaw, T. D.; Wrigley, S.; McCall, C.J.; Lelieveld, P.; Fichtner, I.; Stevens, M.F.G. *J. Med. Chem.* **1996**, *39*, 3375.
<http://dx.doi.org/10.1021/jm9600959>
15. Bradshaw, T. D.; Wrigley, S.; Shi, S. D. -F, Schuitz, R. J.; Paull, K. D.; Stevens, M. F. G. *Br. J. Cancer* **1998**, *77*, 745.
<http://dx.doi.org/10.1038/bjc.1998.122>
16. Abdelall E. K. A.; Kamel G. M. *Eur. J. Med. Chem.* **2016**, *118*, 250.
<http://dx.doi.org/10.1016/j.ejmech.2016.04.049>
17. Abdelhamid, A. O.; Abdelall, E. K. A.; Abdel-Riheem N. A.; Ahmed, S. A. *Phosphorus, Sulfur, Silicon and Relat. Elements* **2010**, *185*, 709.
<http://dx.doi.org/10.1080/10426500902922933>
18. Elgemeie, G. H.; El-Enany, M. M.; Ismail, M. M.; Abdelall E. K. A. *Nucleosides, Nucleotides and Nucleic Acids* **2002**, *21*, 477.
<http://dx.doi.org/10.1081/NCN-120014820>
19. Kandeel, M. M.; Kamal, A. M.; Abdelall, E. K. A.; Elshemy, H. A. H. *Eur. J. Med. Chem.* **2013**, *59*, 183.
<http://dx.doi.org/10.1016/j.ejmech.2012.11.011>
20. Kandeel, M. M.; Abdelhameid, M. K.; Abdelall, E. K. A.; Labib, M. B. *Chem. Pharm. Bull.* **2013**, *61*, 637.
<http://dx.doi.org/10.1248/cpb.c13-00089>
21. Abdelall, E. K. A.; Mahmoud, A. M.; Abdelhamid, A. O. *Phosphorus, Sulfur and Silicon and Related Elements* **2010**, *185*, 1862.
<http://dx.doi.org/10.1080/10426500903348013>