

Supplementary Material

Synthesis of new zinc (II) phthalocyanine conjugates with block copolymers for cancer therapy

Łukasz Lamch,^a Marta Tsirigotis-Maniecka,^a Julita Kulbacka,^b Kazimiera A. Wilk^{a*}

Department of Organic and Pharmaceutical Technology, Faculty of Chemistry, Wrocław University of Science and Technology, Wybrzeże Wyspińskiego 27, 50-370 Wrocław, Poland

E-mail: kazimiera.wilk@pwr.edu.pl

Issue in Honor of Prof. Jacek Młochowski on the occasion of his 80th anniversary

Table of contents

1	Figure S1. Fluorescence spectrum compound 5 (λ_{ex} 354 nm).	S2
2	Figure S2. AFM images and DLS graphs of ZnPc-functionalized polymeric micelles.	S3
4	Figure S3. Fluorescence spectra of ZnPc- functionalized polymeric micelles (λ_{ex} 354 nm).	S4




Figure S1. Fluorescence spectrum of poly(ethylene oxide) modified zinc (II) phthalocyanine (λ_{ex} 354 nm) **5**.



a.**b.**

Figure S2. AFM images and DLS graphs of ZnPc-functionalized Pluronic (**a.** systems 1b) and PLLA (**b.** system 2b) polymeric micelles.




Figure S3. Fluorescence spectra of ZnPc-functionalized Pluronic (**a.** system 1b) and PLLA (**b.** system 2b) polymeric micelles for λ_{ex} 354 nm.