

Supplementary Material

Microwave-assisted synthesis of new imide- and formamide-derivatives of 2(3H)-benzoxazolinones and 2(3H)-benzothiazolinones

Khemissi Seddiki,^{*a,b,c} Wassila Yahia,^b Messaoud Liacha,^b Andrea Defant,^c Denise Sighel,^{c,d} and Ines Mancini^{*c}

^aLaboratoire de Synthèse et Biocatalyse Organique, Faculté des Sciences, Université Badji Mokhtar de Annaba B.P. 12, 23000 Annaba, Algérie

^bEcole Préparatoire Aux Sciences Et Techniques, Citésaf-saf BP-218, 23000, Annaba-Algerie

^cLaboratorio di Chimica Bioorganica, Dipartimento di Fisica, Università di Trento, via Sommarive 14, 38123 Povo-Trento, Italy

^dCentro di Biologia Integrata (CIBIO) Università di Trento, Trento, via Sommarive 9, 38123 Povo-Trento, Italy

E-mail: khemissi_seddiki@yahoo.fr

Tables of Contents

Figure S1. ¹ H NMR spectrum of formamide derivative 8 (400 MHz, CDCl ₃)	S2
Figure S2. ¹ H NMR spectrum of formamide derivative 8 (400 MHz, CD ₃ OD)	S3
Figure S3. ¹ H NMR spectrum of formamide derivative 9 (400 MHz, CDCl ₃)	S4
Figure S4. ¹ H NMR spectrum of formamide derivative 10 (400 MHz, CD ₃ OD)	S5
Figure S5. ¹ H NMR spectrum of formamide derivative 11 (400 MHz, CD ₃ OD)	S6
Figure S6. ¹ H NMR spectrum of formamide derivative 11 (400 MHz, DMSO- <i>d</i> ₆)	S7




Figure S1. ¹H NMR spectrum of formamide derivative **8** (400 MHz, CDCl₃).




Figure S2. ^1H NMR spectrum of formamide derivative **8** (400 MHz, CD_3OD).




Figure S3. ^1H NMR spectrum of formamide derivative **9** (400 MHz, CDCl_3).




Figure S4. ¹H NMR spectrum of formamide derivative **10** (400 MHz, CD₃OD).





Figure S5. ¹H NMR spectrum of formamide derivative **11** (400 MHz, CD₃OD).





Figure S6. ¹H NMR spectrum of formamide derivative **11** (400 MHz, DMSO-*d*₆).