

## Supplementary Material

### ***Tert*-Butylation of naphthalene-2,6-diol and 6-methoxynaphthalen-2-ol**

**Andrey A. Berezin\*, Davide Marinelli**

*School of Chemistry, Cardiff University Main Building, Park Pl, Cardiff CF10 3AT, United Kingdom*


*E-mail: [Berezina@cardiff.ac.uk](mailto:Berezina@cardiff.ac.uk)*

**Dedicated to Professor Oleg A. Rakitin on the occasion of his 65th birthday**


#### **Table of Contents**

1H-NMR of compound <b>2</b>	S2
13C-NMR of compound <b>2</b>	S3
1H-NMR of compound <b>3</b>	S4
13C-NMR of compound <b>3</b>	S5
1H-NMR of compound <b>5</b>	S6
13C-NMR of compound <b>5</b>	S7
1H-NMR of compound <b>6</b>	S8
13C-NMR of compound <b>6</b>	S9
1H-NMR of compound <b>7</b>	S10
13C-NMR of compound <b>7</b>	S11
Crystal data for compound <b>2</b>	S12
Crystal data for compound <b>3</b>	S19
Crystal data for compound <b>5</b>	S24

## 3,7-di-tert-butyl-naphthalene-2,6-diol (2)



3,7-di-tert-butyl-naphthalene-2,6-diol (2)



Current Data Parameters  
 NAME BAB005\_Acetone-d6\_C-13  
 EXPNO 1  
 PROCNO 1


F2 - Acquisition Parameters  
 Date\_ 20170522  
 Time 11:49  
 INSTRUM FOURIER300  
 PROBHD 5 mm DUL 13C-1  
 PULPROG zgpg30  
 TD 65536  
 SOLVENT Acetone  
 NS 512  
 DS 4  
 SWH 24414.063 Hz  
 FIDRES 0.372529 Hz  
 Aq 1.3421773 sec  
 RG 501.187  
 DW 20.480 usec  
 DE 6.50 usec  
 TE 292.3 K  
 D1 2.0000000 sec  
 D11 0.0300000 sec  
 D31 0.00001125 sec  
 D40 0.01745360 sec  
 L4 42  
 L5 60  
 P32 86.00 usec  
 T00 1

----- CHANNEL f1  
 SFO1 75.4861091 MHz  
 NUC1 13C  
 P1 11.25 usec  
 PLW1 31.62299919 W



----- CHANNEL f2  
 SFO2 300.1712007 MHz  
 NUC2 1H  
 CPDPRG2 waltz16  
 PCPD2 86.00 usec  
 PLW2 15.84899998 W  
 PLW12 0.27121001 W  
 PLW13 0.20059000 W

F2 - Processing parameters  
 SI 65536  
 SF 75.4778065 MHz  
 WDW EM  
 SSB 0  
 LB 1.00 Hz  
 GB 0  
 PC 1.40


## 3-tert-butyl-naphthalene-2,6-diol (3)



3-tert-butyl-naphthalene-2,6-diol (3)




Current Data Parameters  
 NAME BAB013\_acetone-d6\_C13  
 EXPNO 1  
 PROCNO 1  
 F2 - Acquisition Parameters  
 Date\_ 20170824  
 Time 13:47  
 INSTRUM FOURIER300  
 PROBHD 5 mm DUL  
 13C-1  
 PULPROG zgpg30  
 TD 64000  
 SOLVENT Acetone  
 NS 512  
 DS 4  
 SWH 34414.083 Hz  
 FREQ0 125.760370 Hz  
 AQ 1.3107200 sec  
 RG 501.187  
 DW 20.480 usec  
 DE 8.50 usec  
 TE 293.2 K  
 CWD2 145.0000000  
 CWDT1 1.0000000  
 D1 1.00000000 sec  
 D11 0.03000000 sec  
 D20 0.00000000 sec  
 D21 0.00000000 sec  
 D30 0.0001125 sec  
 D31 0.0001125 sec  
 D40 0.00020100 sec  
 D42 0.00020250 sec  
 L4 41  
 L5 30  
 P1 11.25 usec  
 P2 22.50 usec  
 TEO 1




===== CHANNEL f1  
 SFO1 75.4681081 MHz  
 NUC1 13C  
 P0 11.25 usec  
 P2 22.50 usec  
 PLW1 31.82389219 W  
 ===== CHANNEL f2  
 SFO2 300.1713007  
 MHz  
 NUC2 1H  
 CPDPRG2 zgpg30  
 P0P02 68.00 usec  
 PLW2 15.64889988 W  
 PLW12 0.27121001 W  
 F2 - Processing parameters  
 SI 32768  
 SF 75.4776085 MHz  
 WDW EM  
 SSB 0  
 LB 1.00 Hz  
 GB 0  
 PC 1.40

## 3,7-di-tert-butyl-6-methoxynaphthalen-2-ol (5)



3,7-di-tert-butyl-6-methoxynaphthalen-2-ol (5)



Current Data Parameters  
 NAME 300\_BABH1\_C003\_C13  
 EXPNO 10  
 PROCNO 1


F2 - Acquisition Parameters  
 Date\_ 20161207  
 Time 20:13  
 INSTRUM FOURIER500  
 PROBNM 5 mm DUL 13C-1  
 PULPROG apt  
 TD 64000  
 SOLVENT CDCl3  
 NS 8000  
 DS 4  
 SWH 34414.000 Hz  
 FIDRES 0.381470 Hz  
 AQ 1.2107200 sec  
 RG 501.187  
 DW 20.480 usec  
 DE 6.50 usec  
 TE 295.1 K  
 CHST0 145.0000000  
 CHST11 1.0000000  
 D1 1.00000000 sec  
 D11 0.02000000 sec  
 D20 0.00000000 sec  
 D21 0.00000000 sec  
 D30 0.00001500 sec  
 D31 0.00001500 sec  
 D40 0.01000100 sec  
 D42 0.00003000 sec  
 LA 44  
 LB 30  
 PH 15.00 usec  
 PD2 80.00 usec  
 TDD 1

CHANNEL F1  
 SFO1 75.4880229 MHz  
 NUC1 13C  
 P0 15.00 usec  
 P2 30.00 usec  
 PLW1 14.18988901 W

CHANNEL F2  
 SFO2 300.1812007 MHz  
 NUC2 1H  
 CPOPRG2 waltz16  
 PCPD2 80.00 usec  
 PLW2 17.00000000 W  
 PLW12 0.32617699 W

F2 - Processing parameters  
 SI 32768  
 SF 75.4803270 MHz  
 WDW no  
 GB 0  
 LB 0 Hz  
 GB 0  
 PC 1.40

3-tert-butyl-6-methoxynaphthalen-2-ol (6)



Current Data Parameters  
 NAME BAS015\_CDCl3\_013  
 EXPNO 1  
 PROCNO 1

F2 - Acquisition Parameters  
 Date\_ 20170524  
 Time 14:39  
 INSTRUM PULPROG300  
 PROBRD 5mm DUL

13C-1  
 PULPROG Apt  
 TD 64000  
 SOLVENT CDCl3  
 NS 512  
 DS 4  
 SWH 24414.063 Hz  
 FIDRES 0.381470 Hz  
 AQ 1.3107200 sec  
 RG 504.187  
 DW 22.480 usec  
 DE 8.33 usec  
 TE 292.3 K  
 CHST2 145.0000000  
 CHST1 1.0000000  
 D1 1.00000000 sec  
 D11 0.00000000 sec  
 D20 0.00000000 sec  
 D21 0.00000000 sec  
 D30 0.00001125 sec  
 D31 0.00001125 sec  
 D40 0.00000100 sec  
 D42 0.00002250 sec  
 L4 41  
 L5 30  
 P1 11.25 usec  
 P2 22.50 usec  
 TDO 1


----- CHANNEL f1  
 SFO1 75.4861001  
 MHz  
 NUC1 13C  
 P0 11.25 usec  
 P2 22.50 usec  
 PLW1 31.02288919 W

----- CHANNEL f2  
 SFO2 300.1713007  
 MHz  
 NUC2 1H  
 CPDPRG2 waltz16  
 PCPD2 88.00 usec  
 PLW2 15.64800000 W  
 PLW12 0.27121001 W


F2 - Processing parameters  
 SI 32768  
 SF 75.4770000 MHz  
 WDW EM  
 GB 0  
 LB 1.00 Hz  
 GB 0  
 PC 1.40



## 2-tert-butoxy-6-methoxynaphthalene (7)




## 2-tert-butoxy-6-methoxynaphthalene (7)



**X-ray measurements:**

Single crystals of **2** and **3** were grown by slow evaporation of  $\text{CHCl}_3$  solutions, while a single crystal of **5** was obtained by evaporation of a hexane solution. Crystallographic studies were undertaken on single crystal mounted in paratone, mounted on the goniometer head with a nylon loop and studied on an Agilent SuperNova Dual three-circle diffractometer using  $\text{Cu-K}\alpha$  ( $\lambda = 1.540598 \text{ \AA}$ ) or  $\text{Mo-K}\alpha$  ( $\lambda = 0.7093187 \text{ \AA}$ ) radiation and a CCD detector. Measurements were typically made at 150(1) K with temperatures maintained using an Oxford Cryostream unless otherwise stated. Data were collected, integrated and corrected for absorption using a numerical absorption correction based on Gaussian integration over a multifaceted crystal model within CrysAlisPro.<sup>[1S]</sup> The structures were solved by direct methods and refined against F2 within SHELXL-2013.<sup>[2S]</sup> A summary of crystallographic data are available as ESI and the structures deposited with the Cambridge Structural Database (CCDC deposition numbers 1558768 (**2**), 1558766(**3**), and 1558767 (**5**)). These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

**Crystal data for compound 2****Table S1.** Crystal data and structure refinement for **2**.

Identification code	1558768 ( <b>2</b> )
Empirical formula	$\text{C}_{36} \text{H}_{52} \text{O}_6$
Formula weight	580.77
Temperature	150(2) K
Wavelength	1.54184 $\text{Å}$
Crystal system	Monoclinic

Space group	I 2/a	
Unit cell dimensions	a = 12.2515(4) Å	$\alpha = 90^\circ$ .
	b = 11.0764(3) Å	$\beta = 104.115(4)^\circ$ .
	c = 24.8986(10) Å	$\gamma = 90^\circ$ .
Volume	3276.8(2) Å <sup>3</sup>	
Z	4	
Density (calculated)	1.177 Mg/m <sup>3</sup>	
Absorption coefficient	0.622 mm <sup>-1</sup>	
F(000)	1264	
Crystal size	0.260 x 0.182 x 0.068 mm <sup>3</sup>	
Theta range for data collection	3.661 to 73.864°.	
Index ranges	-15 ≤ h ≤ 14, -8 ≤ k ≤ 13, -30 ≤ l ≤ 30	
Reflections collected	9291	
Independent reflections	4729 [R(int) = 0.0329]	
Completeness to theta = 67.684°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	1.00000 and 0.52751	
Refinement method	Full-matrix least-squares on F <sup>2</sup>	
Data / restraints / parameters	4729 / 0 / 206	
Goodness-of-fit on F <sup>2</sup>	1.039	
Final R indices [I > 2σ(I)]	R1 = 0.0897, wR2 = 0.2341	
R indices (all data)	R1 = 0.1000, wR2 = 0.2576	
Extinction coefficient	n/a	
Largest diff. peak and hole	0.738 and -0.338 e.Å <sup>-3</sup>	

**Table S2.** Atomic coordinates ( $\times 10^4$ ) and equivalent isotropic displacement parameters ( $\text{Å}^2 \times 10^3$ ) for **2**.  $U(\text{eq})$  is defined as one third of the trace of the orthogonalized  $U_{ij}$  tensor.

	x	y	z	U(eq)
C(1)	4045(2)	3711(2)	5591(1)	26(1)
C(2)	4151(2)	3637(2)	5055(1)	26(1)
C(3)	4754(2)	4512(2)	4830(1)	24(1)
C(4)	4899(2)	4440(2)	4282(1)	25(1)
C(5)	4510(2)	4722(2)	5941(1)	24(1)
C(6)	4335(3)	4844(2)	6531(1)	29(1)
C(7)	4902(3)	3778(3)	6885(1)	38(1)

C(8)	3074(3)	4881(2)	6514(2)	34(1)
C(9)	4863(3)	6013(3)	6811(1)	38(1)
C(10)	3458(2)	-1278(2)	4407(1)	26(1)
C(11)	4096(2)	-1366(2)	4942(1)	26(1)
C(12)	4919(2)	-491(2)	5170(1)	25(1)
C(13)	5616(2)	-574(2)	5716(1)	25(1)
C(14)	3573(2)	-271(2)	4061(1)	26(1)
C(15)	2825(3)	-132(2)	3470(1)	27(1)
C(16)	3040(3)	-1192(3)	3115(1)	38(1)
C(17)	1571(3)	-114(2)	3490(2)	34(1)
C(18)	3058(3)	1047(3)	3196(1)	37(1)
O(1)	3496(1)	2839(2)	5808(1)	33(1)
O(2)	2693(1)	-2156(2)	4188(1)	34(1)
O(3)	2500	1141(2)	5000	30(1)
O(4)	7500	3854(2)	5000	31(1)

---

**Table S3.** Bond lengths [ $\text{\AA}$ ] and angles [ $^{\circ}$ ] for **2**.

---

C(1)-O(1)	1.362(3)
C(1)-C(2)	1.374(3)
C(1)-C(5)	1.447(3)
C(2)-C(3)	1.415(3)
C(2)-H(2)	0.9500
C(3)-C(3)#1	1.414(4)
C(3)-C(4)	1.421(3)
C(4)-C(5)#1	1.375(3)
C(4)-H(4)	0.9500
C(5)-C(4)#1	1.375(3)
C(5)-C(6)	1.541(4)
C(6)-C(7)	1.535(4)
C(6)-C(8)	1.535(5)
C(6)-C(9)	1.537(4)
C(7)-H(7A)	0.9800
C(7)-H(7B)	0.9800
C(7)-H(7C)	0.9800
C(8)-H(8A)	0.9800
C(8)-H(8B)	0.9800
C(8)-H(8C)	0.9800
C(9)-H(9A)	0.9800
C(9)-H(9B)	0.9800
C(9)-H(9C)	0.9800
C(10)-O(2)	1.368(3)
C(10)-C(11)	1.374(3)
C(10)-C(14)	1.438(3)
C(11)-C(12)	1.413(3)
C(11)-H(11)	0.9500
C(12)-C(12)#2	1.420(5)
C(12)-C(13)	1.421(3)
C(13)-C(14)#2	1.379(3)
C(13)-H(13)	0.9500
C(14)-C(13)#2	1.379(3)
C(14)-C(15)	1.539(4)
C(15)-C(18)	1.531(4)
C(15)-C(16)	1.532(4)
C(15)-C(17)	1.549(5)

C(16)-H(16A)	0.9800
C(16)-H(16B)	0.9800
C(16)-H(16C)	0.9800
C(17)-H(17A)	0.9800
C(17)-H(17B)	0.9800
C(17)-H(17C)	0.9800
C(18)-H(18A)	0.9800
C(18)-H(18B)	0.9800
C(18)-H(18C)	0.9800
O(1)-H(1A)	0.8400
O(2)-H(2A)	0.8400
O(3)-H(3)	0.85(4)
O(4)-H(4A)	0.87(4)
O(1)-C(1)-C(2)	121.00(19)
O(1)-C(1)-C(5)	118.0(2)
C(2)-C(1)-C(5)	121.0(2)
C(1)-C(2)-C(3)	121.7(2)
C(1)-C(2)-H(2)	119.1
C(3)-C(2)-H(2)	119.1
C(3)#1-C(3)-C(2)	118.4(3)
C(3)#1-C(3)-C(4)	118.8(2)
C(2)-C(3)-C(4)	122.9(2)
C(5)#1-C(4)-C(3)	123.5(2)
C(5)#1-C(4)-H(4)	118.2
C(3)-C(4)-H(4)	118.2
C(4)#1-C(5)-C(1)	116.6(2)
C(4)#1-C(5)-C(6)	122.3(2)
C(1)-C(5)-C(6)	121.1(2)
C(7)-C(6)-C(8)	110.3(3)
C(7)-C(6)-C(9)	107.8(3)
C(8)-C(6)-C(9)	107.4(3)
C(7)-C(6)-C(5)	109.3(2)
C(8)-C(6)-C(5)	110.5(3)
C(9)-C(6)-C(5)	111.5(2)
C(6)-C(7)-H(7A)	109.5
C(6)-C(7)-H(7B)	109.5
H(7A)-C(7)-H(7B)	109.5
C(6)-C(7)-H(7C)	109.5
H(7A)-C(7)-H(7C)	109.5

H(7B)-C(7)-H(7C)	109.5
C(6)-C(8)-H(8A)	109.5
C(6)-C(8)-H(8B)	109.5
H(8A)-C(8)-H(8B)	109.5
C(6)-C(8)-H(8C)	109.5
H(8A)-C(8)-H(8C)	109.5
H(8B)-C(8)-H(8C)	109.5
C(6)-C(9)-H(9A)	109.5
C(6)-C(9)-H(9B)	109.5
H(9A)-C(9)-H(9B)	109.5
C(6)-C(9)-H(9C)	109.5
H(9A)-C(9)-H(9C)	109.5
H(9B)-C(9)-H(9C)	109.5
O(2)-C(10)-C(11)	120.6(2)
O(2)-C(10)-C(14)	118.0(2)
C(11)-C(10)-C(14)	121.4(2)
C(10)-C(11)-C(12)	121.4(2)
C(10)-C(11)-H(11)	119.3
C(12)-C(11)-H(11)	119.3
C(11)-C(12)-C(12)#2	118.4(3)
C(11)-C(12)-C(13)	122.8(2)
C(12)#2-C(12)-C(13)	118.7(2)
C(14)#2-C(13)-C(12)	123.2(2)
C(14)#2-C(13)-H(13)	118.4
C(12)-C(13)-H(13)	118.4
C(13)#2-C(14)-C(10)	116.8(2)
C(13)#2-C(14)-C(15)	121.4(2)
C(10)-C(14)-C(15)	121.9(2)
C(18)-C(15)-C(16)	108.6(3)
C(18)-C(15)-C(14)	112.2(2)
C(16)-C(15)-C(14)	109.2(2)
C(18)-C(15)-C(17)	107.3(2)
C(16)-C(15)-C(17)	109.8(3)
C(14)-C(15)-C(17)	109.6(3)
C(15)-C(16)-H(16A)	109.5
C(15)-C(16)-H(16B)	109.5
H(16A)-C(16)-H(16B)	109.5
C(15)-C(16)-H(16C)	109.5
H(16A)-C(16)-H(16C)	109.5
H(16B)-C(16)-H(16C)	109.5



C(15)-C(17)-H(17A)	109.5
C(15)-C(17)-H(17B)	109.5
H(17A)-C(17)-H(17B)	109.5
C(15)-C(17)-H(17C)	109.5
H(17A)-C(17)-H(17C)	109.5
H(17B)-C(17)-H(17C)	109.5
C(15)-C(18)-H(18A)	109.5
C(15)-C(18)-H(18B)	109.5
H(18A)-C(18)-H(18B)	109.5
C(15)-C(18)-H(18C)	109.5
H(18A)-C(18)-H(18C)	109.5
H(18B)-C(18)-H(18C)	109.5
C(1)-O(1)-H(1A)	109.5
C(10)-O(2)-H(2A)	109.5


Symmetry transformations used to generate equivalent atoms:

#1 -x+1,-y+1,-z+1 #2 -x+1,-y,-z+1

**Table S4.** Anisotropic displacement parameters ( $\text{\AA}^2 \times 10^3$ ) for **2**. The anisotropic displacement factor exponent takes the form:  $-2 \sum_{i,j} h^i k^j a^i b^j U^{ij}$

	U11	U22	U33	U23	U13	U12
C(1)	25(1)	22(1)	32(1)	2(1)	7(1)	0(1)
C(2)	24(1)	21(1)	31(1)	-1(1)	6(1)	0(1)
C(3)	21(1)	21(1)	30(1)	-1(1)	5(1)	2(1)
C(4)	22(1)	22(1)	30(1)	-3(1)	5(1)	0(1)
C(5)	21(1)	27(1)	26(1)	0(1)	7(1)	4(1)
C(6)	31(2)	31(1)	24(1)	1(1)	8(1)	0(1)
C(7)	43(2)	41(2)	29(1)	5(1)	5(1)	5(1)
C(8)	33(2)	40(2)	35(2)	1(1)	16(1)	1(1)
C(9)	46(2)	39(2)	31(1)	-6(1)	13(1)	-8(1)
C(10)	24(1)	22(1)	32(1)	-4(1)	7(1)	-1(1)
C(11)	26(1)	22(1)	32(1)	1(1)	9(1)	0(1)
C(12)	23(1)	23(1)	29(1)	0(1)	8(1)	2(1)
C(13)	25(1)	22(1)	29(1)	0(1)	9(1)	1(1)
C(14)	24(1)	27(1)	27(1)	-3(1)	8(1)	2(1)
C(15)	26(1)	31(1)	24(1)	-4(1)	5(1)	2(1)

C(16)	47(2)	38(2)	29(1)	-7(1)	9(1)	3(1)
C(17)	28(2)	42(2)	31(2)	-1(1)	2(1)	3(1)
C(18)	42(2)	39(1)	27(1)	4(1)	1(1)	-2(1)
O(1)	40(1)	28(1)	33(1)	0(1)	13(1)	-9(1)
O(2)	37(1)	28(1)	35(1)	-1(1)	3(1)	-10(1)
O(3)	28(1)	22(1)	41(1)	0	14(1)	0
O(4)	23(1)	21(1)	47(1)	0	4(1)	0

**Crystal data for compound 3****Table S5.** Crystal data and structure refinement for **3**.

Identification code	1558766 ( <b>3</b> )	
Empirical formula	C <sub>14</sub> H <sub>16</sub> O <sub>2</sub>	
Formula weight	216.27	
Temperature	293(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P 21/c	
Unit cell dimensions	a = 6.4201(3) Å	α = 90°.

	$b = 7.1231(5) \text{ \AA}$	$\beta = 94.017(6)^\circ$
	$c = 25.2730(14) \text{ \AA}$	$\gamma = 90^\circ$
Volume	$1152.92(12) \text{ \AA}^3$	
Z	4	
Density (calculated)	$1.246 \text{ Mg/m}^3$	
Absorption coefficient	$0.082 \text{ mm}^{-1}$	
F(000)	464	
Crystal size	$0.775 \times 0.646 \times 0.135 \text{ mm}^3$	
Theta range for data collection	2.972 to 29.635°.	
Index ranges	$-6 \leq h \leq 8, -9 \leq k \leq 7, -33 \leq l \leq 34$	
Reflections collected	5223	
Independent reflections	2724 [R(int) = 0.0187]	
Completeness to theta = 25.242°	100.0 %	
Absorption correction	Gaussian	
Max. and min. transmission	1.000 and 0.340	
Refinement method	Full-matrix least-squares on $F^2$	
Data / restraints / parameters	2724 / 0 / 145	
Goodness-of-fit on $F^2$	1.037	
Final R indices [ $I > 2\sigma(I)$ ]	R1 = 0.0474, wR2 = 0.1102	
R indices (all data)	R1 = 0.0707, wR2 = 0.1240	
Extinction coefficient	n/a	
Largest diff. peak and hole	0.222 and -0.223 e. $\text{\AA}^{-3}$	

**Table S6.** Atomic coordinates ( $\times 10^4$ ) and equivalent isotropic displacement parameters ( $\text{\AA}^2 \times 10^3$ ) for **3**.  $U(\text{eq})$  is defined as one third of the trace of the orthogonalized  $U_{ij}$  tensor.

	x	y	z	U(eq)
O(1)	11338(2)	4645(2)	1190(1)	33(1)
O(2)	4243(2)	2605(2)	3480(1)	39(1)
C(5)	6799(2)	3215(2)	2225(1)	20(1)
C(12)	9232(2)	1617(3)	528(1)	37(1)
C(4)	6514(2)	2866(2)	1673(1)	21(1)
C(3)	7961(2)	3323(2)	1318(1)	22(1)
C(11)	7572(2)	2953(2)	721(1)	26(1)
C(10)	8679(2)	4083(2)	2426(1)	22(1)
C(2)	9848(2)	4192(2)	1535(1)	23(1)

C(9)	8974(2)	4437(2)	2978(1)	25(1)
C(1)	10174(2)	4554(2)	2065(1)	23(1)
C(6)	5270(2)	2723(2)	2579(1)	22(1)
C(7)	5613(2)	3077(2)	3109(1)	26(1)
C(14)	5447(2)	2025(3)	591(1)	32(1)
C(8)	7476(2)	3949(2)	3310(1)	27(1)
C(13)	7596(2)	4824(2)	419(1)	35(1)

---

**Table S7.** Bond lengths [Å] and angles [°] for **3**.

---

O(1)-C(2)	1.3767(15)
O(1)-H(1A)	0.8200
O(2)-C(7)	1.3712(16)
O(2)-H(10)	0.8200
C(5)-C(4)	1.4168(18)
C(5)-C(6)	1.4177(18)
C(5)-C(10)	1.4184(19)
C(12)-C(11)	1.534(2)
C(12)-H(12A)	0.9600
C(12)-H(12B)	0.9600
C(12)-H(12C)	0.9600
C(4)-C(3)	1.3747(18)
C(4)-H(4)	0.9300
C(3)-C(2)	1.4350(19)
C(3)-C(11)	1.5331(19)
C(11)-C(14)	1.5313(19)
C(11)-C(13)	1.537(2)
C(10)-C(1)	1.4106(18)
C(10)-C(9)	1.4175(19)
C(2)-C(1)	1.3647(19)
C(9)-C(8)	1.365(2)
C(9)-H(9)	0.9300
C(1)-H(1)	0.9300
C(6)-C(7)	1.3662(19)
C(6)-H(6)	0.9300
C(7)-C(8)	1.410(2)
C(14)-H(14A)	0.9600
C(14)-H(14B)	0.9600
C(14)-H(14C)	0.9600

C(8)-H(8)	0.9300
C(13)-H(13A)	0.9600
C(13)-H(13B)	0.9600
C(13)-H(13C)	0.9600
C(2)-O(1)-H(1A)	109.5
C(7)-O(2)-H(10)	109.5
C(4)-C(5)-C(6)	122.12(12)
C(4)-C(5)-C(10)	118.48(12)
C(6)-C(5)-C(10)	119.40(12)
C(11)-C(12)-H(12A)	109.5
C(11)-C(12)-H(12B)	109.5
H(12A)-C(12)-H(12B)	109.5
C(11)-C(12)-H(12C)	109.5
H(12A)-C(12)-H(12C)	109.5
H(12B)-C(12)-H(12C)	109.5
C(3)-C(4)-C(5)	123.69(12)
C(3)-C(4)-H(4)	118.2
C(5)-C(4)-H(4)	118.2
C(4)-C(3)-C(2)	116.33(12)
C(4)-C(3)-C(11)	121.98(12)
C(2)-C(3)-C(11)	121.69(12)
C(14)-C(11)-C(3)	111.59(11)
C(14)-C(11)-C(12)	106.97(13)
C(3)-C(11)-C(12)	110.57(11)
C(14)-C(11)-C(13)	107.85(12)
C(3)-C(11)-C(13)	109.42(12)
C(12)-C(11)-C(13)	110.38(12)
C(1)-C(10)-C(9)	123.01(12)
C(1)-C(10)-C(5)	118.22(12)
C(9)-C(10)-C(5)	118.78(13)
C(1)-C(2)-O(1)	120.71(12)
C(1)-C(2)-C(3)	121.61(12)
O(1)-C(2)-C(3)	117.67(12)
C(8)-C(9)-C(10)	120.63(13)
C(8)-C(9)-H(9)	119.7
C(10)-C(9)-H(9)	119.7
C(2)-C(1)-C(10)	121.67(12)
C(2)-C(1)-H(1)	119.2
C(10)-C(1)-H(1)	119.2

C(7)-C(6)-C(5)	120.19(13)
C(7)-C(6)-H(6)	119.9
C(5)-C(6)-H(6)	119.9
C(6)-C(7)-O(2)	123.90(13)
C(6)-C(7)-C(8)	120.59(13)
O(2)-C(7)-C(8)	115.51(12)
C(11)-C(14)-H(14A)	109.5
C(11)-C(14)-H(14B)	109.5
H(14A)-C(14)-H(14B)	109.5
C(11)-C(14)-H(14C)	109.5
H(14A)-C(14)-H(14C)	109.5
H(14B)-C(14)-H(14C)	109.5
C(9)-C(8)-C(7)	120.41(13)
C(9)-C(8)-H(8)	119.8
C(7)-C(8)-H(8)	119.8
C(11)-C(13)-H(13A)	109.5
C(11)-C(13)-H(13B)	109.5
H(13A)-C(13)-H(13B)	109.5
C(11)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5

---


Symmetry transformations used to generate equivalent atoms:

**Table S8.** Anisotropic displacement parameters ( $\text{\AA}^2 \times 10^3$ ) for **3**. The anisotropic displacement factor exponent takes the form:  $-2p^2 [ h^2 a^* 2U^{11} + \dots + 2 h k a^* b^* U^{12} ]$

---

	U <sup>11</sup>	U <sup>22</sup>	U <sup>33</sup>	U <sup>23</sup>	U <sup>13</sup>	U <sup>12</sup>
O(1)	21(1)	40(1)	37(1)	-6(1)	8(1)	-8(1)
O(2)	48(1)	39(1)	30(1)	-4(1)	13(1)	-17(1)
C(5)	20(1)	14(1)	27(1)	1(1)	-1(1)	2(1)
C(12)	30(1)	43(1)	38(1)	-12(1)	3(1)	7(1)
C(4)	17(1)	19(1)	27(1)	1(1)	-2(1)	0(1)
C(3)	20(1)	17(1)	27(1)	1(1)	1(1)	4(1)
C(11)	22(1)	29(1)	26(1)	-1(1)	3(1)	3(1)
C(10)	21(1)	14(1)	28(1)	0(1)	-3(1)	3(1)

C(2)	18(1)	19(1)	32(1)	1(1)	4(1)	2(1)
C(9)	25(1)	19(1)	31(1)	-3(1)	-4(1)	1(1)
C(1)	17(1)	18(1)	34(1)	-3(1)	-2(1)	-2(1)
C(6)	22(1)	16(1)	28(1)	-1(1)	1(1)	-2(1)
C(7)	31(1)	19(1)	29(1)	2(1)	7(1)	0(1)
C(14)	26(1)	42(1)	26(1)	-4(1)	-1(1)	-2(1)
C(8)	35(1)	21(1)	24(1)	-2(1)	-3(1)	1(1)
C(13)	34(1)	41(1)	30(1)	6(1)	4(1)	2(1)

**Crystal data for compound 5****Table S9.** Crystal data and structure refinement for **5**.

Identification code	1558767 ( <b>5</b> )	
Empirical formula	C <sub>19</sub> H <sub>26</sub> O <sub>2</sub>	
Formula weight	286.40	
Temperature	150(2) K	
Wavelength	1.54184 Å	
Crystal system	Orthorhombic	
Space group	P b c n	
Unit cell dimensions	a = 21.420(2) Å	α = 90°.

	$b = 6.4137(6) \text{ \AA}$	$\beta = 90^\circ$ .
	$c = 24.380(2) \text{ \AA}$	$\gamma = 90^\circ$ .
Volume	$3349.3(6) \text{ \AA}^3$	
Z	8	
Density (calculated)	$1.136 \text{ Mg/m}^3$	
Absorption coefficient	$0.557 \text{ mm}^{-1}$	
F(000)	1248	
Crystal size	$0.940 \times 0.157 \times 0.084 \text{ mm}^3$	
Theta range for data collection	$3.626$ to $70.134^\circ$ .	
Index ranges	$-18 \leq h \leq 25$ , $-7 \leq k \leq 7$ , $-29 \leq l \leq 23$	
Reflections collected	8334	
Independent reflections	3165 [R(int) = 0.0777]	
Completeness to theta = $67.684^\circ$	99.9 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	1.00000 and 0.28053	
Refinement method	Full-matrix least-squares on $F^2$	
Data / restraints / parameters	3165 / 0 / 198	
Goodness-of-fit on $F^2$	1.096	
Final R indices [ $I > 2\sigma(I)$ ]	$R1 = 0.0968$ , $wR2 = 0.2641$	
R indices (all data)	$R1 = 0.1214$ , $wR2 = 0.2853$	
Extinction coefficient	n/a	
Largest diff. peak and hole	$0.282$ and $-0.383 \text{ e.\AA}^{-3}$	

**Table S10.** Atomic coordinates ( $\times 10^4$ ) and equivalent isotropic displacement parameters ( $\text{\AA}^2 \times 10^3$ ) for **5**.  $U(\text{eq})$  is defined as one third of the trace of the orthogonalized  $U_{ij}$  tensor.

	x	y	z	U(eq)
C(1)	855(2)	9630(6)	1287(1)	38(1)
C(2)	912(2)	8899(6)	766(2)	37(1)
C(3)	1279(2)	9975(6)	363(1)	37(1)
C(4)	1594(2)	11745(6)	539(1)	36(1)
C(5)	1549(2)	12515(6)	1083(1)	34(1)
C(6)	1864(2)	14336(6)	1260(1)	37(1)
C(7)	1791(2)	15066(6)	1787(1)	36(1)
C(8)	1393(2)	14034(6)	2176(1)	37(1)
C(9)	1099(2)	12264(5)	1995(1)	36(1)



C(10)	1168(2)	11461(6)	1460(1)	35(1)
C(11)	264(2)	5969(7)	971(2)	51(1)
C(12)	1304(2)	9254(7)	-233(2)	45(1)
C(13)	1559(3)	6991(10)	-263(2)	71(2)
C(14)	649(2)	9325(7)	-485(1)	44(1)
C(15)	1715(3)	10664(11)	-585(2)	79(2)
C(16)	1318(2)	14873(6)	2764(1)	42(1)
C(17)	1948(2)	14796(8)	3062(2)	56(1)
C(18)	855(2)	13527(7)	3090(2)	51(1)
C(19)	1057(3)	17098(6)	2754(2)	55(1)
O(1)	2097(2)	16822(4)	1963(1)	48(1)
O(2)	619(1)	7134(4)	583(1)	46(1)

---

**Table S11.** Bond lengths [ $\text{\AA}$ ] and angles [ $^\circ$ ] for **5**.

---

C(1)-C(2)	1.361(5)
C(1)-C(10)	1.417(5)
C(1)-H(1)	0.9500
C(2)-O(2)	1.368(5)
C(2)-C(3)	1.434(5)
C(3)-C(4)	1.389(5)
C(3)-C(12)	1.527(5)
C(4)-C(5)	1.419(5)
C(4)-H(4)	0.9500
C(5)-C(10)	1.402(5)
C(5)-C(6)	1.415(5)
C(6)-C(7)	1.376(5)
C(6)-H(6)	0.9500
C(7)-O(1)	1.373(4)
C(7)-C(8)	1.437(5)
C(8)-C(9)	1.371(5)
C(8)-C(16)	1.539(5)
C(9)-C(10)	1.409(5)
C(9)-H(9)	0.9500
C(11)-O(2)	1.424(5)
C(11)-H(11A)	0.9800
C(11)-H(11B)	0.9800

C(11)-H(11C)	0.9800
C(12)-C(15)	1.525(7)
C(12)-C(14)	1.532(6)
C(12)-C(13)	1.552(7)
C(13)-H(13A)	0.9800
C(13)-H(13B)	0.9800
C(13)-H(13C)	0.9800
C(14)-H(14A)	0.9800
C(14)-H(14B)	0.9800
C(14)-H(14C)	0.9800
C(15)-H(15A)	0.9800
C(15)-H(15B)	0.9800
C(15)-H(15C)	0.9800
C(16)-C(19)	1.532(6)
C(16)-C(17)	1.533(7)
C(16)-C(18)	1.536(6)
C(17)-H(17A)	0.9800
C(17)-H(17B)	0.9800
C(17)-H(17C)	0.9800
C(18)-H(18A)	0.9800
C(18)-H(18B)	0.9800
C(18)-H(18C)	0.9800
C(19)-H(19A)	0.9800
C(19)-H(19B)	0.9800
C(19)-H(19C)	0.9800
O(1)-H(1A)	0.8400
C(2)-C(1)-C(10)	121.4(3)
C(2)-C(1)-H(1)	119.3
C(10)-C(1)-H(1)	119.3
C(1)-C(2)-O(2)	123.2(3)
C(1)-C(2)-C(3)	121.5(3)
O(2)-C(2)-C(3)	115.3(3)
C(4)-C(3)-C(2)	116.7(3)
C(4)-C(3)-C(12)	121.6(3)
C(2)-C(3)-C(12)	121.7(4)
C(3)-C(4)-C(5)	122.7(3)
C(3)-C(4)-H(4)	118.6
C(5)-C(4)-H(4)	118.6
C(10)-C(5)-C(6)	118.4(3)

C(10)-C(5)-C(4)	119.0(3)
C(6)-C(5)-C(4)	122.6(3)
C(7)-C(6)-C(5)	120.6(3)
C(7)-C(6)-H(6)	119.7
C(5)-C(6)-H(6)	119.7
O(1)-C(7)-C(6)	121.2(3)
O(1)-C(7)-C(8)	116.9(3)
C(6)-C(7)-C(8)	121.9(3)
C(9)-C(8)-C(7)	116.1(3)
C(9)-C(8)-C(16)	122.8(3)
C(7)-C(8)-C(16)	121.1(3)
C(8)-C(9)-C(10)	123.5(3)
C(8)-C(9)-H(9)	118.2
C(10)-C(9)-H(9)	118.2
C(5)-C(10)-C(9)	119.4(3)
C(5)-C(10)-C(1)	118.7(3)
C(9)-C(10)-C(1)	121.9(3)
O(2)-C(11)-H(11A)	109.5
O(2)-C(11)-H(11B)	109.5
H(11A)-C(11)-H(11B)	109.5
O(2)-C(11)-H(11C)	109.5
H(11A)-C(11)-H(11C)	109.5
H(11B)-C(11)-H(11C)	109.5
C(15)-C(12)-C(3)	112.1(4)
C(15)-C(12)-C(14)	106.6(4)
C(3)-C(12)-C(14)	109.8(3)
C(15)-C(12)-C(13)	109.0(4)
C(3)-C(12)-C(13)	109.9(4)
C(14)-C(12)-C(13)	109.3(4)
C(12)-C(13)-H(13A)	109.5
C(12)-C(13)-H(13B)	109.5
H(13A)-C(13)-H(13B)	109.5
C(12)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5
C(12)-C(14)-H(14A)	109.5
C(12)-C(14)-H(14B)	109.5
H(14A)-C(14)-H(14B)	109.5
C(12)-C(14)-H(14C)	109.5
H(14A)-C(14)-H(14C)	109.5

H(14B)-C(14)-H(14C)	109.5
C(12)-C(15)-H(15A)	109.5
C(12)-C(15)-H(15B)	109.5
H(15A)-C(15)-H(15B)	109.5
C(12)-C(15)-H(15C)	109.5
H(15A)-C(15)-H(15C)	109.5
H(15B)-C(15)-H(15C)	109.5
C(19)-C(16)-C(17)	111.0(4)
C(19)-C(16)-C(18)	107.3(4)
C(17)-C(16)-C(18)	107.7(3)
C(19)-C(16)-C(8)	110.4(3)
C(17)-C(16)-C(8)	109.7(3)
C(18)-C(16)-C(8)	110.7(3)
C(16)-C(17)-H(17A)	109.5
C(16)-C(17)-H(17B)	109.5
H(17A)-C(17)-H(17B)	109.5
C(16)-C(17)-H(17C)	109.5
H(17A)-C(17)-H(17C)	109.5
H(17B)-C(17)-H(17C)	109.5
C(16)-C(18)-H(18A)	109.5
C(16)-C(18)-H(18B)	109.5
H(18A)-C(18)-H(18B)	109.5
C(16)-C(18)-H(18C)	109.5
H(18A)-C(18)-H(18C)	109.5
H(18B)-C(18)-H(18C)	109.5
C(16)-C(19)-H(19A)	109.5
C(16)-C(19)-H(19B)	109.5
H(19A)-C(19)-H(19B)	109.5
C(16)-C(19)-H(19C)	109.5
H(19A)-C(19)-H(19C)	109.5
H(19B)-C(19)-H(19C)	109.5
C(7)-O(1)-H(1A)	109.5
C(2)-O(2)-C(11)	117.6(3)

---

Symmetry transformations used to generate equivalent atoms:

**Table S12.** Anisotropic displacement parameters ( $\text{\AA}^2 \times 10^3$ ) for **5**. The anisotropic displacement factor exponent takes the form:  $-2p^2 [ h^2 a^* 2U^{11} + \dots + 2 h k a^* b^* U^{12} ]$

–	U11	U22	U33	U23	U13	U12
–						
C(1)	57(2)	37(2)	21(2)	3(1)	-1(2)	-2(2)
C(2)	48(2)	39(2)	26(2)	2(1)	-7(2)	2(1)
C(3)	38(2)	53(2)	20(2)	-4(1)	-4(2)	4(2)
C(4)	36(2)	52(2)	20(2)	1(1)	0(1)	1(1)
C(5)	39(2)	43(2)	20(2)	2(1)	-2(1)	2(1)
C(6)	46(2)	44(2)	21(2)	1(1)	2(2)	-1(1)
C(7)	45(2)	40(2)	22(2)	1(1)	0(1)	-1(1)
C(8)	54(2)	40(2)	17(2)	2(1)	1(2)	1(1)
C(9)	53(2)	36(2)	18(2)	4(1)	0(1)	0(1)
C(10)	46(2)	40(2)	20(2)	3(1)	-2(1)	0(1)
C(11)	68(3)	45(2)	38(2)	4(2)	-5(2)	-7(2)
C(12)	40(2)	73(3)	23(2)	-14(2)	1(2)	6(2)
C(13)	66(3)	104(4)	43(2)	-32(3)	-10(2)	34(3)
C(14)	49(2)	66(3)	17(2)	-7(2)	-4(2)	3(2)
C(15)	64(3)	152(6)	22(2)	-30(3)	14(2)	-39(3)
C(16)	64(3)	45(2)	18(2)	-1(1)	3(2)	1(2)
C(17)	75(3)	71(3)	23(2)	-6(2)	-3(2)	2(2)
C(18)	81(3)	54(2)	19(2)	1(2)	9(2)	1(2)
C(19)	88(3)	46(2)	29(2)	-2(2)	12(2)	6(2)
O(1)	67(2)	48(2)	28(1)	-6(1)	7(1)	-14(1)
O(2)	67(2)	41(1)	31(1)	-4(1)	-6(1)	-6(1)

–

**References:**

1S - CrysAlisPro, Agilent Technologies, Version 1.171.37.33 (release 27-03-2014 CrysAlis171.NET)

2S - G. M. Sheldrick, *SHELX-2013: Program for the Solution of Crystal Structures* **2013** University of Göttingen.