

## Synthesis of some novel oxazolopyranoquinolinones from 3-amino-4-hydroxypyrano[3,2-c]quinolinedione

Hany M. Hassanin,<sup>\*a</sup> Ibrahim M. Abdou,<sup>b</sup> and Abdeltawab M. Saeed<sup>a,b</sup>

<sup>a</sup>Department of Chemistry, Faculty of Education, Ain Shams University Roxy 11711 Cairo, Egypt

<sup>b</sup>Department of Chemistry, College of Science, UAE University Al-Ain 15551, UAE

Email: [hanyhassnin@yahoo.com](mailto:hanyhassnin@yahoo.com)


Received 05-29-2017

Accepted 07-04-2017

Published on line 10-08-2017

### Abstract

6-*n*-Butyl-3-nitropyrano[3,2-*c*]quinolinone was chemoselectively reduced to an amine derivative in a good yield (70%) using an efficient and simple tin-hydrochloric acid reaction. The novel 3-aminopyranoquinolinone was utilized as a precursor to produce some new interesting tetracyclic fused oxazolopyranoquinolinone analogues. The structures of the obtained quinolinone derivatives were confirmed using IR, LC-MS, ESI-MS and NMR techniques.




**Keywords:** Tin reduction, oxazole, pyran, quinolinone, heterocyclization

## Introduction

Pyranoquinoline skeletons including both a quinoline ring and a pyran moiety, afford exceptional biological activities.<sup>1-4</sup> Rings annulated to the pyranoquinolinone units showed potential medicinal properties such as antibacterial,<sup>5</sup> anticoagulant,<sup>6</sup> antitumor,<sup>7</sup> and microtubule-targeting agents.<sup>8</sup> Despite the extensive body of published work on the synthesis of this type of compounds, we are not aware of any report that describes a simple procedure for the annulation of an oxazole nucleus to the pyranoquinolinone moiety to give a tetracyclic system. Oxazole derivatives have been found in numerous natural products and many of them have been shown to have an interesting broad range of biological activities,<sup>9,10</sup> such as in inhibition of *Streptococcus mutans* biofilm formation,<sup>11</sup> antipsychotics,<sup>12</sup> HIV inhibitory,<sup>13</sup> and anti-tumoral activities.<sup>14</sup> It is also known that incorporation of fluorine atoms into molecules of heterocyclic compounds leads to a significant increase in their biological activities.<sup>15</sup> Recently, the synthesis of fluoroquinolones has become of increasing interest since the quinoline skeleton is present in many chemotherapeutic agents.<sup>16-18</sup> These findings prompted us to explore the combination of an oxazole nucleus and fluorine atom within the pyranoquinolinone moiety in one molecular framework. In continuation of our research focused on the chemistry of pyrano[3,2-*c*]quinolinedione derivatives,<sup>19,20</sup> we now disclose the preparation of a new synthetically valuable 3-aminopyranoquinolone derivative in order to obtain a novel series of pyranoquinolinones incorporating an oxazole and/or fluorophenylloxazole at face *c*. We hope that these new compounds will have useful biological activities.


## Results and Discussion

The nitration of **1** using a mixture of concentrated nitric acid and sulfuric acid gave 3-nitropyrano[3,2-*c*]quinoline-2,5-dione **2**. The ESI-MS of compound **2** showed four intense peaks at *m/z* 331.1835, 353.1434, 354.2070 and 386.7095, corresponding to  $[M+H]^+$ ,  $[M+Na]^+$ ,  $[M+Na+H]^{2+}$  and  $[2M+Na]^+$ , respectively. After installing a nitro group in the 3-position of pyrano[3,2-*c*]quinoline-2,5-dione, we turned to its conversion into an amino group by an efficient chemoselective reduction method. Attempts to reduce the nitro compound **2** using activated iron<sup>21</sup> or  $FeCl_3/Zn/DMF/H_2O$ ,<sup>22</sup> were not effective; a mixture of products was isolated, which could not be separated in a satisfactory manner to obtain any considerable amount of the amino derivative. Instead, we used tin, as previously described by Jampilek<sup>23</sup> in the synthesis of 4-hydroxy-3-nitroquinolin-2-one. Accordingly, the nitro-derivative **2** was reduced with tin and concentrated hydrochloric acid at 130 °C to produce 3-amino pyrano[3,2-*c*]quinoline-2,5-dione **3** in 70% yield (Scheme 1). The reduction proceeded efficiently with excellent chemoselectivity without affecting other functional carbonyl groups or the pyranoquinolinone system.




Scheme 1

The formation of the amino analogue was confirmed by its IR spectrum showing bands at 3483 and 3395  $\text{cm}^{-1}$  as a double peak due to the  $\text{NH}_2$  group. The mass spectrum of compound **3** revealed a molecular ion peak at  $m/z$  300, corresponding to its formula weight (300.11). Moreover, compound **3** showed a quasimolecular ion peak at  $m/z$  301.2  $[\text{M}+\text{H}]^+$  and a sodiated molecular ion peak at  $m/z$  323.1  $[\text{M}+\text{Na}]^+$  in the positive electron spray ionization-MS corresponding to  $\text{C}_{16}\text{H}_{16}\text{N}_2\text{O}_4$ . There were two singlet signals in the gCOZY spectrum of compound **3**: 3.72 ppm characteristic of an  $\text{NH}_2$  group and 12.31 ppm characteristic of the OH group (figure 1).




**Figure 1.**  $^1\text{H}$  chemical shifts and correlation in COZY spectrum of compound **3**.

The gHSQC, gHMBC and  $^1\text{H}$ ,  $^{13}\text{C}$ -HMBC spectra lead to assignment of the  $^{13}\text{C}$  – chemical shift of compound **3** as shown in Figure 2.



**Figure 2.** The network of  $\text{H-C}$  long-range heteronuclear correlations and  $^{13}\text{C}$  chemical shift assignment of compound **3**.

Heterocyclization at face [c] of the pyrone moiety occurred when compound **3** was mixed with triethyl orthoformate, under solvent-free conditions. The reaction probably proceeds initially *via* condensation to produce the intermediate **4**. This intermediate can undergo intramolecular cyclondensation to give the desired oxazolopyrano[3,2-*c*]quinoline-4,5-dione **5** in low yield (42%) (Scheme 2). The IR spectrum of compound **5** revealed the absence of bands due to the amino group. The  $^1\text{H}$  NMR spectrum of compound **5** showed a new characteristic singlet signal at 8.79 ppm characteristic of the CH of the oxazole ring. The spectrum also revealed the absence of deuterium-exchangeable protons, observed in the starting material as singlet signals, at 3.72 and 12.31 ppm, for the N-H and O-H protons, respectively.  $^{13}\text{C}$ -NMR spectrum of compound **5** demonstrated the presence of thirteen  $sp^2$ -hybridised carbons in the region 100-165 ppm due to the aromatic tetracyclic system. The same compound was obtained from the reaction of the amine **3** with formic acid in moderate yield 54%.




## Scheme 2

Our attempts to improve the yield of the above reaction either by changing the reaction conditions or isolating the intermediate **4** were not successful. We expected that an alternative method for the preparation of compound **5** through a two-step reaction could produce better results. Thus, the intermediate enamine **6** was prepared by treating amine **3** with dimethylformamide dimethylacetal (DMF-DMA), and isolated by chromatography using ethyl acetate/hexane 6:4 as the eluent. The IR spectrum of enamine **6** revealed the absence of the double stretching bands of the amino group, as well as an ion peak at  $m/z$  356.2 in an ESI-MS  $[\text{M}+\text{H}]^+$  measurement in accordance with its structure. The  $^1\text{H}$  NMR spectrum of the product showed two separate singlet signals at 3.05 ppm and 3.14 ppm attributed to two methyl protons of the  $\text{Me}_2\text{N}$  group, while the proton of the  $(\text{CH}=\text{N})$  group was observed at 8.16 ppm. In the  $^{13}\text{C}$  NMR spectrum of enamine **6**, two new  $sp^3$  methyl carbons appeared at  $\delta$  31.1 and 34.9, and a signal at 207.0 ppm assignable to the  $sp^2$  carbon of the

(CH=N) group. Boiling enamine **6**, in glacial acetic acid, (Scheme 2), effected its heterocyclization to afford oxazolopyran **5** in 80% yield. The time, yield and purity showed there are considerable advantages to the two-step sequence.


Homologue **8** was obtained by thermal cyclocondensation of the amine **3** with triethyl orthoacetate. As expected, this conversion occurred without isolation of proposed intermediate **7**, in a low yield (46%) (Scheme 3). Refluxing the amine **3** with acetic anhydride afforded the same oxazole derivative **8** in a somewhat greater yield, 52%. Therefore, we adopted another procedure to synthesize compound **8**, using a two-step synthesis. The first step was acetylation of the amine **3** to produce acetamide **9** as an isolated intermediate. The ESI-MS analysis of compound **9** showed an  $[M+H]^+$  ion at  $m/z$  343.1, and an abundant  $[M+Na]^+$  ion at  $m/z$  365.1. The IR spectrum of acetyl derivative **9** had an absorption broad band characteristic for (N-H) at  $3082\text{ cm}^{-1}$  and a new stretching band at  $1703\text{ cm}^{-1}$  corresponding to the acetamide carbonyl group. In the  $^1\text{H}$  NMR spectrum of **9**, the protons of the acetyl group were observed at 2.06 ppm as a singlet signal, and the (N-H) proton appeared at 9.12 ppm as a singlet signal. The  $^{13}\text{C}$  NMR spectrum of **9** exhibited the presence of two new signals at 23.2 ppm and 207.0 ppm due to the  $\text{CH}_3$  and  $\text{C}=\text{O}$  of the acetyl group. Compound **9** was heated in refluxing AcOH and  $\text{Ac}_2\text{O}$  to afford oxazole derivative **8** (Scheme 3). The disappearance of both hydroxyl group and carbonyl of the acetyl group are the prominent features in the IR spectrum of oxazole derivative **8** and its ESI-MS spectrum showed an  $[M+H]^+$  ion at  $m/z$  325.3 and a  $[M+Na]^+$  ion at 347.1. A  $[2M+Na]^+$  ion was also observed at  $m/z$  671.1.



Scheme 3

The  $^1\text{H}$  NMR signal for the methyl group on the oxazole ring was observed as a singlet at 2.73 ppm, while the (N-H) proton was no longer present. The  $^{13}\text{C}$ -NMR spectrum of compound **8** showed a signal at 13.8 ppm for the methyl carbon atom at position 2 of the oxazole ring.

A nucleophilic addition reaction occurred when **3** was heated with phenyl isothiocyanate in absolute ethanol at reflux, producing phenylthiourea derivative **10** in a good yield (Scheme 4). The IR spectrum of compound **10** displayed characteristic absorption bands at 3342, 3156  $\text{cm}^{-1}$  for NH groups and in the region of 1283  $\text{cm}^{-1}$ , corresponding to C=S vibrations, in addition to an  $\alpha$ -pyran carbonyl group at 1748  $\text{cm}^{-1}$ . The  $^1\text{H}$ -NMR spectrum of **10** contained two singlet signals attributed to two NH protons, at 8.94 and 9.76 ppm. Signals integrating for the nine aromatic protons of the phenyl and benzo groups were observed in the region 7.06–8.19 ppm, confirming the formation of structure **10**. There were two deuterium-exchangeable signals for the two (N-H) protons. The  $^{13}\text{C}$ -NMR of compound **10** had nineteen signals in the region 100-164 ppm belonging to the aromatic carbon atoms and the thiocarbonyl group. The mass spectrum exhibited a molecular ion peak at  $m/z$  435 (15%) ( $\text{M}^+$ ), along with ( $\text{M}+1$ ) at 436 (5%). The base peak at  $m/z$  189 (100%) correspond to a molecular formula of  $\text{C}_{11}\text{H}_{11}\text{NO}_2$ .




#### Scheme 4

We speculated that boiling phenylthiourea derivative **10** in dry toluene containing *p*-toluenesulfonic acid as a catalyst, would effect its intramolecular heterocyclization. The probable products are either oxazole **11** or oxazole **12**. The IR spectrum of the product displayed characteristic absorption bands at 3342 and 3100  $\text{cm}^{-1}$  for NH, in addition to  $\alpha$ -pyrano carbonyl group at 1745  $\text{cm}^{-1}$ . The  $^1\text{H}$ -NMR spectrum of the product was characterized by the presence of only four aromatic signals in the region 7.36–8.05 ppm corresponding to the

protons of the benzene ring of the quinolinone. There was, in addition, a deuterium-exchangeable singlet signal assignable to an NH proton at 10.78. The  $^{13}\text{C}$ -NMR spectrum had a signal at 177 ppm due to  $\text{C}=\text{S}_{\text{oxazole}}$ , and thirteen other signals belonging to the aromatic carbon atoms of the tetracyclic system. The mass spectrum of the product showed a molecular ion peak at  $m/z$  342 ( $\text{M}^+$ ) (50%) corresponding to a molecular formula of  $\text{C}_{17}\text{H}_{14}\text{N}_2\text{O}_4\text{S}$ . Building on the above data, the structure is confirmed as oxazolopyranoquinolinone **12**. The proposed pathway for the formation of compound **12** is thought to involve an intramolecular addition-elimination as depicted in Scheme 4. We suggest that an initial nucleophilic addition of the phenolic group to the thiocarbonyl function of the neighboring side-chain is followed by loss a molecule of aniline. The structure of compound **12** was chemically confirmed *via* its direct preparation from the reaction of **3** and carbon disulfide in boiling pyridine.

In another approach towards the nucleophilic addition followed by cyclization to give an oxazolopyranoquinolinone derivative, was reacting compound **3** with ammonium thiocyanate in boiling pyridine (Scheme 5). Lassaigne's test and elemental microanalysis of the product showed the absence of sulfur, so the product cannot contain sulfur. The IR spectrum revealed strong absorption bands at 3488 and 3394  $\text{cm}^{-1}$  due to stretching vibration of  $\text{NH}_2$ .




**Scheme 5**

The  $^1\text{H}$ -NMR spectrum of product **13** showed the presence of two deuterium-exchangeable protons, as a broad singlet signal at 7.71 ppm due to  $\text{NH}_2$  protons. The  $^{13}\text{C}$ -NMR spectrum of compound **13** revealed the presence of four  $sp^3$  hybridized carbon atoms due to the butyl group and thirteen  $sp^2$  hybridized carbon atoms belonging to the aromatic carbon atoms of oxazolopyrano quinolinone system. The mass spectrum of compound **13** gave a molecular ion peak at  $m/z$  325 ( $\text{M}^+$ ) corresponding to the molecular formula  $\text{C}_{17}\text{H}_{15}\text{N}_3\text{O}_4$ .

A series of a novel fluoro derivatives of pyranoquinolinones **14a-c** were synthesized from the reaction of aminopyranoquinolinone **3** with some fluorobenzoyl chlorides in refluxing dry THF (Scheme 6).

Evidence for the formation of compounds **14a-c** came from their IR spectra where there was characteristic absorption at *ca.* 3447-3389  $\text{cm}^{-1}$  due to an N-H bond, and a new stretching signal due to the  $\text{C}=\text{O}$  group of the amide between 1639 to 1677  $\text{cm}^{-1}$ . The ESI-MS spectra of compounds **14b-c** exhibited

$[M+Na]^+$  ions ( $m/z$ : 445.0197, 445.1098 and 511.2984) in accordance with their structures. Furthermore, the amide (N-H) proton of this series was observed between 9.53 ppm to 9.81 ppm as a singlet signal in the  $^1H$  NMR spectra of compounds **14a-c**. The aryl carbon atoms were observed between 100 and 162 ppm in their  $^{13}C$  NMR spectra.



### Scheme 6

Intramolecular heterocyclization of compounds **14a-c** by heating in a mixture of glacial acetic acid and acetic anhydride (80:20) gave our target fluoro derivatives **15a-c** in good yields (Scheme 6). The IR spectra of all derivatives **15a-c** showed the absence of the stretching absorption bands attributed to OH and C=O amide. The ESI-MS spectrum of compound **15a** shows a  $[M+H]^+$  ion at  $m/z$  405.1330 and a  $[M+2H]^{2+}$  ion at 406.1340, the  $[2M+Na]^+$  ion was also observed at  $m/z$  831.3452. ESI-MS spectrum of compound **15b** showed an  $[M+H]^+$  ion at  $m/z$  405.2520 and the  $[2M+Na]^+$  ion at  $m/z$  831.3457. The ESI-MS spectrum of compound **15c** revealed a quasimolecular ion peak at  $m/z$  471.1  $[M+H]^+$  and a sodiated molecular ion peak at  $m/z$  493.2  $[M+Na]^+$ .

## Experimental Section

**General.** TLC analysis of the reaction mixtures was performed using Fluka analytical silica gel 60 F254 nm TLC plates. For column chromatography, Fluka analytical silica gel 60 0.063-0.2 mm (70–230 mesh ASTM) was used for separations. Melting points were recorded on Sanyo Gallenkamp MPD 350-BM 3.5 Melting Point apparatus. A Thermo Nicolet Nexus 470 FT-IR spectrophotometer was used for IR analyses.  $^1H$ -NMR (400 MHz) and  $^{13}C$ -NMR (101 MHz) measurements were performed using Varian-400 MHz spectrometer, and chemical shifts are expressed in  $\delta$  (ppm) relative to TMS (in  $CDCl_3$  or  $DMSO-d_6$  as solvent) as the internal standard. Elemental microanalyses were performed Perkin-Elmer CHN-2400II at the Chemical War Department, Ministry of Defence, Cairo, Egypt. Mass spectra were recorded on a Gas Chromatographic GCMSqp 1000 ex Shimadzu instrument at 70 eV. For mass spectra, a triple-quadrupole tandem mass spectrometer (Micromass W Quattro micro<sup>TM</sup>, Waters Corp., Milford, MA, USA) or Waters ZMD Quadrupole equipped with electrospray ionization (ESI) were used.



**6-*n*-Butyl-3*H*-pyrano[3,2-*c*]quinoline-2,4,5(6*H*)-trione (1).** Yield (92%), mp 227–229 °C, (mp 228-230 °C).<sup>24</sup>

**6-*n*-Butyl-4-hydroxy-3-nitro-2*H*-pyrano[3,2-*c*]quinoline-2,5(6*H*)-dione (2).** To a solution of compound **1** (2.85 g, 10 mmol) in glacial AcOH acid (50 mL), kept under 50 °C, a mixture of conc. HNO<sub>3</sub> (2.5 mL) and conc. H<sub>2</sub>SO<sub>4</sub> (2.5 mL) was added dropwise with continuous stirring. The precipitate formed was filtered off, dried and crystallized from glacial AcOH to furnish **2** as pale brown needles (2.7 g, 82%), mp 203-205 °C. IR (KBr, cm<sup>-1</sup>): 3444 broad band (OH), 3087 (CH<sub>aromatic</sub>), 2961, 2932, 2872 (CH<sub>aliphatic</sub>), 1742(C=O<sub>α-pyrone</sub>), 1672 (C=O<sub>quinolone</sub>) and 1566 (C=C). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub>: 1.01 (t, *J* 8.0 Hz, 3H, C4'), 1.45-1.54 (m, 2H, C3'), 1.71- 1.79 (m, 2H, C2'), 4.33 (t, 2H, *J* 8.0 Hz, C1'), 7.39 (t, 1H, *J* 8.0 Hz, C9-H), 7.46 (d, *J* 8.0 Hz, 1H, C7-H), 7.65 (t, 1H, *J* 8.0 Hz, C8-H), 8.24 (dd, 1H, *J* 8.0, 4.0 Hz, C10-H), 12.38 (s, 1H, OH). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ<sub>C</sub>: 13.7 (s, C4'), 20.1 (s, C3'), 29.5 (s, C2'), 42.8 (s, C1'), 97.8 (s, C3), 113.1 (s, C4a), 115.7 (s, C10a), 118.4 (s, C7), 125.1 (s, C10), 125.5 (s, C8), 136.0 (s, C6a), 138.9 (s, C9), 153.4 (s, C10b), 159.3 (s, C4), 163.1 (s, C2), 165.4 (s, C5). ESI-MS *m/z*: 331.1835 [M+H]<sup>+</sup>, 353.1434 [M+Na]<sup>+</sup>, 354.2070 [M+Na+H]<sup>2+</sup>, 386.7095 [2M+Na]<sup>+</sup>. Anal. Calcd for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>6</sub> (330.30): C, 58.18; H, 4.27; N, 8.48. Found C, 57.52; H, 4.20; N, 8.39%.

**3-Amino-6-*n*-butyl-4-hydroxy-2*H*-pyrano[3,2-*c*]quinoline-2,5(6*H*)-dione (3).** A mixture of compound **2**, (10 mmol) tin metal powder (4.00 g, 33 mmol) and conc. HCl (30 mL) was stirred at 130 °C for 1 h. MeOH (25 mL) was added to the reaction mixture and the mixture heated under reflux for 2 h until the color became clear yellow. The reaction mixture was filtered hot and the filtrate poured onto ice (200 g). The obtained solid was filtered off, dried under vacuum and crystallized from EtOH (96%) to produce **3** as yellow crystals (2.4 g, 79%), mp 211-212 °C. IR (KBr, cm<sup>-1</sup>): 3384, 3359 (NH<sub>2</sub>), 2981, 2932 (CH<sub>3</sub>, CH<sub>2aliphatic</sub>), 1710 (C=O<sub>α-pyrone</sub>), 1680 (C=O<sub>quinolone</sub>) and 1608 (C=C<sub>aliphatic</sub>). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ<sub>H</sub>: 0.98 (t, *J* 8.0 Hz, 3H, C4'), 1.42-1.49 (m, 2H, C3'), 1.71- 1.75 (m, 2H, C2'), 3.72 (bs, 2H, NH<sub>2</sub>), 4.28 (t, 2H, *J* 8Hz, 1°C), 7.33 (t, 1H, *J* 8.0 Hz, C9- H), 7.38 (d, 1H, *J* 8.0 Hz, C7- H), 7.61 (t, 1H, *J* 8.0 Hz, C8-H), 8.12 (dd, 1H, *J* 1.2, 8.00Hz, C10-H), 12.31(s, 1H, OH exchangeable with D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: 13.7 (s, C4'), 20.2 (s, C3'), 29.5 (s, C2'), 42.2 (s, C1'), 101.4 (s, C3) 113.9 (s, C4a), 114.8 (s, C7), 114.2 (s, C10a), 123.4 (s, C10), 123.7 (s, C9), 131.7 (s, C8), 136.2 (s, C6a), 146.1 (s, C10b), 159.1 (s, C4), 149.8 (s, C2), 162.1 (s, C5). *m/z* (*I*<sub>r</sub> %): 302 (M<sup>+</sup>+2, 2), 301 (M<sup>+</sup>+1, 19), 300 (M<sup>+</sup>, 100), 299 (M<sup>+</sup>-1, 2), 244 (40), 161(75). ESI-MS *m/z*: 301.2 [M+H]<sup>+</sup>, 323.1 [M+Na]<sup>+</sup>. Anal. Calcd for C<sub>16</sub>H<sub>16</sub>N<sub>2</sub>O<sub>4</sub> (300.32): C, 63.99; H, 5.37; N, 9.33. Found C, 63.72; H, 5.26; N, 9.29%.

**5-*n*-Butyl-4*H*-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5*H*)-dione (5).**

**Method a.** A mixture of compound **3** (3 g, 10 mmol) and triethyl orthoformate (8 mL, 50 mmol) was heated for 12 h. The progress of the reaction was monitored by TLC. The solid deposited after cooling was filtered off then washed with hot EtOH then crystallized from glacial AcOH to afford compound **5** (1.31 g, 42%).

**Method b:** A mixture of compound **3** (3 g, 10 mmol) and formic acid (25 mL), was heated for 6 h. The deposited precipitate was filtered off, air dried and crystallized from AcOH to give compound **5** (1.68 g, 54%).

**Method c.** A mixture of compound **6** (3.55 g, 10 mmol) and glacial acetic acid was heated under reflux for 6 h. Progress of the reaction was monitored by TLC. The reaction mixture was filtered and the filtrate poured onto ice. The obtained solid was filtered off, washed with water (3 x 20 mL), then Et<sub>2</sub>O (3 x 20 mL) and crystallized from AcOH to give compound **5** as yellow crystals (2.49 g, 80%), mp 237-239 °C. IR (KBr, cm<sup>-1</sup>): 3451 broad band (OH), 3078 (CH<sub>aromatic</sub>), 2955, 2917, 2848 (CH<sub>aliphatic</sub>), 1742(C=O<sub>α-pyrone</sub>), 1669 (C=O<sub>quinolone</sub>) and 1646 (C=N<sub>oxazolo</sub>), 1591(C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, δ, 400 MHz) δ<sub>H</sub>: 0.90 (t, *J* 8.0 Hz, 3H, C4'), 1.34-1.44 (m, 2H, C3'), 1.59- 1.68 (m, 2H, C2'), 4.36 (t, 2H, *J* 8Hz, 1°C), 7.57 (t, 1H, *J* 8.0 Hz, C9- H), 7.81 (d, 1H, *J* 8.0 Hz, C7- H), 7.91 (t, 1H, *J* 8.0 Hz, C8-H), 8.20 (dd, 1H, *J* 1.2, 8.00Hz, C10-H), 8.79(s, 1H, CH<sub>oxazole</sub>). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: 14.1 (s, C4'), 19.8 (s, C3'), 29.6 (s, C2'), 42.4 (s, C1'), 100.7 (s, C4a) 113.7 (s, C7), 116.6 (s, C10a), 124.4 (s, C10), 133.5 (s, C9), 135.5 (s, 1C, 3), 137.0 (s, 1C, 8), 138.2 (s, C6a), 144.1 (s, C4), 145.8 (s, C2<sub>oxazolo</sub>), 158.7 (s,

C10b), 162.8 (s, C,2), 163.7 (s, C5).  $m/z$  ( $I_r$  %): 311 ( $M^+ + 1$ , 10), 310 ( $M^+$ , 30), 243 (42), 188 (95), 132 (100). Anal. Calcd for  $C_{17}H_{14}N_2O_4$  (310.31): C, 65.80; H, 4.55; N, 9.03. Found C, 65.52; H, 4.20; N, 8.99%.

***N'*-(6-*n*-Butyl-4-hydroxy-2,5-dioxo-5,6-dihydro-2*H*-pyrano[3,2-*c*]quinolin-3-yl)-*N,N*-dimethylformimidamide (6).** A mixture of compound **3** (3 g, 10 mmol) and DMF-DMA (3.0 mL, 10 mmol) in (50 mL) dry toluene was heated under reflux for 6 h. The progress of the reaction was monitored by TLC. After completion of reaction, the solvent was removed under vacuum to produce a yellow oily residue. The targeted compound **4** was obtained using column chromatography (ethyl acetate/hexane 6:4) as yellow crystals (2.98 g, 84%), mp 192-193 °C. IR (KBr,  $cm^{-1}$ ): 3409 broad band (OH), 2953, 2918, 2866 (CH *aliphatic.*), 1715 (C=O  $\alpha$ -pyrone), 1650 (C=O *quinolone*) and 1613 (C=N *imid*), 1539 (C=C *aromatic*).  $^1H$  NMR (DMSO- $d_6$ )  $\delta_H$ : 0.91 (t,  $J$  8Hz, 3H, C4'), 1.33-1.42 (m, 2H, C3'), 1.54 - 1.61 (m, 2H, C2'), 3.05 (s, 3H, N-CH<sub>3</sub>), 3.14 (s, 3H, N-CH<sub>3</sub>), 4.21 (t, 2H,  $J$  8Hz, C1'), 7.34 (t, 1H,  $J$  8Hz, C9-H), 7.60 (d, 1H,  $J$  8Hz, C7-H), 7.70 (t,  $J$  8Hz, 1H, C8-H), 8.04 (dd, 1H,  $J$  1.2, 8Hz, C10-H), 8.16 (s, 1H, CH= N).  $^{13}C$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_C$ : 14.2 (s, C4'), 20.0 (s, C3'), 29.7 (s, C2'), 31.1 (s, N-CH<sub>3</sub>), 34.9 (s, N-CH<sub>3</sub>), 35.6 (s, C1'), 105.9 (s, C3), 113.8 (s, C4a), 115.5 (s, C10a), 122.9 (s, C7), 123.9 (s, C10), 133.0 (s, C9), 138.4 (s, C8), 156.3 (s, C6a), 157.1 (s, C10b), 159.5 (s, C2), 159.9 (s, C4), 160.5 (s, C5), 207.0 (s, N=CH). ESI-MS  $m/z$ : 356.2 [M+H]<sup>+</sup>, 357.2 [M+2H]<sup>2+</sup>, 379.1 [M+H+Na]<sup>2+</sup>. Anal. Calcd for  $C_{19}H_{21}N_3O_4$  (355.40): C, 64.21; H, 5.96; N, 11.82. Found C, 64.19; H, 5.20; N, 11.79%.

**5-*n*-Butyl-2-methyl-4*H*-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5*H*)-dione (8).**

**Method a.** A mixture of compound **3** (3 g, 10 mmol) and triethyl orthoacetate (9 mL, 50 mmol) was heated for 12 h. The progress of the reaction was monitored by TLC. The solid deposited after cooling was filtered off and washed with hot EtOH. The targeted compound crystallized from AcOH to give compound **8** as pale brown crystals (1.45 g, 46%).

**Method b.** A mixture of compound **3** (3 g, 10 mmol) and Ac<sub>2</sub>O (25 mL), was heated for 6 h. The solid deposited was filtered off, air dried and crystallized from AcOH to give compound **8** (1.64 g, 52%).

**Method c.** To a mixture of **9** (3.4 g, 10 mmol) in glacial AcOH (80%) (40 mL), Ac<sub>2</sub>O (20%) (10 mL) was added dropwise with stirring at rt for a period of (10 mL) then heated under reflux till the reaction was complete as judged by TLC (24 h). The reaction mixture was cooled to rt and poured on ice (200 g). The brown precipitate was filtered off, washed with water (3 x 20 mL), dried and crystallized from AcOH to give compound **8** as yellow crystals (2.93 g, 90%), mp 223-225 °C, IR (KBr,  $cm^{-1}$ ): 3081 (CH *aromatic*), 2953, 2924, 2866 (CH *aliphatic.*), 1755 (C=O  $\alpha$ -pyrone), 1650 (C=O *quinolone*), and 1623 (C=N *oxazolo*), 1586 (C=C *aromatic*).  $^1H$  NMR (400 MHz, CDCl<sub>3</sub>)  $\delta_H$ : 1.00 (t, 3 H,  $J$  8.00 Hz, C4'), 1.46-1.57 (m, 2H, C3'), 1.71 - 1.81 (m, 2 H, C2'), 2.73 (s, 3 H, CH<sub>3</sub> *oxazolo*), 4.36 (t, 2 H,  $J$  8.00 Hz, C1'), 7.37 (t, 1 H,  $J$  8.00 Hz, C9-H), 7.45 (d, 1 H,  $J$  8.00 Hz, C7-H), 7.71 (t, 1 H,  $J$  8.00 Hz, C8-H), 8.32 (dd, 1 H,  $J$  8.00, 1.17 Hz, C10-H).  $^{13}C$  NMR (101 MHz, DMSO- $d_6$ )  $\delta_C$ : 13.8 (s, CH<sub>3</sub> *oxazolo*), 14.4 (s, C4'), 20.2 (s, C3'), 29.6 (s, C2'), 42.5 (s, C1'), 100.8 (s, C3), 113.5 (s, C10a), 114.9 (s, C7), 123.2 (s, C10), 124.7 (s, C9), 124.8 (s, C4a), 133.4 (s, C8), 138.7 (s, C6a), 154.5 (s, C4), 155.9 (s, C2 *oxazolo*), 156.7 (s, C10b), 156.9 (s, C2), 164.7 (s, C5). ESI-MS  $m/z$ : 325.3 [M+H]<sup>+</sup>, 347.1 [M+Na]<sup>+</sup>, 671.1 [2M+Na]<sup>+</sup>. Anal. Calcd for  $C_{18}H_{16}N_2O_4$  (324.34): C, 66.66; H, 4.97; N, 8.64. Found C, 66.19; H, 4.99; N, 8.59%.

***N*-(6-*n*-Butyl-4-hydroxy-2,5-dioxo-5,6-dihydro-2*H*-pyrano[3,2-*c*]quinolin-3-yl)acetamide (9).** To a solution of compound **3** (3 g, 10 mmol) in THF (50 mL), acetyl chloride (0.75 mL, 10 mmol) was added and the mixture was heated under reflux for 4 h. After completion of reaction (as indicated by the disappearance of starting material on TLC) the reaction mixture was poured onto crushed ice (100 g). The separated solid was filtered off, washed with water (3 x 10 mL), dried and crystallized from EtOH to give compound **9** as off white crystals (3.25 g, 95%), mp 185-187 °C. IR (KBr,  $cm^{-1}$ ): 3415 (broad band, OH), 3082 (NH), 3050 (CH *aromatic*), 2953, 2924, 2866 (CH *aliphatic.*), 1738 (C=O  $\alpha$ -pyrone), 1703 (C=O *acetyl*), 1668 (C=O *quinolone*), and 1568 (C=C *aromatic*).  $^1H$  NMR (400 MHz, DMSO- $d_6$ )  $\delta_H$ : 0.91 (t, 3 H,  $J$  8.00 Hz, C4'), 1.34 - 1.46 (m, 2 H, C3'), 1.59 - 1.70 (m, 2 H, C2'), 2.06 (s, 3

H, CH<sub>3</sub>acetyl), 4.34 (t, 2 H, *J* 8.00 Hz, C1'), 7.52 (t, 1 H, *J* 8.00 Hz, C9-H), 7.73 (d, 1 H, *J* 8.00 Hz, C7-H), 7.85 (t, 1 H, *J* 8.00 Hz, C8-H), 8.14 (dd, 1 H, *J* 8.00, 1.17 Hz, C10-H), 9.12 (s, 1 H, N-H<sub>amide</sub> exchangeable in D<sub>2</sub>O), 13.64 (bs, 1 H, C4-OH exchangeable in D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ<sub>C</sub>: 13.7 (s, C4'), 20.1 (s, C3'), 23.2 (s, CH<sub>3</sub>acetyl), 29.5 (s, C2'), 42.4 (s, C1'), 100.0 (s, C3), 101.6 (s, C4a), 113.6 (s, C10a), 115.1 (s, C7), 124.2 (s, C10) 124.8 (s, C9), 134.0 (s, C8), 137.9 (s, C6a), 156.9 (s, C4), 159.7 (s, C10b), 162.8 (s, C2), 169.0 (s, C5), 207.0 (s, C=O<sub>acetyl</sub>). ESI-MS *m/z*: 343.1 [M+H]<sup>+</sup>, 365.1 [M+Na]<sup>+</sup>. Anal. Calcd for C<sub>18</sub>H<sub>18</sub>N<sub>2</sub>O<sub>5</sub> (342.35): C, 63.15; H, 5.30; N, 8.18. Found C, 62.81; H, 5.20; N, 8.16%.

**1-(6-*n*-Butyl-4-hydroxy-2,5-dioxo-5,6-dihydro-2H-pyrano[3,2-*c*]quinolin-3-yl)-3-phenylthiourea (10).** To a solution of compound **3** (3 g, 10 mmol) and pyridine (1 mL) in absolute EtOH (50 mL), phenyl isothiocyanate (1.21 mL, 10 mmol) was added. The reaction mixture was heated under reflux for 6 h. The reaction was monitored by TLC. The reaction mixture was filtered hot and the filtrate was poured on ice (200 g). The obtained solid was filtered off, washed with water (3 x 10 mL), dried and crystallized from DMF to give compound **10** as a pale yellow powder (2.8 g, 65%), mp 126-129 °C. IR (KBr, cm<sup>-1</sup>): 3442 broad band (OH), 3342, 3156 (N-H), 2967, 2928, 2850 (CH<sub>aliphatic</sub>), 1748 (C=O<sub>α-pyrone</sub>), 1679 (C=O<sub>quinolone</sub>), 1617 (C=N), 1568 (C=C<sub>aromatic</sub>) and 1283 (C=S). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ<sub>H</sub>: 0.90 (t, *J* 8.00 Hz, 3 H, C4'), 1.34 - 1.43 (m, 2 H, C3'), 1.56 - 1.66 (m, 2 H, C2'), 4.33 (t, 2 H, *J* 8.00 Hz, C1'), 7.05 - 7.14 (m, 1 H, C-H<sub>phenyl</sub>), 7.29 (t, 1 H, *J* 8.00 Hz, C-H<sub>phenyl</sub>), 7.42 (t, 1 H, *J* 8.00 Hz, C-H<sub>phenyl</sub>), 7.53 (t, 1 H, *J* 8.00 Hz, C9-H), 7.68 (d, 1 H, *J* 8.00 Hz, C7-H), 7.77 (d, 1 H, *J* 8.00 Hz, C-H<sub>phenyl</sub>), 7.85 (t, 1 H, *J* 8.00 Hz, C8-H), 8.01 (d, 1 H, *J* 8.00 Hz, C-H<sub>phenyl</sub>), 8.15 (dd, 1 H, *J* 8.00, 1.17 Hz, C10-H), 8.94 (s, 1 H, N-H exchangeable in D<sub>2</sub>O), 9.76 (s, 1 H, N-H exchangeable in D<sub>2</sub>O), 13.73 (bs, 1H, OH exchangeable in D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: 14.1 (s, C4'), 19.9 (s, C3'), 29.6 (s, C2'), 42.0 (s, C1'), 101.6 (s, C3), 113.9 (s, C4a), 114.5 (s, C7), 116.4 (s, C10a), 122.7 (s, C<sub>phenyl</sub>), 123.6 (s, C10), 123.8 (s, C9), 124.1 (s, C<sub>phenyl</sub>), 124.4 (s, C<sub>phenyl</sub>), 124.8 (s, C<sub>phenyl</sub>), 132.2 (s, C8), 136.1 (s, C6a), 139.8 (s, C<sub>phenyl</sub>), 149.0 (s, C10b), 155.9 (s, C4), 159.3 (s, C2), 162.7 (s, C5), 163.0 (s, C=S). MS: *m/z* (relative intensity): 436 [M<sup>+</sup>+1; 5], 435 [M<sup>+</sup>; 15], 189 (100). Anal. Calcd for C<sub>23</sub>H<sub>21</sub>N<sub>3</sub>O<sub>4</sub>S (435.51): C, 63.43; H, 4.86; N, 9.65; S, 7.36. Found C, 63.32; H, 4.76; N, 9.36; S, 7.23%.

#### **5-*n*-Butyl-2-thioxo-4H-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5H)-dione (12)**

**Method a.** To a solution of compound **10** (4.3 g, 10 mmol) in dry toluene (50 mL), *p*-toluenesulfonic acid (0.3 g) was added. The reaction mixture was heated under reflux for 24 h. The progress of the reaction was monitored by TLC. The reaction mixture filtered and poured on ice (200 g). The obtained precipitate was filtered off, washed with water (3 x 10 mL), dried and crystallized from EtOH to produce **12** as light brown crystals (1.95 g, 57%), mp 163-165 °C. IR (KBr, cm<sup>-1</sup>): 3342, 3100 (N-H), 3056 (CH<sub>aromatic</sub>), 2928, 2918, 2849 (CH<sub>aliphatic</sub>), 1745 (C=O<sub>α-pyrone</sub>), 1676 (C=O<sub>quinolone</sub>), 1537 (C=C<sub>aromatic</sub>) and 1296 (C=S). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ<sub>H</sub>: ppm 0.92 (t, 3 H, *J* 8.00 Hz, C4'), 1.33 - 1.45 (m, 2 H, C3'), 1.55 - 1.65 (m, 2 H, C2'), 4.26 (t, 2 H, *J* 8.00 Hz, C1'), 7.36 (t, 1 H, *J* 8.00 Hz, C9-H), 7.48 (d, 1 H, *J* 8.00 Hz, C7-H), 7.77 (t, 1 H, *J* 8.00 Hz, C8-H), 8.05 (dd, 1 H, *J* 8.00, 0.8 Hz, C10-H), 10.78 (s, 1 H, N-H exchangeable in D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: ppm 13.8 (s, C4'), 20.2 (s, C3'), 29.4 (s, C2'), 42.2 (s, C1'), 135.6 (s, C3), 129.7 (s, C4a), 114.6 (s, C7), 119.0 (s, C10a), 122.6 (s, C10), 124.6 (s, C9), 133.7 (s, C8), 136.4 (s, C6a), 156.9 (s, C10b), 141.2 (s, C4), 157.7 (s, C2), 161.8 (s, C5), 177.9 (s, C=S<sub>thioxo</sub>). MS: *m/z* (relative intensity): 343 [M<sup>+</sup>+1; 9], 342 [M<sup>+</sup>; 50], 286 [M<sup>+</sup>-Bu; 24]. Anal. Calcd for C<sub>17</sub>H<sub>14</sub>N<sub>2</sub>O<sub>4</sub>S (342.38): C, 59.64; H, 4.12; N, 8.18; S, 9.37. Found C, 59.12; H, 4.10; N, 8.08; S, 9.20%.

**Method b.** To a solution of compound **3** (3 g, 10 mmol) in pyridine (50 mL), carbon disulfide (0.9 mL, 15 mmol) was added dropwise. The mixture was heated under reflux for 12 h, monitored by TLC. The reaction mixture was filtered hot and the filtrate was poured onto ice (100 g). The resulting precipitate was filtered off, washed with water (3 x 10 mL), dried and crystallized from EtOH to produce **12** as light brown crystals (2.05 g, 60%), mp 163-165 °C.

**2-Amino-5-*n*-butyl-4H-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5H)-dione (13).** Compound **3** (3 g, 10 mmol) was mixed with pyridine (50 mL) and ammonium thiocyanate (1.2 g, 15 mmol). The resulted mixture was heated under reflux till the reaction was complete as judged by TLC (24 h). The reaction mixture was cooled to rt and poured on ice (200 g). The obtained precipitate was filtered off, washed with water (3 x 10 mL), dried and crystallized from EtOH to furnish **13** as yellow crystals (2.45 g, 75%), mp 258-260°C. IR (KBr, cm<sup>-1</sup>): 3488, 3394 double peaks (NH<sub>2aromatic</sub>), 3086 (CH<sub>aromatic</sub>), 2959, 2930, 2864 (CH<sub>aliphatic</sub>), 1710 (C=O<sub>αpyrone</sub>), 1682 (C=O<sub>quinolone</sub>), 1634 (C=N<sub>oxazolo</sub>), 1586 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ<sub>H</sub>: 0.89 (t, 3 H, *J* 8.00 Hz, C4'), 1.28 - 1.45 (m, 2H, C3'), 1.50 - 1.67 (m, 2H, C2'), 4.28 (t, 2H, *J* 8.00 Hz, C1'), 7.42 (t, 1 H, *J* 8.00 Hz, C9-H), 7.56 (d, 1 H, *J* 8.00 Hz, C7-H), 7.71 (bs, 2 H, C-NH<sub>2oxazolo</sub>), 8.01 (t, 1 H, *J* 8.00 Hz, C8-H), 8.06 (dd, 1 H, *J* 8.00, 1.2 Hz, C10-H). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: 14.1 (s, C4'), 19.9 (s, C3'), 29.6 (s, C2'), 42.0 (s, C1'), 101.6 (s, C4a), 114.0 (s, C10a), 116.3 (s, C7), 122.7 (s, C10), 124.3 (s, C9), 124.3 (s, C3), 132.2 (s, C8), 136.1 (s, C6a), 158.9 (s, C4), 158.9 (s, C10b), 162.8 (s, C2), 164.9 (s, C2<sub>oxazolo</sub>), 165.6 (s, C5). MS: *m/z* (relative intensity): 326 [M<sup>+</sup>+1; 5], 325 [M<sup>+</sup>; 10], 309 [M<sup>+</sup>-NH<sub>2</sub>+H; 6], 269 [M<sup>+</sup>-Bu+H; 20], 144 (100). Anal. Calcd for C<sub>17</sub>H<sub>15</sub>N<sub>3</sub>O<sub>4</sub> (325.33): C, 62.76; H, 4.65; N, 12.92. Found C, 62.57; H, 4.63; N, 12.81%.

**General procedure for preparation of compounds 14a-c.** To a solution of compound **3** (3 g, 10 mmol) in THF (25 mL), an equivalent amount of 2-fluorobenzoyl chloride (1.79 mL), 4-fluorobenzoyl chloride (1.60 mL) or 4-trifluoromethoxybenzoyl chloride (1.60 mL) was added and the reaction mixture was heated under reflux overnight. The targeted compounds **14a-c** were precipitated. The obtained solid was filtered off, dried and crystallized from EtOH to produce the corresponding compound **14**.

***N*-(6-*n*-Butyl-4-hydroxy-2,5-dioxo-5,6-dihydro-2H-pyrano[3,2*c*]quinolin-3-yl)-2-fluorobenzamide (14a).** Pale yellow powder (2.9 g, 72%), mp 215-217 °C. IR (KBr, cm<sup>-1</sup>): 3398 (O-H), 3290 (NH), 3068 (CH<sub>aromatic</sub>), 2954, 2930, 2864 (CH<sub>aliphatic</sub>), 1738 (C=O<sub>α-pyrone</sub>), 1666 (C=O<sub>quinolone</sub>), 1615 (C=O<sub>arylcarbonyl</sub>) and 1537 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ<sub>H</sub>: 0.92 (t, *J* 8.00 Hz, 3H, C4'), 1.35 - 1.47 (m, 2H, C3'), 1.60 - 1.72 (m, 2H, C2'), 4.33 (t, *J* 8.00 Hz, 2H, C1'), 7.28 - 7.35 (m, 2H, C9-H, C-H<sub>phenyl</sub>), 7.51 - 7.60 (m, 2H, C7-H, C-H<sub>phenyl</sub>), 7.67 (t, 1 H, *J* 7.43 Hz, C8-H), 7.88 (d, 1 H, *J* 9.80 Hz, C-H<sub>phenyl</sub>) 7.91 (dd, *J* 8.80, 1.20 Hz, 1H, C-H<sub>phenyl</sub>), 8.18 (d, 1H, *J* 7.43 Hz, C10-H), 9.53 (s, 1H, N-H exchangeable in D<sub>2</sub>O), 13.79 (s, 1H, C4-OH exchangeable in D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: 13.7 (s, C4'), 20.1 (s, C3'), 29.5 (s, C2'), 42.4 (s, C1'), 100.0 (s, C3), 101.6 (s, C4a), 113.2 (s, C7), 115.1 (s, C10a), 115.9 (s, C<sub>phenyl</sub>), 116.2 (s, C<sub>phenyl</sub>), 124.1 (s, C10), 124.8 (s, C9), 124.8 (s, C<sub>phenyl</sub>), 124.8 (s, C<sub>phenyl</sub>), 132.6 (s, C8), 133.9 (s, C<sub>phenyl</sub>), 134.1 (s, C<sub>phenyl</sub>) 137.9 (s, C6a), 157.0 (s, C10b), 159.3 (s, C4), 159.6 (s, C2), 162.5 (s, C5), 162.9 (s, C=O<sub>benzoyl</sub>). ESI-MS *m/z*: 423.2240 [M+H]<sup>+</sup>, 445.1098 [M+Na]<sup>+</sup>, 446.1020 [M+H+Na]<sup>2+</sup>, 867.3456 [2M+Na]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>19</sub>FN<sub>2</sub>O<sub>5</sub> (422.42): C, 65.40; H, 4.53; N, 6.63. Found C, 65.15; H, 4.43; N, 6.61%.

***N*-(6-*n*-Butyl-4-hydroxy-2,5-dioxo-5,6-dihydro-2H-pyrano[3,2-*c*]quinolin-3-yl)-4-fluorobenzamide (14b).** Pale yellow powder (3.5g, 87%), mp > 300 °C. IR (KBr, cm<sup>-1</sup>): 3432 (O-H), 3266 (NH), 3064 (CH<sub>aromatic</sub>), 2959, 2930, 2859 (CH<sub>aliphatic</sub>), 1748 (C=O<sub>α-pyrone</sub>), 1677 (C=O<sub>quinolone</sub>), 1639 (C=O<sub>arylcarbonyl</sub>) and 1573 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>) δ<sub>H</sub>: 0.92 (t, *J* 8.00 Hz, 3 H, C4'), 1.35 - 1.46 (m, 2 H, C3'), 1.60-1.70 (m, 2 H, C2'), 4.34 (t, *J* 8.00 Hz, 2H, C1'), 7.34 (t, *J* 8.00 Hz, 2 H, C-H<sub>phenyl</sub>), 7.53 (t, 1H, *J* 7.24 Hz, C9-H), 7.83 - 7.93 (m, 2 H, C7-H, C-H<sub>phenyl</sub>), 8.00 - 8.08 (m, 2 H, C8-H, C-H<sub>phenyl</sub>), 7.88 (d, 1 H, *J* 9.80 Hz, C-H<sub>phenyl</sub>) 7.91 (dd, *J* 8.80, 1.20 Hz, 1 H, C-H<sub>phenyl</sub>), 8.15 (dd, *J* 8.02, 1.37 Hz, 1 H, C10-H), 9.70 (s, 1 H, N-H exchangeable in D<sub>2</sub>O), 13.75 (s, 1 H, C4-OH exchangeable in D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>) δ<sub>C</sub>: 14.1 (s, C4'), 19.9 (s, C3'), 29.6 (s, C2'), 42.3 (s, C1'), 99.8 (s, C3), 102.1 (s, C4a), 113.6 (s, C7), 115.7 (s, C10a), 115.9 (s, C<sub>phenyl</sub>), 116.7 (s, C<sub>phenyl</sub>), 124.2 (s, C10), 124.7 (s, C9), 130.8 (s, C8), 130.9 (s, C<sub>phenyl</sub>), 130.9 (s, C<sub>phenyl</sub>), 134.7 (s, C<sub>phenyl</sub>), 138.1 (s, C6a), 157.2 (s, C10b), 159.3 (s, C4), 163.1 (s, C2), 163.3 (s, C5), 163.5 (s, C=O<sub>benzoyl</sub>), 164.8 (s, C-F). ESI-MS *m/z*: 423.2758 [M+H]<sup>+</sup>, 445.0197 [M+Na]<sup>+</sup>, 446.1732 [M+H+Na]<sup>2+</sup>, 867.3784 [2M+Na]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>19</sub>FN<sub>2</sub>O<sub>5</sub> (422.42): C, 65.40; H, 4.53; N, 6.63. Found C, 65.35; H, 4.39; N, 6.60%.

**N-(6-*n*-Butyl-4-hydroxy-2,5-dioxo-5,6-dihydro-2*H*-pyrano[3,2-*c*]quinolin-3-yl)-4-(trifluoromethoxy)**

**benzamide (14c).** Yellow powder (3 g, 64%), mp 255-257 °C. IR (KBr, cm<sup>-1</sup>): 3447 (O-H), 3292 (NH), 3130 (CH<sub>aromatic</sub>), 2958, 2932, 2876 (CH<sub>aliphatic</sub>), 1728 (C=O <sub>$\alpha$ -pyrone</sub>), 1679 (C=O<sub>quinolone</sub>), 1613 (C=O<sub>arylcarbonyl</sub>) and 1576 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, DMSO-*d*<sub>6</sub>)  $\delta$ <sub>H</sub>: 0.91 (t, 1 H, *J* 7.24 Hz, 3 H, C4'), 1.35 - 1.46 (m, 2 H, C3'), 1.59 - 1.69 (m, 2 H, C2'), 4.33 (t, 2 H, *J* 8.00 Hz, C1'), 7.47 - 7.54 (m, 3 H) 7.81 - 7.85 (m, 1 H), 7.86 - 7.91 (m, 1 H), 8.07 - 8.14 (m, 3 H), (s, N-H exchangeable in D<sub>2</sub>O), 13.75 (s, C4-OH exchangeable in D<sub>2</sub>O). <sup>13</sup>C NMR (101 MHz, DMSO-*d*<sub>6</sub>)  $\delta$ <sub>C</sub>: 14.1 (s, C4'), 19.9 (s, C3'), 29.6 (s, C2'), 42.3 (s, C1'), 99.8 (s, C3), 101.9 (s, C4a), 113.5 (s, C7), 116.7 (s, C10a), 119.1 (s, OCF<sub>3</sub>), 121.1 (s, C<sub>phenyl</sub>), 121.6 (s, C<sub>phenyl</sub>), 124.1 (s, C10), 124.7 (s, C9), 130.5 (s, 2C<sub>phenyl</sub>), 133.1 (s, C<sub>phenyl</sub>), 134.7 (s, C8), 138.1 (s, C6a), 151.0 (s, C<sub>phenyl</sub>), 157.2 (s, C10b), 159.2 (s, C4), 163.0 (s, C2), 163.5 (s, C5), 164.7 (s, C=O<sub>benzoyl</sub>). ESI-MS *m/z*: 489.2773 [M+H]<sup>+</sup>, 511.2984 [M+Na]<sup>+</sup>, 977.2922 [2M+H]<sup>+</sup>, 999.3752 [2M+Na]<sup>+</sup>. Anal. Calcd for C<sub>24</sub>H<sub>19</sub>F<sub>3</sub>N<sub>2</sub>O<sub>6</sub> (488.42): C, 59.02; H, 3.92; N, 5.74. Found C, 58.97; H, 3.89; N, 5.67%.

**General procedure for preparation of oxazolopyranoquinolinone derivatives 15a-c.** A mixture of compound **14a-c** (10 mmol) and glacial AcOH (15 mL) with Ac<sub>2</sub>O (10 mL) was heated under reflux for 12 h. During this time the targeted compounds **15a-c** were precipitated. After completion of the reaction (confirmed by TLC), the reaction mixture was cooled to rt. The obtained solid was filtered off, washed with EtOH (3 x 10 mL) and crystallized from AcOH to give the corresponding compound **15**.

**5-*n*-Butyl-2-(2-fluorophenyl)-4*H*-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5*H*)-dione (15a).** White powder (3.4 g, 84%), mp 260-262 °C. IR (KBr, cm<sup>-1</sup>): 3077 (CH<sub>aromatic</sub>), 2952, 2935, 2869 (CH<sub>aliphatic</sub>), 1771 (C=O <sub>$\alpha$ -pyrone</sub>), 1662 (C=O<sub>quinolone</sub>), 1600 (C=N) and 1560 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ <sub>H</sub>: 1.00 (t, *J* 7.60 Hz, 3 H, C4'), 1.46-1.58 (m, 2 H, C3'), 1.71-1.85 (m, 2 H, C2'), 4.37 (t, *J* 8.00 Hz, 2 H, C1'), 7.2-7.27 (m, 1 H, C-H<sub>phenyl</sub>), 7.28 - 7.33 (m, 1 H, C-H<sub>phenyl</sub>), 7.37 (t, 1 H, *J* 7.63 Hz, C9-H), 7.44 (d, 1 H, *J* 8.61 Hz, C7-H), 7.51 - 7.58 (m, 1 H, C-H<sub>phenyl</sub>), 7.68 - 7.74 (m, 1 H, C8-H), 8.28 (dd, 1 H, *J* 7.60, 1.60 Hz, C-H<sub>phenyl</sub>), 8.32 (dd, 1 H, *J* 8.02, 1.76 Hz, C10-H), <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$ <sub>C</sub>: 13.8 (s, C4'), 20.2 (s, C3'), 29.5 (s, C2'), 42.6 (s, C1'), 100.8 (s, C4a), 113.4 (s, C10a), 113.9 (s, C7), 114.0 (s, C<sub>phenyl</sub>), 114.8 (s, C<sub>phenyl</sub>), 117.0 (s, C<sub>phenyl</sub>), 117.2 (s, C10), 123.1 (s, C9), 124.5 (s, C3), 124.7 (s, C<sub>phenyl</sub>), 130.5 (s, C<sub>phenyl</sub>), 133.6 (s, C4), 133.8 (s, C8), 138.9 (s, C6a), 154.5 (s, C10b), 156.5 (s, CF), 157.3 (s, C2<sub>oxazol</sub>), 159.3 (s, C2), 161.9 (s, C5). ESI-MS *m/z*: 405.1330 [M+H]<sup>+</sup>, 406.1340 [M+2H]<sup>2+</sup>, 831.3452 [2M+Na]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>17</sub>FN<sub>2</sub>O<sub>4</sub> (404.40): C, 68.31; H, 4.24; N, 6.93. Found C, 68.28; H, 4.09; N, 6.88 %.

**5-*n*-Butyl-2-(4-fluorophenyl)-4*H*-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5*H*)-dione (15b).** White powder (3.7 g, 91%), mp 236-238 °C. IR (KBr, cm<sup>-1</sup>): 3080 (CH<sub>aromatic</sub>), 2955, 2932, 2869 (CH<sub>aliphatic</sub>), 1788 (C=O <sub>$\alpha$ -pyrone</sub>), 1669 (C=O<sub>quinolone</sub>), 1600 (C=N) and 1563 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ <sub>H</sub>: 1.00 (t, *J* 7.60 Hz, 3 H, C4'), 1.47 - 1.57 (m, 2 H, C3'), 1.73 - 1.82 (m, 2 H, C2'), 4.36 (t, *J* 7.80 Hz, 2 H, C1'), 7.20 (t, 1 H, *J* 8.80 Hz, C9-H), 7.36 (dd, 1 H, *J* 7.70, 1.20 Hz, C-H<sub>phenyl</sub>), 7.40 (dd, 1 H, *J* 6.80, 0.80 Hz, C-H<sub>phenyl</sub>), 7.44 (d, 1 H, *J* 8.61 Hz, C7-H), 7.46 (d, 1 H, *J* 5.87 Hz, C-H<sub>phenyl</sub>), 7.72 (t, 1 H, *J* 7.60 Hz, C8-H), 8.31 (dd, 1 H, *J* 5.20, 2.00 Hz, C-H<sub>phenyl</sub>), 8.34 (dd, 1 H, *J* 8.00, 1.60 Hz, C10-H). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>)  $\delta$ <sub>C</sub>: 14.4 (s, C4'), 20.3 (s, C3'), 29.6 (s, C2'), 42.5 (s, C1'), 100.9 (s, C4a), 113.5 (s, C10a), 114.8 (s, C7), 116.28 (s, C<sub>phenyl</sub>), 116.5 (s, C<sub>phenyl</sub>), 123.1 (s, C3), 124.7 (s, C10), 125.9 (s, C9), 130.1 (s, C8), 133.4 (s, C<sub>phenyl</sub>), 133.6 (s, C<sub>phenyl</sub>), 138.8 (s, C<sub>phenyl</sub>), 138.9 (s, C6a), 154.8 (s, C4), 156.0 (s, C10b), 156.6 (s, C2<sub>oxazol</sub>), 156.9 (s, C2), 163.8 (s, C5), 164.7 (s, CF). ESI-MS *m/z*: 405.2520 [M+H]<sup>+</sup>, 831.3457 [2M+Na]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>17</sub>FN<sub>2</sub>O<sub>4</sub> (404.40): C, 68.31; H, 4.24; N, 6.93. Found C, 68.22; H, 4.15; N, 6.70%.

**5-*n*-Butyl-2-(4-(trifluoromethoxy)phenyl)-4*H*-oxazolo[5',4':4,5]pyrano[3,2-*c*]quinoline-4,11(5*H*)-dione. (15c).** White powder (4.4 g, 94%), mp 245-247 °C. IR (KBr, cm<sup>-1</sup>): 3084 (CH<sub>aromatic</sub>), 2958, 2932, 2879 (CH<sub>aliphatic</sub>), 1755 (C=O <sub>$\alpha$ -pyrone</sub>), 1666 (C=O<sub>quinolone</sub>), 1596 (C=N) and 1560 (C=C<sub>aromatic</sub>). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$ <sub>H</sub>: 1.02 (t, *J* 7.43 Hz, 3 H, C4'), 1.48 - 1.60 (m, 2 H, C3'), 1.73 - 1.85 (m, 2 H, C2'), 4.38 (t, *J* 8.00 Hz, 2 H, C1'), 7.36 (t, 1 H, *J* 8.00

Hz, C9-H), 7.40 (d, *J* 7.20 Hz, 2 H, C-H<sub>phenyl</sub>), 7.46 (d, 1 H, *J* 8.61 Hz, C7-H), 7.73 (t, 1 H, *J* 7.60 Hz, C8-H), 8.34 (dd, *J* 7.6, 1.2 Hz, 2H, C-H<sub>phenyl</sub>), 8.37 (dd, 1 H, *J* 8.00, 1.6 Hz, C10-H). <sup>13</sup>C NMR (101 MHz, CDCl<sub>3</sub>) δ<sub>C</sub>: 13.8 (s, C4'), 20.3 (s, C3'), 29.5 (s, C2'), 42.6 (s, C1'), 100.8 (s, C4a), 113.4 (s, C10a), 114.9 (s, C7), 121.1 (s, C<sub>phenyl</sub>), 123.2 (s, C10), 123.9 (s, C9), 124.7 (s, C<sub>phenyl</sub>), 125.9 (s, C3), 129.5 (s, C<sub>phenyl</sub>), 133.6 (s, C-F3), 138.9 (s, C6a), 151.9 (s, C4), 154.7 (s, C<sub>phenyl</sub>), 155.9 (s, C10b), 156.6 (s, C<sub>phenyl</sub>), 157.3 (s, C2<sub>oxazol</sub>), 160.8 (s, C2), 162.8 (s, C5). ESI-MS *m/z*: 471.1 [M+H]<sup>+</sup>, 493.2 [M+Na]<sup>+</sup>. Anal. Calcd for C<sub>24</sub>H<sub>17</sub>F<sub>3</sub>N<sub>2</sub>O<sub>5</sub> (470.41): C, 61.28; H, 3.64; N, 5.96. Found C, 61.20; H, 3.57; N, 5.89%.

## References

- Anniyappan, M.; Muralidhran, D.; Perumal, P. T. *Tetrahedron Lett.* **2003**, *44*, 3653.  
[https://doi.org/10.1016/S0040-4039\(03\)00707-X](https://doi.org/10.1016/S0040-4039(03)00707-X)
- Ravindranath, N.; Ramesh, C.; Reddy, M. R.; Das, B. *Chem. Lett.* **2003**, *32*, 222.  
<https://doi.org/10.1246/cl.2003.222>
- Arasakumar, T.; Mathusalini, S.; Gopalan, S.; Shyamsivappan, S.; Ata, A.; Mohan, P. S. *Bioorg. Med. Chem. Lett.* **2017**, 1538.  
<https://doi.org/10.1016/j.bmcl.2017.02.042>
- Tomassoli, I.; Herlem, G.; Picaud, F.; Benchekroun, M.; Bautista-Aguilera, O. M.; Luzet, V.; Jimeno, M.-L.; Gharbi, T.; Refouvelet, B.; Ismaili, L. *Monatsh. Chem.* **2016**, *147*, 1069.  
<https://doi.org/10.1007/s00706-016-1660-7>
- Ramesh, E.; Manian, R. S.; Raghunathan, R.; Sainath, S.; Raghunathan, M. *Bioorg. Med. Chem.* **2009**, *17*, 660.  
<https://doi.org/10.1016/j.bmc.2008.11.058>
- Anniyappan, M.; Muralidhran, D.; Perumal, P. T. *Tetrahedron Lett.* **2003**, *44*, 3653.  
[https://doi.org/10.1016/S0040-4039\(03\)00707-X](https://doi.org/10.1016/S0040-4039(03)00707-X)
- El-Agrody, A. M.; Abd-Rabboh, H. S. M.; Al-Ghamd A. M. *Med. Chem. Res.* **2013**, *22*, 1339.  
<https://doi.org/10.1007/s00044-012-0142-7>
- Magedov I.V.; Manpadi M.; Ogasawara M. A.; Dhawan A.S. Rodelj S.; Van slambrouck S.; Steelant W. F. A.; Evdokimov N. M.; Uglinskii P. Y.; Elias E. M.; Knee E. J.; Tongwa P.; Antipin M. Y.; Kornienko A. *J. Med. Chem.* **2008**, *51*, 2561.  
<https://doi.org/10.1021/jm701499n>
- Giddens, A. C.; Boshoff, H. I. M.; Franzblau, S. G.; Barry, C. E.; Copp, B. R. *Tetrahedron Lett.* **2005**, *46*, 7355.  
<https://doi.org/10.1016/j.tetlet.2005.08.119>
- Zhao, Q.; Liu, S.; Li, Y.; Wang, Q.; *J. Agric. Food Chem.*, **2009**, *57*, 2849.  
<https://doi.org/10.1021/1f803632t>
- Chen, L.; Ren, Z.; Zhou, X.; Zeng, J.; Zou, J.; Li, Y. *Appl. Microbiol. Biotechnol.* **2016**, *100*, 857.  
<https://doi.org/10.1007/s00253-015-7092-1>
- Gradillas, A.; Lopez, B.; Brana, Sancho, I.; Perez-Garkia, C.; Alguacil, L. F. *Arzneim.-Forsch./Drug Res.* **2005**, *55*, 725.
- Zhang, F.; Chapman, K. T.; Schleif, W.A.; Olsen, D. B.; Stahlhut, M.; Rutkowski, C. A.; Kuo, L. C.; Jin, L.; Lin, J. H.; Emini, E. A.; Tata, J. R. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 2573.  
<https://doi.org/10.1016/S0960-894X0300474-8>

14. Minhas, G. S.; Pilch, D. S.; Kerrigan, J. E.; La Voie, E. J.; Rice, J. E. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 3891.  
<https://doi.org/10.1016/j.bmcl.2006.05.038>
15. Lipunova, G. N.; Nosova, E. V.; Charushin, V. N.; Chupakhin, O. N. *J. Fluorine Chem.* **2015**, *175*, 84.  
<https://doi.org/10.1016/j.jfluchem.2015.03.011>
16. Xia, Y.; Yang, Z. Y.; Xia, P.; Hackl, T.; Hamel, E.; Mauger, A.; Wu, J. H.; Lee, K. H., *J. Med. Chem.* **2001**, *44*, 3932.  
<https://doi.org/10.1021/jm0101085>
17. Al-Trawneh, S. A.; Zahra, J. A.; Kamal, M. R.; El-Abadelah, M. M.; Zani, F.; Incerti, M.; Cavazzoni, A.; Alfieri, R. R.; Petronini, P. G.; Vicini, P. *Bioorg. Med. Chem.* **2010**, *18*, 5873.  
<https://doi.org/10.1016/j.bmc.2010.06.098>
18. Nosova, E. V.; Lipunova, G. N.; Charushin, V. N.; Chupakhin, O. N. *J. Fluorine Chem.* **2010**, *131*, 1267.  
<https://doi.org/10.1016/j.jfluchem.2010.09.007>
19. Ibrahim, M. A.; Hassanin, H. M. *Arkivoc* **2013** (iv) 217.  
<https://doi.org/10.1016/j.jfluchem.2010.09.007>
20. Hassanin, H. M. *Arkivoc* **2012**, (vi), 384.  
<https://doi.org/10.1016/j.jfluchem.2010.09.007>
21. Liu, Y.; Lu, Y.; Prashad, M.; Repič, O.; Blacklock, T. J. *Adv. Synth. Catal.* **2005**, *347*, 217.  
<https://doi.org/10.1002/adsc.200404236>
22. Sadavarte, V. S.; Swami, S. S.; Desai, D. G. *Synth. Commun.* **1998**, *28*, 1139.  
<https://doi.org/10.1080/00397919808005954>
23. Jampilek, J.; Musiol, R.; Pesko, M.; Kralova, K.; Vejsova, M.; Carroll, J.; Coffey, A.; Finster, J.; Tabak, D.; Niedbala, H.; Kozik, V.; Polanski, J.; Csollei, J.; Dohnal, J. *Molecules*, **2009**, *14*, 1145.  
<https://doi.org/10.3390/molecules14031145>
24. Kappe, T.; Aigner, R.; Hohengassner, P.; Stadlbauer, W. *J. Prakt. Chem.* **1994**, *336*, 596.  
<https://doi.org/10.1002/prac.19943360707>