

Supplementary Material

Generation and reactions of thiirenium ions by the cation pool method

Akihiro Shimizu, Shun Horiuchi, Ryutaro Hayashi, Kouichi Matsumoto, Yu Miyamoto,
Yusuke Morisawa, Tomonari Wakabayashi, and Jun-ichi Yoshida

*Department of Synthetic Chemistry and Biological Chemistry, Graduate School of Engineering,
Kyoto University, Nishikyo-ku, Kyoto 615-8510, Japan*

*Department of Chemistry, School of Science and Engineering, Kindai University, Kowakae 3-4-1,
Higashi-Osaka, Osaka 577-8502, Japan*

E-mail: yoshida@sbchem.kyoto-u.ac.jp

Table of Contents

| | |
|---|----|
| Figure S1. H-Type divided cell used for electrolysis | S2 |
| Figure S2. The ^1H - ^{13}C HMQC spectrum of thionium ion 1a at $-78\text{ }^\circ\text{C}$ | S3 |
| Figure S3. The ^1H - ^{13}C HMBC spectrum of thionium ion 1a at $-78\text{ }^\circ\text{C}$ | S3 |
| Figure S4. Differential NOE spectrum of thionium ion 1a at $-78\text{ }^\circ\text{C}$ | S4 |
| Figure S5. CSI-MA spectrum of 1a at $0\text{ }^\circ\text{C}$ | S5 |
| Figure S6. Raman spectra of a) 4-octyne and b) <i>cis</i> -4-octene | S6 |
| Figure S7. Vibrational modes of a) 4-octyne, b) 1a' , and c) 4-octene | S6 |
| DFT Calculations | S7 |
| NMR Spectra of compounds 2e , 2g , 4 , 2i , 3a , 3b , 2x | S9 |




Figure S1. An H-type divided cell for electrolysis.




Figure S2. The ^1H - ^{13}C HMQC spectrum of thionium ion **1a** at -78 °C.




Figure S3. The ^1H - ^{13}C HMBC spectrum of thionium ion **1a** at -78 °C.




Figure S4. Differential NOE spectrum of thionium ion **1a** at $-78\text{ }^{\circ}\text{C}$.




Figure S5. CSI-MA spectrum of 1a at 0 °C




Figure S6. Raman spectra of a) 4-octyne and b) *cis*-4-octene.




Figure S7. Vibrational modes of a) 4-octyne, b) **1a'**, and c) 4-octene.

DFT Calculations

Table S1. Cartesian coordinates (Å) of the optimized structure of thiirenium ion **1a** calculated at the RB3LYP/6-31G(d) level of theory.

| Atom | X | Y | Z | Atom | X | Y | Z |
|------|----------|----------|----------|------|----------|----------|----------|
| C | - | - | 0.646589 | H | - | - | 1.858382 |
| C | - | - | - | H | 2.335588 | 2.436115 | 0.000000 |
| C | - | - | - | H | 1.566452 | 4.791052 | 0.000000 |
| C | - | - | 1.998522 | H | - | 3.469844 | 0.000000 |
| C | 0.317094 | 1.640930 | 0.000000 | H | - | 1.109948 | 0.000000 |
| C | 1.272328 | 2.661890 | 0.000000 | H | - | 5.309054 | 0.000000 |
| C | 0.836396 | 3.987904 | 0.000000 | H | 1.217552 | - | 2.950179 |
| C | - | 4.276278 | 0.000000 | H | 0.830982 | - | 2.320267 |
| C | - | 3.241975 | 0.000000 | H | - | - | 4.790296 |
| C | - | 1.911153 | 0.000000 | H | - | - | 4.162394 |
| C | 0.410552 | - | 2.848969 | H | 0.830982 | - | - |
| C | - | - | 4.233908 | H | 1.217552 | - | - |
| C | 0.410552 | - | - | H | - | - | - |
| C | - | - | - | H | - | - | - |
| H | - | - | - | H | 0.729728 | - | 4.820425 |
| H | - | - | - | H | 0.729728 | - | - |
| H | - | - | 2.512526 | S | 1.000474 | - | 0.000000 |

Table S2. Cartesian coordinates (Å) of the optimized structure of thiirenium ion **1a'** calculated at the RB3LYP/6-31G(d) level of theory.

| Atom | X | Y | Z | Atom | X | Y | Z |
|------|-----------|-----------|-----------|------|-----------|-----------|-----------|
| C | 1.436690 | 0.538666 | 0.647012 | H | 3.043300 | 0.003334 | 1.855844 |
| C | 1.436690 | 0.538666 | -0.647012 | H | -3.008163 | 1.052776 | 0.000000 |
| C | 2.019203 | 0.373660 | -1.997520 | H | -4.689283 | -0.791386 | 0.000000 |
| C | 2.019203 | 0.373660 | 1.997520 | H | -1.537344 | -3.704128 | 0.000000 |
| C | -1.318187 | -0.309980 | 0.000000 | H | 0.171364 | -1.887512 | 0.000000 |
| C | -2.679673 | 0.016803 | 0.000000 | H | 0.988442 | 2.037002 | 2.932232 |
| C | -3.626087 | -1.004700 | 0.000000 | H | 2.577067 | 2.447659 | 2.289430 |
| C | -3.185612 | -2.326086 | 0.000000 | H | 2.073130 | 0.700851 | 4.785479 |
| C | -1.831113 | -2.660107 | 0.000000 | H | 3.673496 | 1.115739 | 4.143474 |
| C | -0.885470 | -1.639257 | 0.000000 | H | 2.577067 | 2.447659 | -2.289430 |
| C | 2.019203 | 1.675004 | 2.831197 | H | 0.988442 | 2.037002 | -2.932232 |
| C | 2.633044 | 1.451321 | 4.216208 | H | 3.673496 | 1.115739 | -4.143474 |
| C | 2.019203 | 1.675004 | -2.831197 | H | 2.073130 | 0.700851 | -4.785479 |
| C | 2.633044 | 1.451321 | -4.216208 | H | 2.622730 | 2.382404 | 4.790223 |
| H | 3.043300 | 0.003334 | -1.855844 | H | 2.622730 | 2.382404 | -4.790223 |
| H | 1.464042 | -0.412456 | -2.525680 | F | -4.088037 | -3.307522 | 0.000000 |
| H | 1.464042 | -0.412456 | 2.525680 | S | -0.198093 | 1.101565 | 0.000000 |




Table S3. Cartesian coordinates (Å) of the optimized structure of 4-octyne calculated at the RB3LYP/6-31G(d) level of theory.


| Atom | X | Y | Z | Atom | X | Y | Z |
|------|-----------|-----------|-----------|------|-----------|-----------|-----------|
| C | -4.425829 | 0.002868 | 0.095326 | H | -2.749892 | 0.998324 | 1.044428 |
| C | -2.976515 | 0.495637 | 0.096088 | H | -2.832709 | 1.244467 | -0.692568 |
| C | -1.964795 | -0.650962 | -0.114768 | H | -2.191040 | -1.160698 | -1.062892 |
| C | -0.570794 | -0.201311 | -0.117759 | H | -2.107507 | -1.405988 | 0.672351 |
| C | 0.570794 | 0.201293 | -0.117787 | H | 2.191049 | 1.160519 | -1.063070 |
| C | 1.964796 | 0.650943 | -0.114861 | H | 2.107502 | 1.406104 | 0.672131 |
| C | 2.976513 | -0.495621 | 0.096200 | H | 2.749884 | -0.998142 | 1.044626 |
| C | 4.425828 | -0.002855 | 0.095361 | H | 2.832712 | -1.244587 | -0.692327 |
| H | -5.125550 | 0.832491 | 0.245683 | H | 5.125547 | -0.832452 | 0.245873 |
| H | -4.681886 | -0.480315 | -0.855713 | H | 4.681895 | 0.480157 | -0.855762 |
| H | -4.598791 | -0.727680 | 0.895267 | H | 4.598786 | 0.727837 | 0.895171 |

Table S4. Cartesian coordinates (Å) of the optimized structure of *cis*-4-octene calculated at the RB3LYP/6-31G(d) level of theory.

| Atom | X | Y | Z | Atom | X | Y | Z |
|------|-----------|-----------|-----------|------|-----------|-----------|-----------|
| C | -2.135846 | 3.034243 | -1.113396 | H | -0.268938 | 2.825071 | -0.032964 |
| C | -1.305409 | 2.471133 | 0.043919 | H | -0.902619 | 0.540766 | -0.852689 |
| C | -1.305409 | 0.930254 | 0.089691 | H | -2.350218 | 0.585768 | 0.142292 |
| C | -0.549416 | 0.382665 | 1.271197 | H | -0.956201 | 0.674215 | 2.241581 |
| C | 0.549416 | -0.382665 | 1.271197 | H | 0.956201 | -0.674215 | 2.241581 |
| C | 1.305409 | -0.930254 | 0.089691 | H | 0.902619 | -0.540766 | -0.852689 |
| C | 1.305409 | -2.471133 | 0.043919 | H | 2.350218 | -0.585768 | 0.142292 |
| C | 2.135846 | -3.034243 | -1.113396 | H | 1.690438 | -2.860493 | 0.996837 |
| H | -2.118214 | 4.130050 | -1.121595 | H | 0.268938 | -2.825071 | -0.032964 |
| H | -1.753715 | 2.688065 | -2.081738 | H | 2.118214 | -4.130050 | -1.121595 |
| H | -3.183913 | 2.718032 | -1.040187 | H | 1.753715 | -2.688065 | -2.081738 |
| H | -1.690438 | 2.860493 | 0.996837 | H | 3.183913 | -2.718032 | -1.040187 |

NMR spectra





2e

NOE spectrum

