

Cyanine–cyanine hybrid structure as a stabilized polyelectrochromic system: synthesis, stabilities, and redox behavior of di(1-azulenyl)methylium units connected with electron-accepting π -electron systems

Shunji Ito,^{*a} Ryuta Sekiguchi,^b Akira Mizushima,^a Kohei Kudo,^a Jun Kawakami,^a and Taku Shoji^b

^aGraduate School of Science and Technology, Hirosaki University, Hirosaki 036-8561, Aomori, Japan

^bDepartment of Chemistry, Faculty School of Science, Shinshu University, Matsumoto 390-8621, Nagano, Japan

Email: itsnj@hirosaki-u.ac.jp

Dedicated to Professor Kenneth K. Laali on the occasion of his 65th birthday


Received 05-10-2017

Accepted 09-23-2017

Published on line 11-26-2017

Abstract

This paper describes preparation of the two-types of cations composed of di(1-azulenyl)methylium units based on a new structural principle of cyanine–cyanine (C–C) hybrid for the design of polyelectrochromic materials. Voltammetric analysis of these cations revealed the idealized reversible two-step, two-electron redox properties. Two-step color changes presumed by their C–C hybrid structure were revealed by their electrochemical reduction. The scope of the creation of the stabilized polyelectrochromic materials taking the new structural principle with azulene skeletons is demonstrated by two-types of examples.




Keywords: Azulene, multi-electron redox system, polyelectrochromism, cyanine, voltammetry

Introduction

Electrochromism is observed in reversible redox systems, which exhibit significant color changes in different oxidation states.¹⁻² Stabilization of the redox cycle is very important in construction of potentially useful electrochromic materials, because the molecules utilized for the application require high redox-stability. Construction of organic molecules that contain multiple redox-active chromophores is required for the preparation of novel polyelectrochromic materials, which respond to different potentials with a variety of colors.³

Previously, Hünig et al. proposed a concept of violene–cyanine (V–C) hybrid to produce stabilized organic electrochromic materials.⁴⁻⁷ The hybrid is constructed by the violene-type redox system containing delocalized closed-shell polymethine (e.g., cyanine) dyes as one or two end groups. The hybrid is expected to provide a color of a cyanine dye by an overall two-electron transfer as illustrated by the general structure with a cyanine end group in both terminals (Scheme 1). However, multiple-color changes important for the polyelectrochromic materials will not be established by the hybrid structure.



Scheme 1. General structure of the Hünig's V–C hybrid system. The presumed cyanine substructures generated by the two-electron transfer are represented in bold line in the Scheme.


To produce novel polyelectrochromic materials that exhibit multiple-color changes, we have recently proposed a molecular design from the cyanine-based polymethine dye.⁸⁻⁹ The cyanine dyes, themselves does not possess good electrochemical properties, because of their ionic structures. However, either one or two end groups of the cyanine-based polymethine dye are replaced by another cyanine unit, in that case, two-step redox activities with multiple-color changes should be induced. We call this molecular design cyanine–cyanine (C–C) hybrid. Herein we demonstrate the new structural principle for the polyelectrochromic materials by the preparation of two-types of the C–C hybrid structure composed of di(1-azulenyl)methylum units as a stabilized cyanine end group.¹⁰⁻¹⁷

Results and Discussion


Structural Principle. Electronic nature of the terminal groups constituted the cyanine end group is very important for the design of the C–C hybrid with high redox stability. The hybrid structure in Scheme 2 represents the hybrid constructed by the two cyanine units composed of equal-electronic nature, in which a cyanine substructure should be induced in either reduced or oxidized form.

In contrast, the hybrid structure could be constructed by the two cyanine units with push and pull electronic nature. The general structure illustrated in Scheme 3 represents such an example. In this case, the cyanine structure **A** in Scheme 3 will be transform into structure **C** by two-electron transfer via a neutral radical state **B**. Expected species **A** and **C** in Scheme 3 should represent the closed shell systems. The neutral radical state **B**


may also be stabilized by *capto-dative* substitution effect in common with those of violen-type electrochromics, from which higher persistency would be expected.¹⁸⁻²⁰ Therefore, the three-colored states will be stabilized by the C–C hybrid constructed by two cyanine units with push and pull electronic nature represented in Scheme 3. Di(1-azulenyl)(6-azulenyl)methylm ions (**1a-c⁺**) would exemplify this type of the C–C hybrid, in which cyanine substructure should be induced in both reduced and oxidized forms (Scheme 4).²¹



Scheme 2. General structure of the C–C hybrid prepared by three terminal groups (X) with same electronic nature. The presumed cyanine substructures generated by the two-electron transfer are represented in bold line in the Scheme.




Scheme 3. General structure of the C–C hybrid prepared by three terminal groups (X and Y) with push and pull electronic nature. The presumed cyanine substructures generated by the two-electron transfer are represented in bold line in the Scheme.



Scheme 4. An example of the C–C hybrid prepared by the three terminal groups by azulenyl groups.

When the C–C hybrid is constructed by two cyanine end groups with equal-electronic nature, a defect of conjugation will arise in the core cyanine unit in either reduced or oxidized form because the core unit is

constructed by odd number of polymethine chain.⁸⁻⁹ In contrast, the C–C hybrid structure constructed by two cyanine end groups with push and pull electronic nature should avoid such disadvantage in the conjugation in the core cyanine unit as illustrated in Scheme 5. Both the colored species **D** and **F** in Scheme 5 should also represent the closed shell systems. Therefore, the stabilized three-colored states will be achieved by the two-types of the C–C hybrid structure constructed by the cyanine units with push and pull electronic nature illustrated by the general structures in Schemes 3 and 5 with oxidation levels varying from 1+ to 1–.



Scheme 5. General structure of the C–C hybrid prepared by four terminal groups (X and Y) with push and pull electronic nature. The presumed cyanine substructures generated by the two-electron transfer are represented in bold line in the Scheme.

For the construction of the new hybrid system represented the general structures in Schemes 3 and 5, it is very important to select highly colored polymethine end groups with high stability. We selected di(1-azulenyl)methylm units as an end group for the construction of the C–C hybrid due to their high stability and also their strong absorption in visible region owing to the cyanine-like structure as demonstrated by our previous work.¹⁰⁻¹⁷

Two di(1-azulenyl)methylm units with bis(3,6-di-*tert*-butyl and 3-methoxycarbonyl) substituents were applied to examine the substituent effect on the azulene rings (Schemes 6–9). Introduction of the *tert*-butyl group into the azulene rings should increase their thermodynamic stabilities and also reversibility of both electrochemical reduction and oxidation states. 3-Methoxycarbonyl derivatives might increase the stability in their presumed anionic state to decrease the strong electron-donating character of the 1-azulenyl groups.


As the electron-withdrawing core cyanine unit for the preparation of the C–C hybrid illustrated in Scheme 3, we selected 4-nitrophenyl and 5-(4-nitrophenyl)-2-thienyl groups to examine the effect of the core cyanine unit (Schemes 6 and 7). Introduction of 2,5-thiophenediyl spacer might further stabilize the presumed two-electron reduction state by the extension of the π -electron system with the thienoquinoidal form.²²⁻²⁵

To exemplify the C–C hybrid illustrated in Scheme 5, we have examined the preparation of 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propenylium ion that employed 4-nitrophenyl groups as the electron-withdrawing terminal groups. We have also examined the substituent effects at the 1,1-diaryl moiety, i.e., diphenyl, bis(4-fluorophenyl), and fluorenylidene in addition to the bis(4-nitrophenyl) terminal groups (Schemes 8 and 9). Thus, the two-types of C–C hybrids constructed by di(1-azulenyl)methylm units as an end group **5a⁺**, **5b⁺**, **8a⁺**, **8b⁺**, **11a–d⁺**, and **14⁺** (Schemes 6–9) were designed as the new examples for the C–C hybrid with the general structure in Schemes 3 and 5.


Synthesis. Synthesis of **5a⁺**, **5b⁺**, **8a⁺**, **8b⁺**, **11a–d⁺**, and **14⁺** constructed by di(1-azulenyl)methylm units was established by the hydride abstraction from the corresponding hydro derivatives **4a**, **4b**, **7a**, **7b**, **10a–d**, and **13** with DDQ and obtained as hexafluorophosphates by the treatment with hexafluorophosphoric acid as illustrated in Schemes 6–9.

The reaction of 1,6-di-*tert*-butylazulene (**3a**)¹⁰ with 4-nitrobenzaldehyde (**2**) in acetic acid at room temperature afforded bis(3,6-di-*tert*-butyl-1-azulenyl)(4-nitrophenyl)methane (**4a**) in 76% yield (Scheme 6). Similarly, bis(3,6-di-*tert*-butyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (**7a**) was obtained by the reaction of **3a** with 5-(4-nitrophenyl)thiophene-2-carbaldehyde (**6**),²⁶ which was prepared by Miyaura-Suzuki cross-coupling of 2-formylthiophene-5-boronic acid with 4-iodonitrobenzene, in acetic acid at room temperature in 62% yield (Scheme 7).

Analogously, preparation of the corresponding bis(3-methoxycarbonyl) derivatives **4b** and **7b** for the precursors of **5b**⁺ and **8b**⁺ was accomplished via the reaction of methyl azulene-1-carboxylate (**3b**) with **2** and **6** in refluxing acetic acid 84% and 11% yields, respectively. The yield of **7b** has been further improved to 35% by performing the reaction in a boiling mixture of acetic acid and toluene, probably due to the increasing solubility of **6** in the mixed solvent.



Scheme 6. Preparation of **5a**⁺ PF_6^- and **5b**⁺ PF_6^- .




Scheme 7. Preparation of **8a**⁺ PF_6^- and **8b**⁺ PF_6^- .


Furthermore, reaction of **3a** with β -phenylcinnamaldehyde (**9a**) in acetic acid at room temperature afforded 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-diphenylpropene (**10a**) in 76% yield (Scheme 8). Preparation

of the corresponding bis(3-methoxycarbonyl) derivative **10b** for the precursor of **11b**⁺ was accomplished by the similar reaction of **3b** with **9a** in refluxing acetic acid in 27% yield.

As similar to the reaction with **9a**, 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-fluorophenyl)propene (**10b**) was obtained in 43% yield by the reaction of **3a** with 3,3-bis(4-fluorophenyl)-2-propenal (**9b**), which was prepared from 4,4'-difluorobenzophenone by the reaction with Et₃N in the presence of TiCl₄ followed by the hydration with saturated NH₄Cl solution as reported in the literature.²⁷ Preparation of the 4,4'-dinitro derivative, 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propene (**10d**), for the precursor of **11d**⁺ was established in 30% yield by the similar reaction of **3a** with 3,3-bis(4-nitrophenyl)-2-propenal (**9c**), which was prepared in 27% yield by the reaction of 4,4'-dinitrobenzophenone²⁸ with Et₃N in the presence of TiCl₄ followed by the hydration with saturated NH₄Cl solution by utilizing the preparation method for **9b**. 9-[2,2-Bis(3,6-di-*tert*-butyl-1-azulenyl)ethylidene]-9*H*-fluorene (**13**) was also obtained by the reaction of **3a** with 9*H*-fluoren-9-ylidene acetaldehyde (**12**)²⁷ in acetic acid at room temperature in 64% yield (Scheme 9).



Scheme 8. Preparation of **11a-d**⁺PF₆⁻.




Scheme 9. Preparation of **14**⁺PF₆⁻.

Synthesis of **5a⁺**, **5b⁺**, **8a⁺**, **8b⁺**, **11a–d⁺**, and **14⁺** was established by the hydride abstraction from **4a**, **4b**, **7a**, **7b**, **10a–d**, and **13** with DDQ at room temperature and obtained as hexafluorophosphates by the treatment with hexafluorophosphoric acid. The reaction of **4a**, **4b**, **7a**, and **7b** with DDQ yielded **5a⁺** (58%), **5b⁺** (74%), **8a⁺** (77%), and **8b⁺** (33%) as a hexafluorophosphate (Schemes 6 and 7). Likewise, the oxidation of **10a**, **10c**, and **13** with DDQ afforded the corresponding hexafluorophosphates **11a⁺PF₆⁻**, **11c⁺PF₆⁻**, and **14⁺PF₆⁻** in 74%, 95%, and 88% yields, respectively (Schemes 8 and 9). The hexafluorophosphates **11b⁺PF₆⁻** and **11d⁺PF₆⁻** were obtained by the hydride abstraction from **10b** and **10d** with DDQ in the two-phase reaction of CH₂Cl₂ and HPF₆ solution at room temperature in 92% and 66% yields, respectively. These new salts are stable, deep-colored crystals, and are storable in the crystalline state in spite of their cationic structure with electron-withdrawing nitro group as an end group in some cases.

Spectroscopic Properties. The two-types of C–C hybrids **5a⁺**, **5b⁺**, **8a⁺**, **8b⁺**, **11a–d⁺**, and **14⁺** were fully characterized by the spectral data as shown in the Experimental Section. Mass spectra of the cation salts **5a⁺PF₆⁻**, **5b⁺PF₆⁻**, **8a⁺PF₆⁻**, **8b⁺PF₆⁻**, **11a–d⁺PF₆⁻**, and **14⁺PF₆⁻** ionized by ESI showed the correct (M – PF₆)⁺ ion peaks, which indicated the cationic structures of these products. The characteristic bands of the hexafluorophosphates were observed at 839–842 (strong) and 557–558 (medium) cm⁻¹ in their IR spectra, which also supported the ionic structure of these compounds.

UV–vis spectra of these cations in acetonitrile are shown in Figures 1 and 2. As expected by their cyanine-like substructures, these cations showed characteristic charge-transfer absorption bands in the visible region. Their absorption maxima (nm) and coefficients (log ε) with those of di(1-azulenyl)phenylmethylium ions **15a⁺** and **15b⁺**¹⁰ without the nitro substituent are summarized in Table 1. The longest wavelength absorption of **5a⁺** (704 nm) and **5b⁺** (637 nm) showed bathochromic shift by 23 nm and 21 nm, respectively, compared with that of **15a⁺** (681 nm) and **15b⁺** (616 nm). However, **8a⁺** and **8b⁺** inserted with thienylene spacer for the extension of π-electron system exhibited a slight hypsochromic shift as compared with that of **5a⁺** and **5b⁺**. Characteristic feature of the UV–vis spectra for **8a⁺** (540 nm and 702 nm) and **8b⁺** (536 nm and 630 nm) was represented by two absorption bands, although that for **5a⁺** and **5b⁺** was characterized by a strong absorption band in this region.



The UV–vis spectra of **11a⁺** and **11b⁺** in visible region were characterized by two strong absorption bands, whereas, **5a⁺** and **5b⁺** exhibited an absorption band in this region. The longest wavelength absorption of **11a⁺** (722 nm) and **11b⁺** (655 nm) showed bathochromic shift by 41 nm and 39 nm, respectively, compared with that for **15a⁺** and **15b⁺**. Electron-withdrawing substituents on the terminal phenyl groups showed a bathochromic shift by, 2 nm in the fluoro derivative **11c⁺**, 14 nm in nitro derivative **11d⁺**, and 7 nm in the

fluorenilidene derivative **14**⁺ compared with that of **11a**⁺. As expected, 3-methoxycarbonyl-1-azulenyl groups exhibited hypsochromic shift in the longest wavelength absorption band compared with that of *tert*-butyl derivative **11a**⁺. The extinction coefficients of these cations **11a–d**⁺ and **14**⁺ are approximately as large as those of cations **5a**⁺, **5b**⁺, **8a**⁺, and **8b**⁺.




Figure 1. UV–vis Spectra: (a) **5a**⁺ (black line) and **5b**⁺ (red line); (b) **8a**⁺ (black line) and **8b**⁺ (red line) in acetonitrile.




Figure 2. UV–vis Spectra: (a) **11a**⁺ (black line) and **11b**⁺ (red line); (b) **11c**⁺ (black line), **11d**⁺ (blue line), and **14**⁺ (red line) in acetonitrile.

Table 1. Longest wavelength absorption maxima (nm) and their coefficients of **5a**⁺, **5b**⁺, **8a**⁺, **8b**⁺, **11a–d**⁺, **14**⁺, **15a**⁺, and **15b**⁺ in acetonitrile

Sample	λ_{\max} (log ϵ)	Sample	λ_{\max} (log ϵ)	Ref.
5a ⁺	704 (4.79)	11c ⁺	535 (4.23)	
			724 (4.63)	
5b ⁺	637 (4.63)	11d ⁺	507 (3.85)	
			736 (4.35)	
8a ⁺	540 (4.55)	14 ⁺	544 (4.15)	
	702 (4.69)		729 (4.49)	

Table 1. Continued

Sample	λ_{\max} (log ϵ)	Sample	λ_{\max} (log ϵ)	Ref.
8b⁺	536 (4.57)	15a⁺	681 (4.61)	10
	630 (4.50)			
11a⁺	534 (4.18)	15b⁺	616 (4.60)	10
	722 (4.58)			
11b⁺	543 (4.27)			
	655 (4.40)			

Thermodynamic stability. As a measure of the thermodynamic stability, the pK_R^+ value of these cations was determined spectrophotomerically in a buffer solution prepared in 50% aqueous acetonitrile. The K_R^+ scale is defined by the equilibrium constant for the reaction of a carbocation with a water molecule ($K_R^+ = [\text{ROH}][\text{H}_3\text{O}^+]/[\text{R}^+]$). Therefore, the larger pK_R^+ index value ($pK_R^+ = -\log K_R^+$) indicates higher stability of the carbocation. However, the neutralization of these cations was not completely reversible attributable to the instability of the neutralized products under the highly basic conditions for the pK_R^+ measurement. After the measurement, acidification of the alkaline solutions with HCl regenerated the characteristic absorption in the visible region in 1–89% (Table 2). Therefore, the reported values correspond to the decomposition point started by the reaction with water molecule. The values are summarized in Table 2.

Table 2. pK_R^+ values^a and redox potentials^b of **5a⁺**, **5b⁺**, **8a⁺**, **8b⁺**, **11a–d⁺**, **14⁺**, **15a⁺**, and **15b⁺**

Sample	pK_R^+	E_1^{red}	E_2^{red}	E_1^{ox}	E_2^{ox}	$E_1^{\text{red}} - E_2^{\text{red}}$	Ref.
5a⁺	9.5 ± 0.1 (1%)	-0.59	-0.94	0.94	(1.44)	0.35	
		(-0.55) ^c	(-0.92) ^c	(0.92) ^c	(1.41) ^c		
5b⁺	1.4 ± 0.1 (89%)	-0.31	-0.79	(1.38)		0.48	
		(-0.28) ^c	(-0.76) ^c	(1.35) ^c			
8a⁺	10.9 ± 0.1 (3%)	-0.63	-1.08	0.95	(1.33)	0.45	
		(-0.61) ^c	(-1.06) ^c	(0.93) ^c	(1.29) ^c		
8b⁺	2.9 ± 0.1 (83%)	-0.36	-0.99	(1.43)	(1.76)	0.63	
		(-0.34) ^c	(-0.96) ^c	(1.36) ^c	(1.68) ^c		
11a⁺	12.1 ± 0.1 (5%)	-0.67	(-1.48)	0.84	(1.40)	0.81	
		(-0.64) ^c	(-1.45) ^c	(0.82) ^c	(1.30) ^c		
11b⁺	5.3 ± 0.1 (60%)	-0.37	(-1.19)	(1.27)	(1.62)	0.82	
		(-0.34) ^c	(-1.15) ^c	(1.24) ^c	(1.59) ^c		
11c⁺	11.7 ± 0.1 (5%)	-0.65	(-1.54)	0.85		0.89	
		(-0.62) ^c	(-1.42) ^c	(0.82) ^c	(1.41) ^c		
11d⁺	8.7 ± 0.1 (1%)	-0.50	-0.86	0.88		0.36	
		(-0.48) ^c	(-0.84) ^c	(0.86) ^c	(1.38) ^c		
14⁺	9.3 ± 0.1 (10%)	-0.49	(-1.12)	0.93	(1.48)	0.63	
		(-0.47) ^c	(-1.08) ^c	(0.90) ^c	(1.41) ^c		

Table 2. Continued

Sample	pK_R^+	E_1^{red}	E_2^{red}	E_1^{ox}	E_2^{ox}	$E_1^{\text{red}} - E_2^{\text{red}}$	Ref.
15a ^d	12.4	-0.78	(-1.64)	0.88	(1.38)	0.86	10
15b ^d	3.4	(-0.49) ^e	(-1.31)	(1.32)		0.82	10

^a The pK_R^+ values were determined spectrophotometrically in a buffered solution prepared in 50% aqueous acetonitrile.¹⁰ Regenerated absorption maxima (%) of the cations in visible region by acidification of the alkaline solution with HCl are shown in parentheses. ^b Redox potentials were measured by CV and DPV [V vs. Ag/AgNO₃, 1 mM in benzonitrile containing Et₄NClO₄ (0.1 M), Pt electrode (ID: 1.6 mm), scan rate 100 mV s⁻¹, and Fc/Fc⁺ = 0.15 V]. ^c The peak potentials measured by DPV are shown in parentheses. ^d Potentials were the value measured in acetonitrile. ^e The wave was accompanied by second peak at -0.57 V probably due to the deposition of reduced species on the electrode in acetonitrile.

Cation **5a**⁺ (pK_R^+ = 9.5) is not strongly destabilized by the 4-nitro substituent compared with that of **15a**⁺ (pK_R^+ = 12.4),¹⁰ in spite of the existence of the strong electron-withdrawing nitro substituent as an end group. Bis(3-methoxycarbonyl) derivative **5b**⁺ (pK_R^+ = 1.4) exhibited significant destabilization due to the substitution of an electron-withdrawing methoxycarbonyl group at both 1-azulenyl groups, but still exhibited reasonably high pK_R^+ value. Cations **8a**⁺ and **8b**⁺ is slightly stabilized by the insertion of thienylene spacer, since **8a**⁺ (pK_R^+ = 10.9) and **8b**⁺ (pK_R^+ = 2.9) exhibited slightly larger pK_R^+ value, compared with that of **5a**⁺ and **5b**⁺.

The pK_R^+ value of **11a**⁺ (pK_R^+ = 12.1) and **11b**⁺ (pK_R^+ = 5.3) is almost equal to that of **15a**⁺ and **15b**⁺ (pK_R^+ = 3.4),¹⁰ in spite of the existence of two phenyl substituents as end groups with allylic cationic system. As expected by destabilization by the electron-withdrawing substituents as the end groups, bisfluorophenyl substituted cation **11c**⁺ (pK_R^+ = 11.7), bisnitrophenyl substituted cation **11d**⁺ (pK_R^+ = 8.7), fluorenylidene substituted cation **14**⁺ (pK_R^+ = 9.3) is less stable than **11a**⁺, as similar to the results on the cations between **5a**⁺ and **15a**⁺ owing to the destabilization by the electron-withdrawing groups.

Redox potentials. Subsequently, the redox behavior of **5a**⁺, **5b**⁺, **8a**⁺, **8b**⁺, **11a-d**⁺, and **14**⁺ was examined by cyclic voltammetry (CV) and differential pulse voltammetry (DPV). The first and second redox potentials (in volts vs Ag/AgNO₃) of these cations are summarized in Table 2. The redox potentials observed at higher potentials are summarized in the Supporting Information. The first and second reduction waves of **5a**⁺, **8a**⁺, **11a**⁺, and **11d**⁺ are shown in Figures 3 and 4 and those of the others are summarized in the Supporting Information.

As it is shown in Table 2, the characteristic feature of the cations with electron-accepting nitrophenyl groups was small $E_1^{\text{red}} - E_2^{\text{red}}$ values, compared with those without the electron-withdrawing group. Indeed, **5a**⁺ showed a reversible two-step, two-electron reduction wave at -0.59 V and -0.94 V on CV due to the formation of a radical and an anionic species. The first reduction potentials of **5a**⁺ are slightly less negative compared with that of **15a**⁺; this indicates the electrochemical destabilization of the methyl cation by the 4-nitro substituent as similar to the results on the pK_R^+ measurements. The less negative second reduction potential of **5a**⁺ by 0.70 V, compared with that of **15a**⁺, corresponds to the stabilization of anionic state by the 4-nitro substituent. The potential difference between the first and the second reduction waves (0.35 V) is significantly small as compared with that of **15a**⁺ (0.86 V). These results should correspond to the characteristic features of the C-C hybrid structure. The electrochemical oxidation of **5a**⁺ showed a reversible

wave at 0.94 V on CV, due to the oxidation of an azulene ring to give a dication radical. The oxidation potential is almost equal to that of **15a**⁺ ($E_1^{\text{ox}} = 0.88$ V).¹⁰

The first reduction potential of bis(3-methoxycarbonyl) derivative **5b**⁺ is slightly less negative compared with that of **5a**⁺, as similar to the results on those of cations **15a**⁺ and **15b**⁺ ($E_1^{\text{red}} = -0.49$ V).¹⁰ The oxidation of **5b**⁺ also exhibited a wave at 1.38 V on CV ascribed to the oxidation of an azulene ring to generate a dicationic species, but without reversibility. The reduction of **8a**⁺ and **8b**⁺ showed same tendencies for **5a**⁺ and **5b**⁺ except for the slightly more negative reduction potentials as expected by the stabilization by the thienylene insertion.

The electrochemical reduction of **11a**⁺ showed two-step, two-electron reduction wave at -0.67 V and -1.48 V on CV due to the formation of a radical and an anionic species. The reduction potentials of **11b**⁺ are slightly less negative compared with those of **11a**⁺; this indicates the electrochemical destabilization of the methyl cations by the methoxycarbonyl substituents as similar to the results on **5a**⁺ and **5b**⁺.

The characteristic feature of the cations with electron-accepting groups **11c**⁺, **11d**⁺, and **14**⁺ was small $E_1^{\text{red}} - E_2^{\text{red}}$ values, compared with those without the electron-withdrawing group, although difluoro derivative **11c**⁺ showed almost same value with that of **11a**⁺. The less negative second reduction potential of **11d**⁺ by 0.62 V, compared with that of **11a**⁺, corresponds to the increase of electron affinity of the cation by the two 4-nitrophenyl substituents. The high reversibility of the CV waves and small potential difference (0.36 V) of **11d**⁺ indicate the characteristic features of the presumed C-C hybrid structure. The electrochemical oxidation of **11a-c**⁺ and **14**⁺ showed almost similar tendency on CV with those of **5a**⁺, **5b**⁺, **8a**⁺, and **8b**⁺, due to the oxidation of an azulene ring to give a dication radical.




Figure 3. Cyclic voltammograms of (a) reduction of **5a**⁺ (1 mM) and (b) **8a**⁺ (1 mM) in benzonitrile containing Et₄NClO₄ (0.1 M) as a supporting electrolyte; scan rate, 100 mV s⁻¹.




Figure 4. Cyclic voltammograms of (a) reduction of **11a**⁺ (1 mM) and (b) **11d**⁺ (1 mM) in benzonitrile containing Et₄NClO₄ (0.1 M) as a supporting electrolyte; scan rate, 100 mV s⁻¹.

Electrochromic analysis. Visible spectra of the two-types of C–C hybrids were monitored to clarify the color changes under the electrochemical reduction conditions. A constant-current reduction was applied to the solutions of **5a**⁺, **5b**⁺, **8a**⁺, **8b**⁺, **11a–d**⁺, and **14**⁺ with a platinum mesh for working and a wire counter electrodes.

When the visible spectra of **5a**⁺ were monitored in benzonitrile containing Et₄NClO₄ (0.1 M) as a supporting electrolyte at room temperature under the electrochemical reduction conditions, the strong absorption band at 710 nm in the visible region was gradually decreased along with the development of new absorption band at around 590 nm (Figure 5). Further reduction developed a broad absorption band around the new absorption band (525 nm). Thus, the spectral changes should correspond to the two-step, two-electron reduction wave on CV arising from the formation of a radical and an anionic species. The color of the solution of **5a**⁺ gradually changed from dark green to deep purple during the electrochemical reduction. Absence of a clear isosbestic point during the electrochemical reduction suggests some decomposition of the reduced species of **5a**⁺ under the spectrophotometric measurements. Indeed, reverse oxidation of the purple-colored solution did not regenerate the spectrum of **5a**⁺ (regeneration 33%), completely, although the CV analysis revealed good reversibility in the two step reduction. Instability of the reduced species might be attributable to the reduction of nitro function under the electrochemical conditions.

Similar two-step spectral changes were observed during the electrochemical reduction of **5b**⁺ (see the Supporting Information). The dark green color of the solution of **5b**⁺ also changed to purple one during the electrochemical reduction. Reverse oxidation of the purple-colored solution regenerated the UV–vis spectra of the deep colored **5b**⁺ in 30%. Therefore, the two step color changes of **5a**⁺ and **5b**⁺ should be concluded to the formation of presumed radical and anionic species with some instability under the electrochemical conditions. We also tried electrochemical reduction of **8a**⁺ and **8b**⁺ under visible spectral monitoring (Figure 5 for **8a**⁺ and see the Supporting Information for **8b**⁺). We anticipated that the formation of the thienoquinoid forms during the redox reaction might improve their reversibility. In these cases, however, we also observed reversibility for the reduction of **8a**⁺ (regeneration 24%) and **8b**⁺ (regeneration 51%), insufficiently.




Figure 5. Continuous change in the visible spectrum of (a) cation **5a**⁺: constant-current electrochemical reduction (70 μ A) in benzonitrile (2.5×10^{-4} M, 2 mL) and (b) cation **8a**⁺: constant-current electrochemical reduction (70 μ A) in benzonitrile (2.6×10^{-4} M, 2 mL) containing Et₄NClO₄ (0.1 M) at 1 min intervals.




Figure 6. Continuous change in the visible spectrum of (a) cation **11a**⁺: constant-current electrochemical reduction (70 μ A) in benzonitrile (2.9×10^{-4} M, 2 mL) and (b) cation **11d**⁺: constant-current electrochemical reduction (30 μ A) in benzonitrile (4.1×10^{-4} M, 2 mL) containing Et₄NClO₄ (0.1 M) at 1 min intervals.

The strong absorption band of **11a**⁺ at 732 nm in the visible region was gradually decreased during the electrochemical reduction (Figure 6). Absence of a clear isosbestic point suggests some instability of the reduced species of **11a**⁺ under the spectrophotometric measurements. The electrochemical reduction of **11b**⁺ also caused gradual color change of the solution from purple to pink (see the Supporting Information). Reverse oxidation of the reduced solution did not regenerate the spectrum of **11a**⁺ (regeneration 57%) and **11b**⁺ (regeneration 45%), completely.

We also tried electrochemical reduction of **11c**⁺, **11d**⁺, and **14**⁺ under visible spectral monitoring (Figure 6 for **11d**⁺ and see the Supporting Information for **11c**⁺ and **14**⁺). We anticipated that the electron-withdrawing nature of the end groups should improve the two-step color changes. The purple colored solution of **11c**⁺ exhibits color change to pink one under the reduction conditions. The spectral features of **11c**⁺ under the electrochemical reduction resembled to those of **11a**⁺ and the reversibility was still low in this case (regeneration 39%). In the case of **11d**⁺, the two-strong absorption bands at 509 nm and 741 nm in the visible

region was gradually decreased along with the development of new absorption band at around 580 nm under the electrochemical reduction conditions (Figure 6). Further reduction developed a broad absorption band spread into the wide range of visible region. As expected, the spectral changes should correspond to a two-step color changes arising from the formation of a radical and an anionic species. Absence of a clear isosbestic point during the electrochemical reduction suggests some decomposition during the two-step reduction of **11d**⁺ under the spectrophotometric measurements. Reverse oxidation of the green-colored solution did not regenerate the spectrum of **11d**⁺ (regeneration 48%), completely, although the CV analysis revealed good reversibility in the two step reduction. Similarly, color change was observed during the electrochemical reduction of **14**⁺ (see the Supporting Information), but still low reversibility (regeneration 61%).

Conclusions

The scope of the creation of stabilized polyelectrochromic materials containing azulene skeletons has been demonstrated by the two-types of the C–C hybrid structure **5a**⁺, **5b**⁺, **8a**⁺, **8b**⁺, **11a–d**⁺, and **14**⁺ constructed by the cyanine units with push and pull electronic nature illustrated by the general structures in Schemes 3 and 5 with oxidation levels varying from 1+ to 1–. The system is consisted with the polymethine dye containing the moieties of either one or two polymethine dyes as end groups.^{8–9} The two-types of C–C hybrids exhibit the presumed reversible two-step, two-electron redox properties on CV in the cases of the hybrids with nitro function. Two-step color changes presumed by their C–C hybrid structure were revealed by their electrochemical reduction, although the reversibility of the redox reaction on electrochromic measurements was still low for all cases. Thus, the electrochemical behavior was not ideal for the presumed C–C hybrid, since the stabilization in the anionic state is insufficient by the introduction of nitro substituents, but the use of di(1-azulenyl)methylum units as a stabilized redox active polymethine units would be highly effective in the point of their high stability and their strong absorption in visible region with their redox activity. Preparations of the other-types of hybrid structures by using the di(1-azulenyl)methylum units as cyanine end groups are now in progress in our groups.

Experimental Section

General. Melting points were determined on a Stuart Scientific melting point apparatus SMP3 or a Yanagimoto micro melting point apparatus MP-S3 and are uncorrected. Mass spectra were obtained with a Bruker APEX II instrument under ESI conditions or a Bruker Daltonics autoflex III TOF/TOF instrument under MALDI conditions. IR and UV/Vis spectra were measured on a BIO-RAD FTS-30 or a JUSCO FT/IR-6100 infrared spectrophotometer, and a JASCO V-670 spectrophotometer, respectively. ¹H NMR spectra (¹³C NMR spectra) were recorded on a JEOL ECA500 or a JEOL ECZR500 spectrometer at 500 MHz (125 MHz). The peak assignment of ¹H and ¹³C NMR spectra reported was accomplished by HH COSY, NOE, DEPT, HMQC, and/or HMBC experiments. ¹H chemical shifts in CDCl₃ are reported in parts per million (ppm) downfield from internal tetramethylsilane. ¹³C chemical shifts in CDCl₃ are referred by the solvent signals as 77.0 ppm. Chemical shifts in (CDCl₂)₂ are referred by the solvent signals as 5.90 ppm in ¹H and 74.2 ppm in ¹³C. Gel permeation chromatography (GPC) was performed on a JAI LC-9110 NEXT with JAIGEL-1H–3H using CHCl₃ as an eluent. Column chromatography on silica gel and Al₂O₃ was performed using MERCK Silica gel 60 Art. 7734 or Cica

Silica gel 60 Art. 37564 and MERCK Aluminium oxide 90 Art. 1097, respectively. Elemental analyses were performed at the Instrumental Analysis Center of Hirosaki University.

Bis(3,6-di-*tert*-butyl-1-azulenyl)(4-nitrophenyl)methane (4a). A mixture of 4-nitrobenzaldehyde (**2**) (151 mg, 1.00 mmol) and 1,6-di-*tert*-butylazulene (**3a**) (481 mg, 2.00 mmol) in acetic acid (12 mL) was stirred at room temperature for 24 h under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH₂Cl₂. The organic layer was washed with 5% NaHCO₃ and water, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH₂Cl₂ and GPC with CHCl₃ to afford bis(3,6-di-*tert*-butyl-1-azulenyl)(4-nitrophenyl)methane (**4a**) (464 mg, 76%). green crystals. mp 201.0–203.2 °C (MeOH). IR (KBr disk, ν_{\max} , cm⁻¹): 2964 (s), 2905 (m), 2869 (m), 1603 (w), 1576 (s), 1519 (s), 1489 (w), 1479 (w), 1458 (w), 1422 (w), 1390 (w), 1363 (m), 1344 (s), 1245 (m), 1228 (m), 1197 (w), 1105 (w), 1070 (w), 1012 (w), 987 (w), 852 (w), 835 (m), 726 (w), 672 (w), 596 (w), 535 (w). UV–vis (CH₂Cl₂, λ_{\max} , nm): 243 (log ϵ 4.45), 289 sh (4.80), 300 (4.80), 343 sh (3.97), 355 (4.02), 373 (3.94), 560 sh (2.68), 607 (2.82), 669 sh (2.70), 744 sh (2.19). ¹H NMR (500 MHz, CDCl₃): δ_{H} 8.59 (d, 2H, *J* 10.7 Hz, 4-H), 8.12 (d, 2H, *J* 10.6 Hz, 8-H), 8.08 (d, 2H, *J* 8.7 Hz, 3',5'-H), 7.32 (s, 2H, 2-H), 7.26 (d, 2H, *J* 8.7 Hz, 2',6'-H), 7.21 (dd, 2H, *J* 10.7, 1.8 Hz, 5-H), 7.10 (dd, 2H, *J* 10.7, 1.8 Hz, 7-H), 6.71 (s, 1H, CH), 1.50 (s, 18H, 3-*t*-Bu), 1.41 (s, 18H, 6-*t*-Bu). ¹³C NMR (125 MHz, CDCl₃): δ_{C} 160.9 (C-6), 154.2 (C-1'), 146.1 (C-4'), 138.0 (C-3), 135.8 (C-2), 134.9 (C-4), 134.7 (C-8a), 134.2 (C-3a), 132.0 (C-8), 129.5 (C-2',6'), 128.5 (C-1), 123.5 (C-3',5'), 119.6 (C-7), 118.8 (C-5), 42.0 (CH), 38.3 (s, 6-*t*-Bu), 33.2 (s, 3-*t*-Bu), 32.2 (q, 3-*t*-Bu), 31.8 (q, 6-*t*-Bu). HRMS (ESI): calcd for C₄₃H₅₁NO₂ + Na⁺ 636.3812, found 636.3810. Anal. calcd for C₄₃H₅₁NO₂ (613.8705): C, 84.13; H, 8.37; N, 2.28. Found: C, 84.05; H, 8.37; N, 2.38.

Bis(3-methoxycarbonyl-1-azulenyl)(4-nitrophenyl)methane (4b). A mixture of 4-nitrobenzaldehyde (**2**) (452 mg, 2.99 mmol) and methyl azulene-1-carboxylate (**3b**) (402 mg, 2.16 mmol) in acetic acid (12 mL) was stirred at 90 °C for 2 days under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH₂Cl₂. The organic layer was washed with 5% NaHCO₃ and water, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH₂Cl₂ and recrystallization from hexane to afford bis(3-methoxycarbonyl-1-azulenyl)(4-nitrophenyl)methane (**4b**) (461 mg, 84%). purple crystals. mp 231.2–232.2 °C (hexane). IR (KBr disk, ν_{\max} , cm⁻¹): 2940 (w), 1691 (s), 1594 (w), 1574 (w), 1536 (w), 1519 (m), 1485 (w), 1457 (m), 1442 (m), 1418 (s), 1400 (w), 1367 (w), 1346 (m), 1310 (w), 1210 (s), 1150 (w), 1125 (w), 1103 (w), 1043 (m), 948 (w), 866 (w), 855 (w), 813 (w), 777 (m), 741 (w), 561 (w). UV–vis (CH₂Cl₂, λ_{\max} , nm): 238 (log ϵ 4.75), 260 (4.57), 288 sh (4.91), 294 (4.94), 306 (4.97), 370 (4.28), 383 (4.30), 516 sh (2.60), 554 (2.72), 597 sh (2.63), 659 sh (2.13). ¹H NMR (500 MHz, CDCl₃): δ_{H} 9.70 (d, 2H, *J* 9.7 Hz, 4-H), 8.31 (d, 2H, *J* 9.6 Hz, 8-H), 8.16 (d, 2H, *J* 8.6 Hz, 3',5'-H), 7.81 (dd, 2H, *J* 10.0, 9.8 Hz, 6-H), 7.78 (s, 2H, 2-H), 7.57 (dd, 2H, *J* 10.0, 9.7 Hz, 5-H), 7.35 (dd, 2H, *J* 9.8, 9.6 Hz, 7-H), 7.34 (d, 2H, *J* 8.6 Hz, 2',6'-H), 6.71 (s, 1H, CH), 3.86 (s, 6H, 3-CO₂Me). ¹³C NMR (125 MHz, CDCl₃): δ_{C} 165.5 (s, 3-CO₂Me), 151.6 (C-1'), 146.7 (C-4'), 141.7 (C-3a), 140.6 (C-2), 139.9 (C-8a), 139.6 (C-6), 138.4 (C-4), 135.0 (C-8), 129.8 (C-1), 129.6 (C-2',6'), 128.2 (C-5), 126.6 (C-7), 124.0 (C-3',5'), 115.4 (C-3), 51.1 (q, 3-CO₂Me), 42.4 (CH). HRMS (ESI): calcd for C₃₁H₂₃NO₆ + Na⁺ 528.1418, found 528.1415. Anal. calcd for C₃₁H₂₃NO₆ (505.5174): C, 73.65; H, 4.59; N, 2.77. Found: C, 73.61; H, 4.71; N, 2.88.

Bis(3,6-di-*tert*-butyl-1-azulenyl)(4-nitrophenyl)methylum hexafluorophosphate (5a⁺PF₆⁻). To a solution of bis(3,6-di-*tert*-butyl-1-azulenyl)(4-nitrophenyl)methane (**4a**) (201 mg, 0.327 mmol) in CH₂Cl₂ (20 mL) was added DDQ (87 mg, 0.38 mmol) at room temperature. After the solution was stirred at the same temperature for 30 min, 60% HPF₆ (2 mL) and water (20 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed

with water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was crystallized from hexane to give bis(3,6-di-*tert*-butyl-1-azulenyl)(4-nitrophenyl)methylum hexafluorophosphate (**5a**⁺ PF_6^-) (143 mg, 58%). deep green crystals. mp 272.8–275.1 °C (CH_3CN /hexane). IR (KBr disk, ν_{max} , cm^{-1}): 2964 (m), 2898 (w), 2872 (w), 1578 (w), 1542 (w), 1524 (m), 1477 (s), 1417 (m), 1400 (w), 1367 (m), 1345 (w), 1333 (s), 1311 (s), 1283 (w), 1244 (s), 1205 (w), 1179 (m), 1120 (m), 1083 (m), 1047 (w), 1026 (w), 952 (w), 869 (m), 841 (s), 810 (m), 739 (w), 707 (w), 688 (w), 614 (w), 558 (m), 505 (w). UV-vis (CH_3CN , λ_{max} , nm): 196 (log ϵ 4.68), 237 (4.75), 252 (4.75), 268 sh (4.69), 333 (4.66), 350 (4.53), 480 sh (3.78), 652 sh (4.48), 704 (4.79). ^1H NMR (500 MHz, CDCl_3): δ_{H} 9.04 (d, 2H, J 11.0 Hz, 4-H), 8.42 (d, 2H, J 8.3 Hz, 3',5'-H), 8.19 (dd, 2H, J 11.0, 1.8 Hz, 5-H), 7.90 (d, 2H, J 10.7 Hz, 8-H), 7.71 (dd, 2H, J 10.7, 1.8 Hz, 7-H), 7.67 (d, 2H, J 8.3 Hz, 2',6'-H), 7.42 (s, 2H, 2-H), 1.56 (s, 18H, 3-*t*-Bu), 1.45 (s, 18H, 6-*t*-Bu). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 169.6 (C-6), 156.2 (C⁺), 149.8 (C-3a), 149.7 (C-1' or C-4'), 148.5 (C-8a), 148.3 (C-3), 147.8 (C-1' or C-4'), 141.9 (C-2), 139.4 (C-4), 138.8 (C-8), 134.8 (C-2',6'), 133.2 (C-5), 133.0 (C-7), 131.8 (C-1), 123.8 (C-3',5'), 39.5 (s, 6-*t*-Bu), 33.3 (s, 3-*t*-Bu), 31.5 (q, 6-*t*-Bu), 31.0 (q, 3-*t*-Bu). ^1H NMR (500 MHz, $(\text{CDCl}_2)_2$, 80 °C): δ_{H} 8.97 (d, 2H, J 11.0 Hz, 4-H), 8.36 (d, 2H, J 8.6 Hz, 3',5'-H), 8.11 (dd, 2H, J 11.0, 1.7 Hz, 5-H), 7.78 (d, 2H, J 10.7 Hz, 8-H), 7.60 (dd, 2H, J 10.7, 1.7 Hz, 7-H), 7.60 (d, 2H, J 8.6 Hz, 2',6'-H), 7.42 (s, 2H, 2-H), 1.52 (s, 18H, 3-*t*-Bu), 1.41 (s, 18H, 6-*t*-Bu). ^{13}C NMR (125 MHz, $(\text{CDCl}_2)_2$, 80 °C): δ_{C} 170.3 (C-6), 156.7 (C⁺), 150.3 (C-1' or C-4'), 150.2 (C-3a), 149.1 (C-3), 148.8 (C-8a), 147.7 (C-1' or C-4'), 142.5 (C-2), 139.7 (C-4), 138.7 (C-8), 134.9 (C-2',6'), 133.4 (C-5), 132.9 (C-7), 132.1 (C-1), 124.1 (C-3',5'), 39.7 (s, 6-*t*-Bu), 33.6 (s, 3-*t*-Bu), 31.7 (q, 6-*t*-Bu), 31.3 (q, 3-*t*-Bu). HRMS (ESI): calcd for $\text{C}_{43}\text{H}_{50}\text{NO}_2^+$ 612.3836, found 612.3834. Anal. calcd for $\text{C}_{43}\text{H}_{50}\text{F}_6\text{NO}_2\text{P}$ (757.8268): C, 68.15; H, 6.65; N, 1.85. Found: C, 67.73; H, 6.44; N, 2.07.

Bis(3-methoxycarbonyl-1-azulenyl)(4-nitrophenyl)methylum hexafluorophosphate (5b⁺ PF_6^-). To a solution of bis(3-methoxycarbonyl-1-azulenyl)(4-nitrophenyl)methane (**4b**) (200 mg, 0.396 mmol) in CH_2Cl_2 (40 mL) was added DDQ (109 mg, 0.480 mmol) at room temperature. After the solution was stirred at the same temperature for 30 min, 60% HPF_6 (4 mL) and water (40 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was dissolved in a small amount of CH_2Cl_2 and then poured into Et_2O . The precipitated crystals were collected by filtration, washed with Et_2O , and dried in vacuo to give bis(3-methoxycarbonyl-1-azulenyl)(4-nitrophenyl)methylum hexafluorophosphate (**5b**⁺ PF_6^-) (190 mg, 74%). deep red crystals. mp 173.2–173.6 °C ($\text{CH}_2\text{Cl}_2/\text{Et}_2\text{O}$). IR (KBr disk, ν_{max} , cm^{-1}): 2947 (w), 1705 (s), 1590 (w), 1524 (m), 1474 (s), 1452 (s), 1417 (s), 1349 (s), 1321 (m), 1301 (w), 1260 (w), 1240 (s), 1228 (s), 1196 (m), 1162 (s), 1089 (s), 1064 (m), 1011 (w), 913 (w), 877 (m), 841 (s), 782 (w), 757 (w), 738 (w), 707 (w), 587 (w), 557 (w), 460 (w). UV-vis (CH_2Cl_2 , λ_{max} , nm): 201 (log ϵ 4.53), 230 (4.64), 262 (4.61), 293 (4.55), 303 (4.55), 348 (4.26), 367 sh (4.23), 438 (3.67), 591 sh (4.18), 637 (4.63). ^1H NMR (500 MHz, $(\text{CDCl}_2)_2$, 80 °C): δ_{H} 10.01 (d, 2H, J 11.1 Hz, 4-H), 8.37 (d, 2H, J 8.5 Hz, 3',5'-H), 8.30–8.23 (m, 4H, 5,6-H), 8.20 (s, 2H, 2-H), 8.17 (d, 2H, J 10.0 Hz, 8-H), 7.80 (dd, 2H, J 10.0, 9.1 Hz, 7-H), 7.62 (d, 2H, J 8.5 Hz, 2',6'-H), 3.92 (s, 6H, 3- CO_2Me). ^{13}C NMR (125 MHz, $(\text{CDCl}_2)_2$, 80 °C): δ_{C} 165.4 (C⁺), 163.8 (s, 3- CO_2Me), 152.5 (C-3a), 151.2 (C-1' or C-4'), 149.9 (C-8a), 149.6 (C-2), 146.6 (C-1' or C-4'), 146.5 (C-6), 143.2 (C-4), 141.5 (C-8), 139.5 (C-5), 138.4 (C-7), 135.7 (C-2',6'), 132.2 (C-1), 126.4 (C-3), 124.5 (C-3',5'), 52.6 (q, 3- CO_2Me). HRMS (ESI): calcd for $\text{C}_{31}\text{H}_{22}\text{NO}_6^+$ 504.1442, found 504.1441. Anal. calcd for $\text{C}_{31}\text{H}_{22}\text{F}_6\text{NO}_6\text{P}$ (649.4737): C, 57.33; H, 3.41; N, 2.16. Found: C, 57.20; H, 3.49; N, 2.38.

Bis(3,6-di-*tert*-butyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (7a). A mixture of 5-(4-nitrophenyl)thiophene-2-carbaldehyde (**6**) (76 mg, 0.339 mmol) and 1,6-di-*tert*-butylazulene (**3a**) (157 mg, 0.653 mmol) in acetic acid (5 mL) was stirred at room temperature for 24 h under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH_2Cl_2 . The organic layer was washed with 5% NaHCO_3 and water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was purified by

column chromatography on silica gel with CH_2Cl_2 and GPC with CHCl_3 to afford bis(3,6-di-*tert*-butyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (**7a**) (140 mg, 62%). green crystals. mp 248.5–251.1 °C (MeOH). IR (KBr disk, ν_{max} , cm^{-1}): 2963 (s), 2895 (m), 2868 (m), 1592 (s), 1577 (s), 1514 (s), 1488 (w), 1458 (w), 1424 (w), 1384 (w), 1363 (m), 1342 (s), 1246 (w), 1226 (w), 1111 (w), 1060 (w), 1051 (w), 984 (w), 949 (w), 879 (w), 851 (m), 831 (m), 820 (w), 795 (w), 751 (m), 681 (w), 669 (w), 451 (w). UV–vis (CH_2Cl_2 , λ_{max} , nm): 244 (log ϵ 4.53), 284 (4.83), 302 (4.80), 343 sh (4.36), 358 (4.46), 373 (4.45), 562 sh (2.74), 608 (2.85), 661 sh (2.76), 741 sh (2.23). ^1H NMR (500 MHz, CDCl_3): δ_{H} 8.59 (d, 2H, J 10.6 Hz, 4-H), 8.27 (d, 2H, J 10.6 Hz, 8-H), 8.14 (d, 2H, J 9.0 Hz, 3'',5''-H), 7.60 (s, 2H, 2-H), 7.59 (d, 2H, J 9.0 Hz, 2'',6''-H), 7.26 (d, 1H, J 3.8 Hz, 4'-H), 7.21 (dd, 2H, J 10.6, 1.8 Hz, 5-H), 7.15 (dd, 2H, J 10.6, 1.8 Hz, 7-H), 6.84 (s, 1H, CH), 6.65 (dd, 1H, J 3.8, 1.0 Hz, 3'-H), 1.53 (s, 18H, 3-*t*-Bu), 1.42 (s, 18H, 6-*t*-Bu). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 160.7 (C-6), 154.9 (C-2'), 146.0 (C-4''), 141.1 (C-1''), 139.4 (C-5'), 137.9 (C-3), 135.4 (C-2), 134.8 (C-4), 134.45 (C-8a), 134.38 (C-3a), 131.8 (C-8), 129.1 (C-1), 126.9 (C-3'), 125.4 (C-4'), 125.3 (C-2'',6''), 124.3 (C-3'',5''), 119.7 (C-7), 118.7 (C-5), 38.2 (s, 6-*t*-Bu), 37.5 (d, CH), 33.3 (s, 3-*t*-Bu), 32.2 (q, 3-*t*-Bu), 31.8 (q, 6-*t*-Bu). HRMS (ESI): calcd for $\text{C}_{47}\text{H}_{53}\text{NO}_2\text{S} + \text{Na}^+$ 718.3689, found 718.3687. Anal. calcd for $\text{C}_{47}\text{H}_{53}\text{NO}_2\text{S}$ (695.9942): C, 81.11; H, 7.68; N, 2.01; S, 4.61. Found: C, 80.81; H, 7.82; N, 2.04; S, 4.46.

Bis(3-methoxycarbonyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (7b). A mixture of 5-(4-nitrophenyl)thiophene-2-carbaldehyde (**6**) (644 g, 2.76 mmol) and methyl azulene-1-carboxylate (**3b**) (1.03 g, 5.54 mmol) in acetic acid (50 mL) and toluene (20 mL) was refluxed for 8 days under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH_2Cl_2 . The organic layer was washed with 5% NaHCO_3 and water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH_2Cl_2 to afford bis(3-methoxycarbonyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (**7b**) (572 mg, 35%). reddish purple crystals. mp 230.0–232.1 °C (hexane). IR (KBr disk, ν_{max} , cm^{-1}): 2926 (w), 2837 (w), 1692 (s), 1593 (m), 1580 (w), 1536 (w), 1514 (m), 1486 (w), 1457 (m), 1441 (m), 1418 (s), 1363 (w), 1344 (m), 1337 (s), 1310 (w), 1251 (w), 1210 (s), 1155 (w), 1110 (w), 1043 (m), 951 (w), 852 (m), 816 (w), 776 (m), 745 (w), 740 (w), 680 (w), 566 (w), 455 (w). UV–vis (CH_2Cl_2 , λ_{max} , nm): 238 (log ϵ 4.63), 259 (4.44), 288 sh (4.71), 294 (4.73), 305 (4.76), 371 (4.46), 381 (4.46), 514 sh (2.78), 553 (2.89), 593 sh (2.80), 657 sh (2.30). ^1H NMR (500 MHz, CDCl_3): δ_{H} 9.69 (d, 2H, J 9.5 Hz, 4-H), 8.44 (d, 2H, J 9.6 Hz, 8-H), 8.15 (d, 2H, J 9.0 Hz, 3'',5''-H), 8.05 (s, 2H, 2-H), 7.80 (dd, 2H, J 9.9, 9.8 Hz, 6-H), 7.61 (d, 2H, J 9.0 Hz, 2'',6''-H), 7.56 (dd, 2H, J 9.9, 9.5 Hz, 5-H), 7.38 (dd, 2H, J 9.8, 9.6 Hz, 7-H), 7.32 (d, 1H, J 3.8 Hz, 4'-H), 6.85 (s, 1H, CH), 6.77 (d, 1H, J 3.8 Hz, 3'-H), 3.87 (s, 6H, 3-CO₂Me). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 165.6 (s, 3-CO₂Me), 151.4 (C-2'), 146.3 (C-4''), 141.7 (C-3a), 140.6 (C-1''), 140.3 (C-2 and C-5'), 139.7 (C-8a), 139.5 (C-6), 138.3 (C-4), 134.9 (C-8), 130.6 (C-1), 128.1 (C-5), 127.5 (C-3'), 126.6 (C-7), 125.6 (C-4'), 125.5 (C-2'',6''), 124.3 (C-3'',5''), 115.3 (C-3), 51.1 (q, 3-CO₂Me), 37.8 (CH). HRMS (ESI): calcd for $\text{C}_{35}\text{H}_{25}\text{NO}_6\text{S} + \text{Na}^+$ 610.1295, found 610.1292. Anal. calcd for $\text{C}_{35}\text{H}_{25}\text{NO}_6\text{S}$ (587.6411): C, 71.54; H, 4.29; N, 2.38. Found: C, 71.25; H, 4.44; N, 2.14.

Bis(3,6-di-*tert*-butyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methylum hexafluorophosphate (8a⁺PF₆⁻). To a solution of bis(3,6-di-*tert*-butyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (**7a**) (100 mg, 0.144 mmol) in CH_2Cl_2 (15 mL) was added DDQ (34 mg, 0.15 mmol) at room temperature. After the solution was stirred at the same temperature for 30 min, 60% HPF₆ (2 mL) and water (20 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was dissolved in a small amount of CH_2Cl_2 and then poured into Et_2O . The precipitated crystals were collected by filtration, washed with Et_2O , and dried in vacuo to give bis(3,6-di-*tert*-butyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methylum hexafluorophosphate (**8a⁺PF₆⁻**) (93 mg, 77%). deep purple crystals. mp 208.1–211.2 °C ($\text{CH}_3\text{CN}/\text{Et}_2\text{O}$). IR (KBr disk, ν_{max} , cm^{-1}): 2963 (m), 2916 (w), 2872 (w), 1594 (w), 1578 (w), 1543 (w), 1523 (m),

1472 (s), 1422 (s), 1397 (w), 1376 (m), 1360 (m), 1333 (s), 1312 (s), 1263 (w), 1245 (m), 1207 (m), 1189 (w), 1177 (w), 1113 (m), 1079 (w), 1022 (w), 950 (w), 868 (m), 840 (s), 807 (m), 743 (w), 705 (w), 558 (m), 521 (w). UV-vis (CH₃CN, λ_{max} , nm): 199 sh (log ϵ 4.75), 241 (4.72), 270 sh (4.64), 306 (4.60), 337 (4.50), 381 sh (4.33), 409 (4.46), 429 sh (4.40), 540 (4.55), 650 sh (4.40), 702 (4.69). ¹H NMR (500 MHz, (CDCl₂)₂, 100 °C): δ_{H} 8.97 (d, 2H, *J* 11.0 Hz, 4-H), 8.25 (d, 2H, *J* 8.9 Hz, 3'',5''-H), 8.05 (dd, 2H, *J* 11.0, 1.9 Hz, 5-H), 7.88 (d, 2H, *J* 10.8 Hz, 8-H), 7.84 (d, 2H, *J* 8.9 Hz, 2'',6''-H), 7.79 (d, 1H, *J* 4.0 Hz, 4'-H), 7.75 (s, 2H, 2-H), 7.60 (dd, 2H, *J* 10.8, 1.9 Hz, 7-H), 7.53 (d, 1H, *J* 4.0 Hz, 3'-H), 1.58 (s, 18H, 3-*t*-Bu), 1.42 (s, 18H, 6-*t*-Bu). ¹³C NMR (125 MHz, (CDCl₂)₂, 100 °C): δ_{C} 169.7 (C-6), 154.4 (C-5'), 150.6 (C⁺), 149.0 (C-3a), 148.6 (C-4''), 148.4 (C-8a), 148.1 (C-3), 146.0 (C-2'), 142.3 (C-2), 140.2 (C-3'), 139.4 (C-4), 139.0 (C-1''), 138.3 (C-8), 132.0 (C-5), 131.6 (C-7), 131.2 (C-1), 128.6 (C-4'), 127.2 (C-2'',6''), 124.9 (C-3'',5''), 39.6 (s, 6-*t*-Bu), 33.6 (s, 3-*t*-Bu), 31.7 (q, 6-*t*-Bu), 31.5 (q, 3-*t*-Bu). HRMS (ESI): calcd for C₄₇H₅₂NO₂S⁺ 694.3713, found 694.3714. Anal. calcd for C₄₇H₅₂F₆NO₂PS (839.9505): C, 67.21; H, 6.24; N, 1.67; S, 3.82. Found: C, 67.16; H, 6.36; N, 1.61; S, 3.76.

Bis(3-methoxycarbonyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methylum hexafluorophosphate (8b⁺PF₆⁻).

To a solution of bis(3-methoxycarbonyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methane (**7b**) (101 mg, 0.172 mmol) in CH₂Cl₂ (20 mL) was added DDQ (47 mg, 0.21 mmol) at room temperature. After the solution was stirred at the same temperature for 30 min, 60% HPF₆ (2 mL) and water (20 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried over MgSO₄, and concentrated under reduced pressure. The residue was dissolved in a small amount of CHCl₃ and then poured into hexane. The precipitated crystals were collected by filtration, washed with hexane, and dried in vacuo to give bis(3-methoxycarbonyl-1-azulenyl)[5-(4-nitrophenyl)-2-thienyl]methylum hexafluorophosphate (**8b⁺PF₆⁻**) (42 mg, 33%). deep purple crystals. mp 208.1–209.9 °C (CH₂Cl₂/Et₂O). IR (KBr disk, ν_{max} , cm⁻¹): 2947 (w), 1703 (s), 1593 (w), 1580 (w), 1534 (w), 1518 (m), 1459 (s), 1417 (s), 1350 (s), 1340 (s), 1274 (w), 1241 (s), 1224 (s), 1163 (s), 1101 (w), 1085 (m), 1068 (m), 1064 (w), 1037 (w), 1012 (w), 914 (w), 877 (w), 842 (s), 783 (w), 752 (w), 685 (w), 600 (w), 558 (m). UV-vis (CH₃CN, λ_{max} , nm): 197 (log ϵ 4.72), 229 (4.73), 252 (4.64), 272 (4.62), 300 (4.64), 331 sh (4.42), 365 sh (4.30), 385 sh (4.25), 416 sh (4.12), 536 (4.57), 630 (4.50). ¹H NMR (500 MHz, (CDCl₂)₂, 80 °C): δ_{H} 10.01 (d, 2H, *J* 9.9 Hz, 4-H), 8.57 (s, 2H, 2-H), 8.25 (d, 2H, *J* 8.7 Hz, 3'',5''-H), 8.21 (dd, 2H, *J* 9.7, 9.3 Hz, 6-H), 8.16 (dd, 2H, *J* 9.9, 9.7 Hz, 5-H), 8.14 (d, 2H, *J* 9.8 Hz, 8-H), 7.90 (d, 1H, *J* 4.2 Hz, Th), 7.86 (d, 2H, *J* 8.7 Hz, 2'',6''-H), 7.73 (d, 1H, *J* 4.2 Hz, Th), 7.71 (dd, 2H, *J* 9.8, 9.3 Hz, 7-H), 3.97 (s, 6H, 3-CO₂Me). ¹³C NMR (125 MHz, (CDCl₂)₂, 80 °C): δ_{C} 164.2 (s, 3-CO₂Me), 160.2 (C-5'), 158.3 (C⁺), 151.1 (C-3a), 149.2 (C-8a or C-4''), 149.1 (C-8a or C-4''), 148.9 (C-2), 146.0 (C-2'), 145.8 (C-6), 143.8 (d, Th), 143.0 (C-4), 140.5 (C-8), 138.1 (C-1''), 137.9 (C-5), 136.6 (C-7), 130.5 (C-1), 130.2 (d, Th), 127.8 (C-2'',6''), 125.0 (C-3'',5''), 124.8 (C-3), 52.5 (q, 3-CO₂Me). HRMS (ESI): calcd for C₃₅H₂₄NO₆S⁺ 586.1319, found 586.1318.

3,3-Bis(4-nitrophenyl)-2-propenal (9c). To a solution of 4,4'-dinitrobenzophenone (709 mg, 2.60 mmol) in CH₂Cl₂ (25 mL) was added a solution of TiCl₄ (1.1 mL, 10.0 mmol) in CH₂Cl₂ (1.1 mL) and Et₃N (1.4 mL, 10.1 mmol) at 0 °C under an N₂ atmosphere. The resulting mixture was stirred at 30 °C for 24 h. The reaction mixture was poured into saturated NH₄Cl (50 mL) and stirred at the same temperature for 3 h. The mixture was extracted with CH₂Cl₂. The organic layer was washed with brine, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH₂Cl₂ to afford 3,3-bis(4-nitrophenyl)-2-propenal (**9c**) (208 mg, 27%). yellow crystals. mp 173.2–174.0 °C (MeOH). IR (KBr disk, ν_{max} , cm⁻¹): 3109 (w), 3077 (w), 2926 (w), 2849 (w), 1670 (s), 1599 (w), 1518 (s), 1430 (w), 1348 (s), 1300 (w), 1253 (w), 1117 (w), 1124 (m), 1168 (w), 1013 (w), 921 (w), 860 (m), 846 (m), 759 (w), 746 (w), 706 (w). UV-vis (CH₂Cl₂, λ_{max} , nm): 299 (log ϵ 4.65). ¹H NMR (500 MHz, CDCl₃): δ_{H} 9.57 (d, 1H, *J* 7.7 Hz, 1-H), 8.39 (d, 2H, *J* 8.8 Hz, 3',5'-H or 3'',5''-H), 8.27 (d, 2H, *J* 9.1 Hz, 3',5'-H or 3'',5''-H), 7.53 (d, 2H, *J* 8.8 Hz, 2',6'-H or 2'',6''-

H), 7.50 (d, 2H, J 9.1 Hz, 2',6'-H or 2'',6''-H), 6.74 (d, 1H, J 7.7 Hz, 2-H). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 191.2 (C-1), 156.2 (C-3), 149.9 (C-4' or C-4''), 148.6 (C-4' or C-4''), 144.4 (C-1' or C-1''), 141.8 (C-1' or C-1''), 131.4 (C-2',6' or C-2'',6''), 130.3 (C-2), 129.2 (C-2',6' or C-2'',6''), 124.1 (C-3',5' or C-3'',5''), 124.0 (C-3',5' or C-3'',5''). HRMS (MALDI): calcd for $\text{C}_{15}\text{H}_{10}\text{N}_2\text{O}_5 - \text{H}^- [\text{M} - \text{H}]^+$ 297.0506, found 297.0468. Anal. Calcd for $\text{C}_{15}\text{H}_{10}\text{N}_2\text{O}_5$ (298.2503): C, 60.41; H, 3.38; N, 9.39. Found: C, 60.30; H, 3.33; N, 9.43.

3,3-Bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-diphenylpropene (10a). A mixture of β -phenylcinnamaldehyde (**9a**) (208 mg, 1.00 mmol) and 1,6-di-*tert*-butylazulene (**3a**) (482 mg, 2.01 mmol) in acetic acid (12 mL) was stirred at room temperature for 24 h under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH_2Cl_2 . The organic layer was washed with 5% NaHCO_3 and water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH_2Cl_2 and GPC with CHCl_3 to afford 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-diphenylpropene (**10a**) (507 mg, 76%). blue crystals. mp 210.1–211.1 °C (MeOH). IR (KBr disk, ν_{max} , cm^{-1}): 3079 (w), 3050 (w), 3018 (w), 2963 (s), 2904 (m), 2868 (m), 1576 (s), 1507 (w), 1492 (w), 1455 (m), 1441 (w), 1424 (m), 1390 (m), 1363 (s), 1245 (m), 1228 (m), 1198 (w), 1072 (w), 874 (w), 833 (m), 766 (w), 702 (m), 673 (w), 580 (w), 531 (w), 458 (w), 442 (w). UV-vis (CH_2Cl_2 , λ_{max} , nm): 246 (log ϵ 4.57), 288 (4.83), 302 sh (4.80), 345 sh (3.92), 357 (4.00), 375 (3.94), 560 sh (2.57), 604 (2.74), 674 sh (2.63), 748 sh (2.04). ^1H NMR (500 MHz, CDCl_3): δ_{H} 8.52 (d, 2H, J 10.6 Hz, 4'-H), 7.81 (d, 2H, J 10.6 Hz, 8'-H), 7.66 (s, 2H, 2'-H), 7.40–7.39 (m, 3H, Ph), 7.22–7.18 (m, 7H, Ph), 7.12 (dd, 2H, J 10.6, 1.9 Hz, 5'-H), 6.92 (dd, 2H, J 10.6, 1.9 Hz, 7'-H), 6.81 (d, 1H, J 10.5 Hz, 2-H), 5.95 (d, 1H, J 10.5 Hz, 3-H), 1.54 (s, 18H, 3'-*t*-Bu), 1.40 (s, 18H, 6'-*t*-Bu). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 160.2 (C-6'), 142.8 (C-1'' or C-1'''), 140.2 (C-1'' or C-1'''), 138.3 (C-1), 137.9 (C-3'), 134.7 (C-2'), 134.3 (C-4'), 134.14 (C-3'a), 134.08 (C-8'a), 132.9 (C-2), 132.3 (C-8'), 130.7 (C-1'), 130.6 (d, Ph), 128.2 (d, Ph), 128.0 (d, Ph), 127.7 (d, Ph), 127.2 (d, Ph), 126.8 (d, Ph), 118.9 (C-7'), 118.0 (C-5'), 38.1 (s, 6'-*t*-Bu), 36.8 (C-3), 33.3 (s, 3'-*t*-Bu), 32.2 (q, 3'-*t*-Bu), 31.8 (q, 6'-*t*-Bu). HRMS (ESI): calcd for $\text{C}_{51}\text{H}_{58} + \text{Na}^+$ 693.4431, found 693.4427. Anal. calcd for $\text{C}_{51}\text{H}_{58}$ (671.0062): C, 91.29; H, 8.71. Found: C, 91.04; H, 8.56.

3,3-Bis(3-methoxycarbonyl-1-azulenyl)-1,1-diphenylpropene (10b). A mixture of β -phenylcinnamaldehyde (**9a**) (514 mg, 2.47 mmol) and methyl azulene-1-carboxylate (**3b**) (300 mg, 1.61 mmol) in acetic acid (20 mL) was refluxed for 2 days under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH_2Cl_2 . The organic layer was washed with 5% NaHCO_3 and water, dried over MgSO_4 , and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH_2Cl_2 and ethyl acetate to afford 3,3-bis(3-methoxycarbonyl-1-azulenyl)-1,1-diphenylpropene (**10b**) (244 mg, 27%). purple crystals. mp 201.1–203.3 °C (Et_2O /hexane). IR (KBr disk, ν_{max} , cm^{-1}): 3016 (w), 2946 (w), 1692 (s), 1587 (w), 1577 (w), 1536 (w), 1513 (w), 1489 (w), 1457 (s), 1441 (s), 1418 (s), 1371 (w), 1309 (w), 1207 (s), 1154 (w), 1136 (w), 1042 (w), 944 (w), 886 (w), 871 (w), 863 (w), 777 (m), 749 (m), 703 (m), 621 (w), 561 (w), 538 (w). UV-vis (CH_2Cl_2 , λ_{max} , nm): 239 (log ϵ 4.72), 261 sh (4.54), 289 sh (4.80), 294 (4.81), 306 (4.82), 372 (4.12), 385 (4.16), 519 sh (2.83), 559 (2.94), 596 sh (2.86), 664 (2.34). ^1H NMR (500 MHz, CDCl_3): δ_{H} 9.65 (d, 2H, J 9.5 Hz, 4'-H), 8.19 (s, 2H, 2'-H), 8.01 (s, 2H, J 9.5 Hz, 8'-H), 7.73 (dd, 2H, J 9.8, 9.8 Hz, 6'-H), 7.51 (dd, 2H, J 9.8, 9.5 Hz, 5'-H), 7.40–7.34 (m, 3H, 4''-H or 4'''-H and 3'',5''-H or 3''',5'''-H), 7.25–7.16 (m, 9H, 7'-H, 2'',6''-H, 2''',6'''-H, and 4''-H or 4'''-H and 3'',5''-H or 3''',5'''-H), 6.73 (d, 1H, J 10.3 Hz, 2-H), 5.94 (d, 1H, J 10.3 Hz, 3-H), 3.90 (s, 6H, 3'- CO_2Me). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 165.7 (s, 3'- CO_2Me), 142.1 (C-1'' or C-1'''), 141.6 (C-3'a), 140.4 (C-1), 139.8 (C-8'a), 139.7 (C-1'' or C-1'''), 139.4 (C-2'), 139.1 (C-6'), 137.8 (C-4'), 135.2 (C-8'), 132.0 (C-1'), 130.9 (C-2), 130.1 (d, Ph), 128.4 (d, Ph), 128.1 (d, Ph), 127.8 (d, Ph), 127.64 (d, Ph), 127.62 (C-5'), 127.3 (d, Ph), 126.1 (C-7'), 115.4 (C-3'), 51.0 (q, 3'- CO_2Me), 37.1 (C-3). HRMS (ESI): calcd for $\text{C}_{39}\text{H}_{30}\text{O}_4 + \text{Na}^+$ 585.2036, found 585.2034. Anal. calcd for $\text{C}_{39}\text{H}_{30}\text{O}_4$ (562.6531): C, 83.25; H, 5.37. Found: C, 83.03; H, 5.58.

3,3-Bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-fluorophenyl)propene (10c). A mixture of 3,3-bis(4-fluorophenyl)-2-propenal (**9c**) (218 mg, 0.893 mmol) and 1,6-di-*tert*-butylazulene (**3a**) (421 mg, 1.75 mmol) in acetic acid (10 mL) was stirred at room temperature for 2 days under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH₂Cl₂. The organic layer was washed with 5% NaHCO₃ and water, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH₂Cl₂/hexane (1:5) and GPC with CHCl₃ to afford 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-fluorophenyl)propene (**10c**) (264 mg, 43%). blue crystals. mp 143.0–144.2 °C (MeOH). IR (KBr disk, ν_{\max} , cm⁻¹): 2964 (s), 2895 (m), 2869 (m), 1602 (w), 1576 (s), 1509 (s), 1456 (m), 1424 (m), 1387 (m), 1363 (m), 1228 (s), 1200 (w), 1158 (m), 1086 (w), 1064 (w), 1010 (w), 872 (w), 834 (s), 669 (w), 570 (w). UV-vis (CH₂Cl₂, λ_{\max} , nm): 246 (log ϵ 4.42), 288 (4.70), 300 sh (4.67), 344 sh (3.80), 357 (3.88), 374 (3.81), 564 sh (2.58), 613 (2.71), 675 sh (2.60), 749 sh (2.10). ¹H NMR (500 MHz, CDCl₃): δ_{H} 8.53 (d, 2H, *J* 10.6 Hz, 4'-H), 7.79 (d, 2H, *J* 10.7 Hz, 8'-H), 7.64 (s, 2H, 2'-H), 7.15–7.12 (m, 6H, 5'-H, 2'',6''-H, and 2''',6'''-H), 7.07 (dd, 2H, *J* 8.7, 8.7 Hz, 3'',5''-H or 3''',5'''-H), 6.94 (dd, 2H, *J* 10.6, 1.9 Hz, 7'-H), 6.91 (dd, 2H, *J* 8.7, 8.7 Hz, 3'',5''-H or 3''',5'''-H), 6.74 (d, 1H, *J* 10.4 Hz, 2-H), 5.90 (d, 1H, *J* 10.4 Hz, 3-H), 1.54 (s, 18H, 3'-*t*-Bu), 1.40 (s, 18H, 6'-*t*-Bu). ¹³C NMR (125 MHz, CDCl₃): δ_{C} 162.2 (*J* 246.6 Hz, C-4'' or C-4'''), 162.1 (*J* 246.6 Hz, C-4'' or C-4'''), 160.4 (C-6'), 138.7 (*J* 3.2 Hz, C-1'' or C-1'''), 138.0 (C-3'), 136.4 (C-1), 135.9 (*J* 3.2 Hz, C-1'' or C-1'''), 134.5 (C-2'), 134.4 (C-4'), 134.2 (C-3'a), 134.0 (C-8'a), 133.2 (C-2), 132.10 (C-8'), 132.07 (*J* 7.8 Hz, C-2'',6'' or C-2''',6'''), 130.4 (C-1'), 129.2 (*J* 7.8 Hz, C-2'',6'' or C-2''',6'''), 118.9 (C-7'), 118.2 (C-5'), 115.2 (*J* 21.0 Hz, C-3'',5'' or C-3''',5'''), 114.8 (*J* 21.0 Hz, C-3'',5'' or C-3''',5'''), 38.2 (s, 6'-*t*-Bu), 36.8 (C-3), 33.3 (s, 3'-*t*-Bu), 32.2 (q, 3'-*t*-Bu), 31.8 (q, 6'-*t*-Bu). HRMS (ESI): calcd for C₅₁H₅₆F₂ + Na⁺ 729.4242, found 729.4240. Anal. calcd for C₅₁H₅₆F₂ (706.9871): C, 86.64; H, 7.98. Found: C, 86.39; H, 8.09.

3,3-Bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propene (10d). A mixture of 3,3-bis(4-nitrophenyl)-2-propenal (**9c**) (208 mg, 0.697 mmol), 1,6-di-*tert*-butylazulene (**3a**) (329 mg, 1.37 mmol) in acetic acid (12 mL) was stirred at 35 °C for 4 days under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH₂Cl₂. The organic layer was washed with 5% NaHCO₃ and water, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH₂Cl₂/hexane (1:1) and recrystallization from ethyl acetate/methanol to afford 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propene (**10d**) (157 mg, 30%). green crystals. mp 158.0–158.9 °C (MeOH). IR (KBr disk, ν_{\max} , cm⁻¹): 2962 (m), 2950 (w), 2931 (w), 1577 (m), 1520 (s), 1454 (w), 1417 (w), 1384 (w), 1360 (m), 1341 (s), 1242 (w), 1226 (w), 1109 (w), 1069 (w), 855 (w), 834 (w), 704 (w). UV-vis (CH₂Cl₂, λ_{\max} , nm): 243 (log ϵ 4.59), 284 sh (4.84), 293 (4.86), 303 (4.88), 356 sh (4.19), 373 (4.09), 558 sh (2.87), 607 sh (2.82), 669 sh (2.65), 749 sh (2.08). ¹H NMR (500 MHz, CDCl₃): δ_{H} 8.56 (d, 2H, *J* 10.7 Hz, 4'-H), 8.24 (d, 2H, *J* 8.7 Hz, 3'',5''-H or 3''',5'''-H), 8.10 (d, 2H, *J* 9.0 Hz, 3'',5''-H or 3''',5'''-H), 7.72 (d, 2H, *J* 10.6 Hz, 8'-H), 7.63 (s, 2H, 2'-H), 7.30 (d, 2H, *J* 8.7 Hz, 2'',6''-H or 2''',6'''-H), 7.29 (d, 2H, *J* 9.0 Hz, 2'',6''-H or 2''',6'''-H), 7.20 (dd, 2H, *J* 10.7, 1.8 Hz, 5'-H), 7.08 (d, 1H, *J* 10.2 Hz, 2-H), 6.97 (dd, 2H, *J* 10.6, 1.8 Hz, 7'-H), 5.87 (d, 1H, *J* 10.2 Hz, 3-H), 1.55 (s, 18H, 3'-*t*-Bu), 1.41 (s, 18H, 6'-*t*-Bu). ¹³C NMR (125 MHz, CDCl₃): δ_{C} 161.0 (C-6'), 147.7 (C-1'' or C-1'''), 147.4 (C-4'' or C-4'''), 146.8 (C-4'' or C-4'''), 145.8 (C-1'' or C-1'''), 138.5 (C-2), 138.3 (C-3'), 135.2 (C-1), 134.8 (C-4'), 134.3 (C-3'a), 134.2 (C-2'), 134.1 (C-8'a), 131.7 (C-8'), 131.2 (C-2'',6'' or C-2''',6'''), 128.7 (C-1'), 128.1 (C-2'',6'' or C-2''',6'''), 123.8 (C-3'',5'' or C-3''',5'''), 123.6 (C-3'',5'' or C-3''',5'''), 119.3 (C-7'), 118.7 (C-5'), 38.2 (s, 6'-*t*-Bu), 37.1 (C-3), 33.3 (s, 3'-*t*-Bu), 32.2 (q, 3'-*t*-Bu), 31.8 (q, 6'-*t*-Bu). HRMS (ESI): calcd for C₅₁H₅₆N₂O₄ + Na⁺ 783.4132, found 783.4130. Anal. Calcd for C₅₁H₅₆N₂O₄ (761.0013): C, 80.49; H, 7.42; N, 3.68. Found: C, 80.17; H, 7.24; N, 3.54.

3,3-Bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-diphenylpropenylium hexafluorophosphate (11a⁺PF₆⁻). To a solution of 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-diphenylpropene (**10a**) (200 mg, 0.298 mmol) in CH₂Cl₂ (30 mL) was added DDQ (81 mg, 0.36 mmol) at room temperature. After the solution was stirred at the same temperature

for 30 min, 60% HPF₆ (3 mL) and water (20 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried with MgSO₄, and concentrated under reduced pressure. The residue was crystallized from hexane to give 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-diphenylpropenylium hexafluorophosphate (**11a**⁺PF₆⁻) (179 mg, 74%). deep purple powder. mp 152.3–154.0 °C (Et₂O/hexane). IR (KBr disk, ν_{\max} , cm⁻¹): 2963 (m), 2896 (w), 2872 (w), 1577 (w), 1545 (w), 1490 (m), 1476 (s), 1440 (m), 1416 (s), 1396 (w), 1366 (m), 1335 (s), 1311 (s), 1281 (w), 1239 (m), 1232 (m), 1200 (w), 1181 (w), 1178 (w), 1101 (w), 1076 (w), 1018 (w), 984 (w), 956 (w), 874 (w), 839 (s), 807 (m), 699 (w), 557 (m). UV-vis (CH₃CN, λ_{\max} , nm): 195 (log ϵ 4.75), 241 (4.67), 309 (4.51), 323 sh (4.48), 416 (4.09), 435 sh (4.02), 534 (4.18), 722 (4.58). ¹H NMR (500 MHz, (CDCl₂)₂, 120 °C): δ_{H} 8.75 (d, 2H, *J* 10.9 Hz, 4'-H), 8.47 (d, 2H, *J* 10.7 Hz, 8'-H), 7.94 (d, 2H, *J* 10.9 Hz, 5'-H), 7.71 (s, 1H, 2-H), 7.67 (d, 2H, *J* 10.7 Hz, 7'-H), 7.62 (s, 2H, 2'-H), 7.50–7.45 (m, 5H, Ph), 6.82–6.74 (m, 5H, Ph), 1.47 (s, 18H, 3'-*t*-Bu), 1.44 (s, 18H, 6'-*t*-Bu). ¹³C NMR (125 MHz, (CDCl₂)₂, 120 °C): δ_{C} 168.9 (C-6'), 161.5 (C-1), 156.7 (C-3), 148.5 (C-3'a), 148.0 (C-8'a), 147.8 (C-3'), 142.1 (C-2'), 141.7 (C-1'' or C-1'''), 139.1 (C-1'' or C-1'''), 138.5 (C-4'), 137.8 (C-8'), 131.7 (C-1'), 131.6 (C-5'), 131.5 (C-7'), 131.1 (d, Ph), 130.4 (d, Ph), 130.2 (C-2), 129.6 (d, Ph), 129.3 (d, Ph), 128.7 (d, Ph), 127.9 (d, Ph), 39.4 (s, 6'-*t*-Bu), 33.5 (s, 3'-*t*-Bu), 31.7 (q, 6'-*t*-Bu), 31.2 (q, 3'-*t*-Bu). HRMS (ESI): calcd for C₅₁H₅₇⁺ 669.4455, found 669.4454. Anal. calcd for C₅₁H₅₇F₆P·1/2H₂O (814.9625): C, 74.34; H, 7.09. Found: C, 74.45; H, 6.82.

3,3-Bis(3-methoxycarbonyl-1-azulenyl)-1,1-diphenylpropenylium hexafluorophosphate (11b⁺PF₆⁻). To a mixture of 3,3-bis(3-methoxycarbonyl-1-azulenyl)-1,1-diphenylpropene (**10b**) (201 mg, 0.357 mmol) and 60% HPF₆ (3 mL) in CH₂Cl₂ (30 mL) and water (30 mL) was added DDQ (97 mg, 0.43 mmol) at room temperature. The resulting mixture was stirred at room temperature for 30 min. The resulting suspension was filtered with suction. The organic layer was separated, dried with MgSO₄, and concentrated under reduced pressure. The residue was recrystallized from Et₂O/hexane to give 3,3-bis(3-methoxycarbonyl-1-azulenyl)-1,1-diphenylpropenylium hexafluorophosphate (**11b**⁺PF₆⁻) (232 mg, 92%). purple crystals. mp 189.2–190.3 °C (Et₂O/hexane). IR (KBr disk, ν_{\max} , cm⁻¹): 2932 (w), 2816 (w), 1704 (s), 1568 (w), 1537 (w), 1512 (m), 1481 (w), 1466 (s), 1446 (s), 1415 (s), 1398 (w), 1363 (s), 1294 (w), 1224 (s), 1163 (s), 1090 (m), 1067 (w), 1025 (w), 870 (m), 841 (s), 787 (w), 701 (m), 699 (w), 558 (m), 455 (w). UV-vis (CH₃CN, λ_{\max} , nm): 193 (log ϵ 4.80), 228 (4.64), 253 sh (4.55), 269 sh (4.50), 303 (4.48), 365 sh (4.09), 543 (4.27), 655 (4.40). ¹H NMR (500 MHz, (CDCl₂)₂, 80 °C): δ_{H} 9.75 (d, 2H, *J* 10.0 Hz, 4'-H), 8.69 (d, 2H, *J* 9.9 Hz, 8'-H), 8.37 (s, 2H, 2'-H), 8.18 (dd, 2H, *J* 9.8, 9.7 Hz, 6'-H), 8.07 (dd, 2H, *J* 10.0, 9.8 Hz, 5'-H), 7.83 (s, 1H, 2-H), 7.82 (dd, 2H, *J* 9.9, 9.7 Hz, 7'-H), 7.57–7.45 (m, 5H, Ph''), 6.83–6.72 (m, 5H, Ph'''), 3.90 (s, 6H, 3'-CO₂Me). ¹³C NMR (125 MHz, (CDCl₂)₂, 80 °C): δ_{C} 168.4, 164.9, 164.0, 150.9, 148.83, 148.82, 145.1, 142.0, 141.0, 140.3, 138.3, 137.7, 136.6, 132.8, 131.4, 130.8, 130.6, 130.2, 129.7, 129.6, 128.3, 124.6, 52.4. HRMS (ESI): calcd for C₃₉H₂₉O₄⁺ 561.2060, found 561.2062. Anal. calcd for C₃₉H₂₉F₆O₄P (706.6093): C, 66.29; H, 4.14. Found: C, 66.03; H, 4.38.

3,3-Bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-fluorophenyl)propenylium hexafluorophosphate (11c⁺PF₆⁻). To a solution of 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-fluorophenyl)propene (**10c**) (150 mg, 0.212 mmol) in CH₂Cl₂ (20 mL) was added DDQ (59 mg, 0.26 mmol) at room temperature. After the solution was stirred at the same temperature for 30 min, 60% HPF₆ (2 mL) and water (20 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried with MgSO₄, and concentrated under reduced pressure. The residue was crystallized from hexane to give 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-fluorophenyl)propenylium hexafluorophosphate (**11c**⁺PF₆⁻) (172 mg, 95%). deep purple crystals. mp 170.1–171.0 °C (hexane). IR (KBr disk, ν_{\max} , cm⁻¹): 2965 (m), 2900 (w), 2782 (w), 1596 (w), 1574 (w), 1545 (w), 1503 (m), 1477 (s), 1441 (m), 1416 (m), 1400 (w), 1367 (m), 1336 (s), 1312 (m), 1281 (w), 1232 (s), 1203 (w), 1184

(w), 1176 (w), 1160 (w), 1120 (w), 1105 (w), 1081 (w), 1021 (w), 991 (w), 956 (w), 869 (w), 840 (s), 810 (w), 802 (w), 705 (w), 566 (w), 558 (m). UV-vis (CH₃CN, λ_{max}, nm): 196 (log ε 4.81), 243 (4.73), 313 (4.55), 323 sh (4.55), 415 (4.17), 434 sh (4.10), 535 (4.23), 724 (4.63). ¹H NMR (500 MHz, (CDCl₂)₂, 80 °C): δ_H 8.74 (d, 2H, *J* 10.8 Hz, 4'-H), 8.42 (br, 2H, 8'-H), 7.94 (d, 2H, *J* 10.8 Hz, 5'-H), 7.68 (d, 2H, *J* 10.5 Hz, 7'-H), 7.64 (s, 1H, 2-H), 7.56 (s, 2H, 2'-H), 7.47 (dd, 2H, *J* 8.4, 5.3 Hz, 2'',6''-H or 2''',6'''-H), 7.14 (dd, 2H, *J* 8.4, 8.4 Hz, 3'',5''-H or 3''',5'''-H), 6.77 (dd, 2H, *J* 8.4, 5.3 Hz, 2'',6''-H or 2''',6'''-H), 6.44 (dd, 2H, *J* 8.4, 8.4 Hz, 3'',5''-H or 3''',5'''-H), 1.45 (s, 18H, 3'-*t*-Bu), 1.41 (s, 18H, 6'-*t*-Bu). ¹³C NMR (125 MHz, (CDCl₂)₂, 80 °C): δ_C 168.9 (C-6'), 164.6 (*J* 253.7 Hz, C-4'' or C-4'''), 162.5 (*J* 251.3 Hz, C-4'' or C-4'''), 158.6 (C-1), 155.7 (C-3), 148.5 (C-3'a), 147.9 (C-3' or C-8'a), 147.8 (C-3' or C-8'a), 141.8 (C-2'), 138.7 (C-4'), 137.9 (C-8'), 137.7 (*J* 3.0 Hz, C-1'' or C-1'''), 135.0 (*J* 3.3 Hz, C-1'' or C-1'''), 132.2 (*J* 8.7 Hz, C-2'',6'' or C-2''',6'''), 131.9 (C-5' and C-7'), 131.7 (*J* 8.4 Hz, C-2'',6'' or C-2''',6'''), 131.4 (C-1'), 130.1 (C-2), 116.6 (*J* 21.6 Hz, C-3'',5'' or C-3''',5'''), 115.2 (*J* 21.6 Hz, C-3'',5'' or C-3''',5'''), 39.5 (s, 6'-*t*-Bu), 33.4 (s, 3'-*t*-Bu), 31.8 (q, 6'-*t*-Bu), 31.2 (q, 3'-*t*-Bu). HRMS (ESI): calcd for C₅₁H₅₅F₈P·1/2H₂O (850.9434): C, 71.23; H, 6.56. Found: C, 71.25; H, 6.29.

3,3-Bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propenylium hexafluorophosphate (11d⁺PF₆⁻). To a mixture of 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propene (**10d**) (75 mg, 99 μmol) and 60% HPF₆ (2 mL) in CH₂Cl₂ (10 mL) and water (20 mL) was added DDQ (30 mg, 0.13 mmol) at room temperature. The resulting mixture was stirred at room temperature for 30 min. The resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried with MgSO₄, and concentrated under reduced pressure. The residue was recrystallized from ethanol to give 3,3-bis(3,6-di-*tert*-butyl-1-azulenyl)-1,1-bis(4-nitrophenyl)propenylium hexafluorophosphate (**11d⁺PF₆⁻**) (59 mg, 66%). deep green crystals. mp 258.2–260.5 °C (EtOH). IR (KBr disk, ν_{max}, cm⁻¹): 2965 (m), 2908 (w), 2864 (w), 1571 (w), 1519 (m), 1475 (m), 1415 (m), 1366 (m), 1335 (s), 1310 (m), 1237 (m), 1205 (w), 1178 (m), 1112 (m), 1082 (m), 1023 (w), 956 (w), 840 (s), 800 (w), 699 (w), 557 (m). UV-vis (CH₃CN, λ_{max}, nm): 194 (log ε 4.64), 235 (4.49), 250 sh (4.48), 281 (4.41), 304 (4.42), 323 (4.37), 407 (4.15), 507 (3.85), 736 (4.35). ¹H NMR (500 MHz, (CDCl₂)₂, 100 °C): δ_H 8.75 (d, 2H, *J* 10.9 Hz, 4'-H), 8.42 (d, 2H, *J* 10.7 Hz, 8'-H), 8.25 (d, 2H, *J* 8.3 Hz, 3'',5''-H or 3''',5'''-H), 8.02 (s, 1H, 2-H), 7.98 (d, 2H, *J* 10.9 Hz, 5'-H), 7.79 (d, 2H, *J* 10.7 Hz, 7'-H), 7.64 (d, 2H, *J* 8.3 Hz, 2'',6''-H or 2''',6'''-H), 7.55 (d, 2H, *J* 8.1 Hz, 3'',5''-H or 3''',5'''-H), 7.55 (s, 2H, 2'-H), 6.89 (d, 2H, *J* 8.1 Hz, 2'',6''-H or 2''',6'''-H), 1.46 (s, 18H, 3'-*t*-Bu), 1.43 (s, 18H, 6'-*t*-Bu). ¹³C NMR (125 MHz, (CDCl₂)₂, 100 °C): δ_C 169.9 (C-6'), 153.8 (C-1), 152.9 (C⁺), 149.3 (C-4'' or C-4'''), 149.0 (C-3' or C-3'a), 148.9 (C-3' or C-3'a), 148.1 (C-8'a), 147.3 (C-4'' or C-4'''), 146.1 (C-1'' or C-1'''), 144.8 (C-1'' or C-1'''), 141.4 (C-2'), 138.8 (C-4'), 138.4 (C-8'), 135.1 (C-2), 133.0 (C-7'), 132.8 (C-5'), 131.8 (C-1'), 130.7 (C-2'',6'' or C-2''',6'''), 130.1 (C-2'',6'' or C-2''',6'''), 124.6 (C-3'',5'' or C-3''',5'''), 122.9 (C-3'',5'' or C-3''',5'''), 39.6 (s, 6'-*t*-Bu), 33.5 (s, 3'-*t*-Bu), 31.7 (q, 6'-*t*-Bu), 31.2 (q, 3'-*t*-Bu). HRMS (ESI): calcd for C₅₁H₅₅N₂O₄⁺ 759.4156, found 759.4154. Anal. Calcd for C₅₁H₅₅F₆N₂O₄P (904.9576): C, 67.69; H, 6.13; N, 3.10. Found: C, 67.68; H, 5.92; N, 3.06.

9-[2,2-Bis(3,6-di-*tert*-butyl-1-azulenyl)ethylidene]-9H-fluorene (13). A mixture of 9H-fluoren-9-ylidene acetaldehyde (**12**) (99 mg, 0.48 mmol) and 1,6-di-*tert*-butylazulene (**3a**) (228 mg, 0.948 mmol) in acetic acid (6 mL) was stirred at room temperature for 24 h under an Ar atmosphere. The reaction mixture was poured into water and extracted with CH₂Cl₂. The organic layer was washed with 5% NaHCO₃ and water, dried over MgSO₄, and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel with CH₂Cl₂/hexane (1:5) and recrystallization from Et₂O/hexane to afford 9-[2,2-bis(3,6-di-*tert*-butyl-1-azulenyl)ethylidene]-9H-fluorene (**13**) (204 mg, 64%). blue crystals. mp 255.4–256.7 °C (Et₂O/hexane). IR (KBr disk, ν_{max}, cm⁻¹): 3060 (w), 2963 (s), 2904 (m), 2868 (m), 1576 (s), 1508 (w), 1475 (w), 1455 (m), 1443 (m), 1439 (w), 1424 (m), 1390 (m), 1361 (s), 1294 (w), 1245 (m), 1227 (m), 1198 (w), 1070 (w), 872 (w), 833 (s), 823 (w), 776 (s), 730 (s), 669 (w), 614 (w), 528 (w), 442 (s). UV-vis (CH₂Cl₂, λ_{max}, nm): 239 (log ε 4.69), 249 (4.65), 259

(4.65), 293 (4.80), 299 (4.80), 343 sh (4.04), 355 (4.07), 375 (4.00), 564 sh (2.74), 610 (2.86), 673 sh (2.75), 750 sh (2.21). ^1H NMR (500 MHz, CDCl_3): δ_{H} 8.56 (d, 2H, J 10.7 Hz, 4-H), 8.19 (d, 2H, J 10.6 Hz, 8-H), 7.90 (d, 1H, J 7.8 Hz, 8'-H), 7.76 (d, 1H, J 7.4 Hz, 5'-H), 7.71 (d, 1H, J 7.6 Hz, 4'-H), 7.70 (d, 1H, J 7.7 Hz, 1'-H), 7.58 (s, 2H, 2-H), 7.33 (dd, 1H, J 7.5, 7.4 Hz, 6'-H), 7.30 (dd, 1H, J 7.6, 7.4 Hz, 3'-H), 7.29 (d, 1H, 10'-H), 7.24 (dd, 1H, J 7.8, 7.4 Hz, 2'-H), 7.21 (dd, 1H, J 7.8, 7.5 Hz, 7'-H), 7.17 (dd, 2H, J 10.7, 1.8 Hz, 5-H), 7.03 (dd, 2H, J 10.6, 1.8 Hz, 7-H), 6.99 (d, 1H, CH), 1.51 (s, 18H, 3-*t*-Bu), 1.39 (s, 18H, 6-*t*-Bu). ^{13}C NMR (125 MHz, CDCl_3): δ_{C} 160.5 (C-6), 141.0 (C-4'b), 134.0 (C-9'a), 138.8 (C-4'a), 137.9 (C-3), 137.3 (C-8'a), 134.9 (C-8a), 134.8 (C-2), 134.5 (C-4), 134.3 (C-3a), 133.9 (C-10'), 133.3 (C-9'), 132.2 (C-8), 129.6 (C-1), 127.6 (C-6'), 127.3 (C-3'), 127.2 (C-7'), 126.7 (C-2'), 125.4 (C-8'), 120.4 (C-1'), 119.7 (C-5'), 119.5 (C-7), 119.4 (C-4'), 118.4 (C-5), 38.2 (s, 6-*t*-Bu), 36.9 (CH), 33.3 (s, 3-*t*-Bu), 32.2 (q, 3-*t*-Bu), 31.8 (q, 6-*t*-Bu). HRMS (ESI): calcd for $\text{C}_{51}\text{H}_{56} + \text{Na}^+$ 691.4274, found 691.4272. Anal. calcd for $\text{C}_{51}\text{H}_{56}$ (668.9903): C, 91.56; H, 8.44. Found: C, 91.30; H, 8.53.

1,1-Bis(3,6-di-*tert*-butyl-1-azulenyl)-2-(9*H*-fluorenylidene)ethylium hexafluorophosphate (14^+PF_6^-). To a solution of 9-[2,2-bis(3,6-di-*tert*-butyl-1-azulenyl)ethylidene]-9*H*-fluorene (**13**) (121 mg, 0.18 mmol) in CH_2Cl_2 (20 mL) was added DDQ (56 mg, 0.25 mmol) at room temperature. After the solution was stirred at the same temperature for 30 min, 60% HPF_6 (2 mL) and water (20 mL) were added to the mixture. After stirring the mixture for an additional 10 min, the resulting suspension was filtered with suction. The organic layer was separated, washed with water, dried with MgSO_4 , and concentrated under reduced pressure. The residue was recrystallized from Et_2O /hexane to give 1,1-bis(3,6-di-*tert*-butyl-1-azulenyl)-2-(9*H*-fluorenylidene)ethylium hexafluorophosphate (14^+PF_6^-) (130 mg, 88%). deep purple powder. mp 182.2–186.3 °C (Et_2O /hexane). IR (KBr disk, ν_{max} , cm^{-1}): 2964 (m), 2898 (w), 2871 (w), 1576 (m), 1544 (w), 1478 (s), 1450 (w), 1416 (s), 1396 (w), 1367 (m), 1332 (s), 1311 (s), 1279 (w), 1240 (m), 1205 (m), 1184 (m), 1178 (m), 1112 (m), 1082 (m), 1025 (w), 990 (w), 953 (w), 869 (m), 841 (s), 806 (m), 782 (w), 726 (w), 705 (w), 558 (m). UV-vis (CH_3CN , λ_{max} , nm): 206 (log ϵ 4.68), 230 (4.76), 248 (4.80), 254 sh (4.79), 280 (4.63), 317 (4.47), 412 (4.14), 446 (4.02), 544 (4.15), 729 (4.49). ^1H NMR (500 MHz, $(\text{CDCl}_2)_2$, 80 °C): δ_{H} 8.90 (d, 2H, J 11.1 Hz, 4-H), 8.55 (d, 2H, J 10.4 Hz, 8-H), 8.04 (s, 1H, 10'-H), 8.03 (dd, 2H, J 11.1, 1.9 Hz, 5-H), 7.86 (d, 1H, J 7.6 Hz, 1'-H), 7.72 (s, 2H, 2-H), 7.68 (dd, 2H, J 10.4, 1.9 Hz, 7-H), 7.62 (d, 1H, J 7.4 Hz, 4'-H), 7.49 (d, 1H, J 7.4 Hz, 5'-H), 7.46 (dd, 1H, J 7.5, 7.4 Hz, 3'-H), 7.37 (dd, 1H, J 7.6, 7.5 Hz, 2'-H), 7.08 (dd, 1H, J 7.5, 7.4 Hz, 6'-H), 6.50 (dd, 1H, J 7.9, 7.5 Hz, 7'-H), 5.89 (d, 1H, J 7.9 Hz, 8'-H), 1.47 (s, 18H, 3-*t*-Bu), 1.41 (s, 18H, 6-*t*-Bu). ^{13}C NMR (125 MHz, $(\text{CDCl}_2)_2$, 80 °C): δ_{C} 169.4 (C-6), 153.5 (C⁺), 151.4 (C-9'), 149.2 (C-3a), 148.9 (C-3), 147.9 (C-8a), 143.1 (C-4'b), 142.5 (C-2), 140.9 (C-4'a), 139.2 (C-4), 139.1 (C-9'a), 138.2 (C-8), 135.5 (C-8'a), 132.6 (C-5), 132.3 (C-7), 132.0 (C-1), 131.9 (C-3'), 131.1 (C-6'), 128.4 (C-2'), 127.4 (C-7'), 126.4 (C-8'), 126.2 (C-10'), 122.6 (C-1'), 120.6 (C-4'), 120.5 (C-5'), 39.6 (s, 6-*t*-Bu), 33.6 (s, 3-*t*-Bu), 31.7 (q, 3-*t*-Bu), 31.4 (q, 6-*t*-Bu). HRMS (ESI): calcd for $\text{C}_{51}\text{H}_{55}^+$ 667.4298, found 667.4300. Anal. calcd for $\text{C}_{51}\text{H}_{55}\text{F}_6\text{P}$ (812.9466): C, 75.35; H, 6.82. Found: C, 75.64; H, 6.69.

Acknowledgements

The present work was supported by JSPS KAKENHI Grant Number JP24550037.

Supplementary Material

Experimental details for physical measurements, pK_{R}^+ data, CV and DVP waves, and spectroelectrograms of **5a⁺**, **5b⁺**, **8a⁺**, **8b⁺**, **11a–d⁺**, and **14⁺**, and copies of ^1H and ^{13}C NMR spectra of reported compounds.

References

1. Monk, P. M. S.; Mortimer, R. J.; Rosseinsky, D. R. *Electrochromism: Fundamentals and Applications*; VCH: Weinheim, Germany, 1995.
<https://doi.org/10.1002/9783527615377>
2. Mortimer, R. J. *Chem. Soc. Rev.* **1997**, 26, 147-156.
<https://doi.org/10.1039/cs9972600147>
3. Deuchert, K.; Hünig, S. *Angew. Chem., Int. Ed. Engl.* **1978**, 17, 875-886.
<https://doi.org/10.1002/anie.197808753>
4. Hünig, S.; Kemmer, M.; Wenner, H.; Perepichka, I. F.; Bäuerle, P.; Emge, A.; Gescheid, G. *Chem. Eur. J.* **1999**, 5, 1969-1973.
[https://doi.org/10.1002/\(SICI\)1521-3765\(19990702\)5:7<1969::AID-CHEM1969>3.0.CO;2-5](https://doi.org/10.1002/(SICI)1521-3765(19990702)5:7<1969::AID-CHEM1969>3.0.CO;2-5)
5. Hünig, S.; Kemmer, M.; Wenner, H.; Barbosa, F.; Gescheidt, G.; Perepichka, I. F.; Bäuerle, P.; Emge, A.; Peters, K. *Chem. Eur. J.* **2000**, 6, 2618-2632.
[https://doi.org/10.1002/1521-3765\(20000717\)6:14<2618::AID-CHEM2618>3.0.CO;2-W](https://doi.org/10.1002/1521-3765(20000717)6:14<2618::AID-CHEM2618>3.0.CO;2-W)
6. Hünig, S.; Perepichka, I. F.; Kemmer, M.; Wenner, H.; Bäuerle, P.; Emge, A. *Tetrahedron* **2000**, 56, 4203-4211.
[https://doi.org/10.1016/S0040-4020\(00\)00345-8](https://doi.org/10.1016/S0040-4020(00)00345-8)
7. Hünig, S.; Langels, A.; Schmittel, M.; Wenner, H.; Perepichka, I. F.; Peters, K. *Eur. J. Org. Chem.* **2001**, 1393-1399.
[https://doi.org/10.1002/1099-0690\(200104\)2001:7<1393::AID-EJOC1393>3.0.CO;2-Z](https://doi.org/10.1002/1099-0690(200104)2001:7<1393::AID-EJOC1393>3.0.CO;2-Z)
8. Ito, S.; Morita, N.; Kubo, T. *J. Syn. Org. Chem.* **2004**, 62, 766-777.
<https://doi.org/10.5059/yukigoseikyokaishi.62.766>
9. Ito, S.; Morita, N. *Eur. J. Org. Chem.* **2009**, 4567-4579.
<https://doi.org/10.1002/ejoc.200900393>
10. Ito, S.; Morita, N.; Asao, T. *Bull. Chem. Soc. Jpn.* **1995**, 68, 1409-1436.
<https://doi.org/10.1246/bcsj.68.1409>
11. Ito, S.; Kobayashi, H.; Kikuchi, S.; Morita, N.; Asao, T. *Bull. Chem. Soc. Jpn.* **1996**, 69, 3225-3237.
<https://doi.org/10.1246/bcsj.69.3225>
12. Ito, S.; Kikuchi, S.; Morita, N.; Asao, T. *Bull. Chem. Soc. Jpn.* **1999**, 72, 839-849.
<https://doi.org/10.1246/bcsj.72.839>
13. Ito, S.; Kikuchi, S.; Morita, N.; Asao, T. *J. Org. Chem.* **1999**, 64, 5815-5821.
<https://doi.org/10.1021/jo990029p>
14. Ito, S.; Morita, N.; Asao, T. *Bull. Chem. Soc. Jpn.* **2000**, 73, 1865-1874.
<https://doi.org/10.1246/bcsj.73.1865>
15. Ito, S.; Kubo, T.; Kondo, M.; Kabuto, C.; Morita, N.; Asao, T.; Fujimori, K.; Watanabe, M.; Harada, N.; Yasunami, M. *Org. Biomol. Chem.* **2003**, 1, 2572-2580.
<https://doi.org/10.1039/b302688d>
16. Ito, S.; Kawakami, J.; Tajiri, A.; Ryuzaki, D.; Morita, N.; Asao, T.; Watanabe, M.; Harada, N. *Bull. Chem. Soc. Jpn.* **2005**, 78, 2051-2065.
<https://doi.org/10.1246/bcsj.78.2051>
17. Ito, S.; Akimoto, K.; Kawakami, J.; Tajiri, A.; Shoji, T.; Satake, H.; Morita, N. *J. Org. Chem.* **2007**, 72, 162-172.
<https://doi.org/10.1021/jo0618324>

18. Stella, L.; Janousek, Z.; Merényi, R.; Viehe, H. G. *Angew. Chem., Int. Ed. Engl.* **1978**, *17*, 691-692.
<https://doi.org/10.1002/anie.197806911>
19. Viehe, H. G.; Merényi, R.; Stella, L.; Janousek, Z. *Angew. Chem., Int. Ed. Engl.* **1979**, *18*, 917-932.
<https://doi.org/10.1002/anie.197909171>
20. Viehe, H. G.; Janousek, Z.; Merényi, R. *Acc. Chem. Res.* **1985**, *18*, 148-154.
<https://doi.org/10.1021/ar00113a004>
21. Ito, S.; Kubo, T.; Morita, N.; Ikoma, T.; Tero-Kubota, S.; Tajiri, A. *J. Org. Chem.* **2003**, *68*, 9753-9762.
<https://doi.org/10.1021/jo035053o>
22. Takahashi, K.; Nihira, T. *Tetrahedron Lett.* **1989**, *30*, 5903-5906.
[https://doi.org/10.1016/S0040-4039\(01\)93501-4](https://doi.org/10.1016/S0040-4039(01)93501-4)
23. Takahashi, K.; Nihira, T.; Tomitani, K. *J. Chem. Soc., Chem. Commun.* **1993**, 1617-1619.
<https://doi.org/10.1039/c39930001617>
24. Ito, S.; Kikuchi, S.; Okujima, T.; Morita, N.; Asao, T. *J. Org. Chem.* **2001**, *66*, 2470-2479.
<https://doi.org/10.1021/jo001709r>
25. Shoji, T.; Higashi, J.; Ito, S.; Okujima, T.; Morita, N. *Eur. J. Org. Chem.* **2011**, 584-592.
<https://doi.org/10.1002/ejoc.201001212>
26. Hosoya, T.; Aoyama, H.; Ikemoto, T.; Kihara, Y.; Hiramatsu, T.; Kondo, M.; Suzuki, M. *Bioorg. Med. Chem.* **2003**, *11*, 663-673.
[https://doi.org/10.1016/S0968-0896\(02\)00600-4](https://doi.org/10.1016/S0968-0896(02)00600-4)
27. Bharathi, P.; Periasamy, M. *Org. Lett.* **1999**, *1*, 857-859.
<https://doi.org/10.1021/ol990745d>
28. Adams, D. J.; Clark, J. H.; McFarland, H. *J. Fluorine Chem.* **1998**, *92*, 127-129.
[https://doi.org/10.1016/S0022-1139\(98\)00274-7](https://doi.org/10.1016/S0022-1139(98)00274-7)