

Supplementary Material

Palladium complexes with 3-methyl-8-thia-1,3-diaza-spiro[4.5]decane-2,4-dione: experimental and theoretical studies

Emiliya D. Cherneva, Rossen T. Buyukliev, and Adriana G. Bakalova

*Department of Chemistry, Faculty of Pharmacy, Medical University - Sofia, 2 Dunav Str., 1000
Sofia, Bulgaria*

Email: e.d.cherneva@gmail.com

Table of Contents

- | | |
|---|------|
| 1. ^1H and ^{13}C NMR spectra of compound 3-Methyl-8-thia-1,3-diaza-spiro[4.5]decane-2,4-dione (2) | S2-3 |
| 2. ^1H NMR spectrum of complex <i>cis</i> -bis(3-methyl-8-thia-1,3-diaza-spiro[4.5]decane-2,4-dione)-dichlorido palladium(II) – <i>cis</i> -[PdL ₂ Cl ₂] (3) | S4 |

1. **Figure S2:** ^1H NMR spectrum of compound 3-Methyl-8-thia-1,3-diaza-spiro[4.5]decane-2,4-dione (**2**)





Figure S3: ^{13}C NMR spectrum of compound **2**

Figure S4: ^1H NMR spectrum of complex *cis*-bis(3-methyl-8-thia-1,3-diaza-spiro[4.5]decane-2,4-dione)-dichlorido palladium(II) – *cis*-[PdL₂Cl₂] (**3**)

