

Supplementary Material

Synthesis of 1,4- and 1,4,4-substituted piperidines for the inhibition of neuronal T-type Ca²⁺ channels and mitigation of neuropathic pain in mice

Medha J. Gunaratna,^a Duy H. Hua,^{*a} Bende Zou,^b Conrado Pascual,^b William Cao,^b Man Zhang,^a Sahani Weerasekara,^a Thi D. T. Nguyen,^a Kui Xiao,^b and Xinmin Simon Xie^{*b}

^aDepartment of Chemistry, Kansas State University, Manhattan, KS 66506, U.S.A.

^bAfaSci Research Laboratories, Redwood City, CA, U.S.A.

Email: duy@ksu.edu; simonxie@afasci.com

Table of Contents

Biological evaluation of T-type channel inhibitors	S2
Biological Experimental Section	S5
References	S6
¹ H and ¹³ C NMR and MS spectra.....	S7

Biological evaluation of T-type channel inhibitors

For the primary screen of our synthesized molecules for T-type channel activities, we used patch clamp recordings of either dissociated mouse dorsal root ganglion (DRG) neurons or $Ca_{v3.2}$ channel expressed in human embryonic kidney (HEK)-293 cells from a tetracycline inducible expression system (Invitrogen; see Experimental Section shown below). The active compounds were then evaluated *in vivo* for resistance of neuropathic pain by either one of the following two methods. Firstly, because knockout or antisense knockdown of the $Ca_{v3.2}$ isoform produces analgesic effects in neuropathic and inflammatory pain in mice,¹ a reported Complete Freund adjuvant (CFA)-inflammatory pain mouse model¹ was used to examine the drug effects. Secondly, because an over activation of T-channel results in the generation of seizure activity,² prolonged latencies to seizures and decreasing the fatality rate in pentylenetetrazole (PTZ)-induced seizure mouse model was also used to determine the inhibition of T-channel activity. Two previously reported T-type calcium channel inhibitors ML218 and Z944 were compared, and gabapentin was used as a standard in the analgesic study. Results of the *in vitro* and *in vivo* studies are summarized in Table S1. From the *in vitro* inhibition of T-type and $Ca_{v3.2}$ channel data (second and third columns of Table S1), the synthesized molecules are not as strong inhibitors as ML218 or Z944 with the exception of molecule **6**, which possessed similar activity in DRG neurons. However, because molecule **6** showed a weak inhibitory activity in PTZ-induced fatality model, it was not investigated further. Compounds **2**, **5**, and **6** possessed low inhibitions of seizure; therefore, they were also not investigated further. Notably, the two reported calcium channel inhibitors ML218 and Z944 showed only weak protective effect in PTZ-induced fatality model (17 and 33%, respectively). Compounds **3**, **7**, **9**, and **10** did not show promising inhibition of either T-channels or $Ca_{v3.2}$ channel; hence, they were not studied further. Molecules **4** and **8** displayed acceptable analgesic activity (strong and medium, respectively) and inhibition of seizure (86 and 89%, respectively). Therefore, these two compounds were investigated further for their analgesic effects using a spared nerve injury (SNI) model, which is an acute pain and neurogenic inflammatory pain rat model.

Table S1. Bioactivities *in vitro* and *in vivo* of molecules **1** – **10** and ML218 and Z944. (NT = not tested)

Compound	Inhibition (%) of T-channel in DRG neurons at 1 μ M (unless specified)	Inhibition (%) of Ca _v 3.2-channel in HEK 293 cells at 1 μ M (unless specified)	Analgesic Max: ++++ at 30 mg/Kg, (unless specified) (n = 6 - 8)	Inhibition (%) of seizure in PTZ-induced fatality, at 30 mg/Kg, via i.p. (n = 6)
1	43.6 \pm 13.2 (n=3)	NT	+	80
2	47	NT	NT	0
3	NT	40 (at 10 μ M)	NT	NT
4	NT	62.6 (n=2)	+++	86
5	83 \pm 6.4 (n=3)	NT	+	0
6	100 (n = 2)	NT	NT	14
7	43.6 (n=2)	NT	NT	63
8	40.1 \pm 6.9 (n=7)	47 (n=2)	++	89
9	0	NT	NT	NT
10	0	NT	NT	NT
ML218	99	91.8 \pm 1.8 (n=3)	+++	17
Z944	93.7 (n=2)	82.8 \pm 3,3 (n=19)	+++	33
Gabapentin	NT	NT	+++ (100 mg/kg)	0

The tested compounds **4** and **8** showed increases of thermal and mechanical thresholds compared to vehicle. The drug actions started around one hour post-injection and peaked between 3 and 4 hours. The drug effects gradually declined to pre-treatment levels over 7-8 hours (Figures S1 and S2). These tested compounds produced analgesic effects on both thermal and mechanical pain thresholds, while the Spared Nerve Injury (SNI)-vehicle group remained thermal hyperalgesia and mechanical allodynia conditions. Therefore, compounds **4** and **8** have the ability to mitigate neuropathic pain induced in rats and **4** appears to have a greater effect than **8** at a 3-hour time point (an 80% increase in mechanical threshold for **4** and ~60% for **8**)

In the SNI studies, baseline thermal and mechanical thresholds of male rats (Sprague-Dawley rats) were taken prior to surgery, and baseline thresholds were measured on the rats' left hind paws. A group of rats underwent surgery without injury of the nerves, i.e., a sham surgery group (n = 6 – 8/group). Two weeks post-surgery, the thermal and mechanical pain thresholds of each rat were assessed at pre-dosing, and 1, 2, 3, 4, 5, 6, 7, and 8 hours post-treatment (of drug) using Hargreaves test and von Frey monofilament test, respectively. Results are depicted in Figures S1 and S2.




Figure S1. Spared Nerve Injury (SNI) rats' left hind paw withdrawal latencies in response to thermal stimulation (Hargreaves test; number of rats: n = 6 - 8).




Figure S2. Mechanical pain threshold assessment (von Frey monofilament test) of SNI rats' left hind paws (number of rats: n = 6 - 8).

Biological Experimental Section.

Bio-evaluation methods

Inhibition of T-channel in DRG neurons. Mouse dorsal root ganglion (DRG) neurons were prepared from 1 – 2 months old C57/BC6 mice. All the procedures related to animal handling were conducted at AfaSci Research Laboratory and were in strict accordance with NIH guidelines and IACUC approved protocols. The spine was removed and split into two halves from the middle line after sacrificing the mice by decapitation. Lumbar DRG neurons were collected into a modified Krebs solution (130 mM NaCl, 10 mM HEPES-Na, 5 mM KCl, 1 mM CaCl₂, 10 mM glucose, 2 mM MgCl₂, pH adjusted to 7.35 with 1 N HCl) in a 1.5 mL-tube. The DRG neurons were removed for digestion into 0.5 mL of Hank's balanced salt solution (HBSS) with 1 mg/mL Collagenase and 0.5 mg/mL Trypsin added. The DRG neurons were minced with fine scissors and incubated at 35°C for 50 min. After removing the HBSS solution, the DRG neurons were dispersed into modified Krebs solution and triturated with fire polished glass pipettes until no clumps were visible. The cells were then dispersed onto poly-L-ornithine-coated cover slips and maintained in a modified Krebs solution and streptomycin sulfate (0.2 mM), Penicillin G Sodium (0.3 mM), and Gentamycin (0.1 mM) at 21°C.

Preparation of HEK293 cells expressing Ca_{v3.2} channels. Ca_{v3.2} channels were expressed in human embryonic kidney (HEK)-293 cells using a tetracycline inducible expression system (Invitrogen). Cells were placed in 25-cm² tissue culture flasks at 37°C, 5% CO₂, and 100% relative humidity in D-MEM/F12 (Dulbecco's Modified Eagle Medium/Nutrient Mixture F-12; Gibco) supplemented with fetal bovine serum (FBS, 10% v/v), sodium pyruvate (0.5 mM, Gibco), penicillin-streptomycin (100 U/mL, 100 µg/mL) and Geneticin^R Selective antibiotic (G418; 0.5 mg/mL). Cells were detached from the flask base using a non-enzymatic cell dissociation solution (CellStripper, Corning), removed, and reseeded onto poly-D-lysine coated glass coverslips in 35-mm dishes. Cells in dishes were further supplemented with G418 at a final concentration of 1 mg/mL and after 24 hours, expression of Ca_{v3.2} was induced with 1 µg/mL tetracycline. The medium for the HEK-293 was composed of D-MEM/F-12 and 15 mM HEPES buffer, with L-glutamine and pyridoxine hydrochloride. It was supplemented with FBS, P/S (penicillin/streptomycin) (Gibco, #15140-122), and sodium pyruvate 100 mM.

Patch clamp recordings. Whole-cell voltage clamp recordings were performed on mouse DRG neurons within 2 days after acutely dissociation or cultured HEK293 cells expressing T-type channels (encoded by Ca_{v3.2} channels). All experiments were performed at ~21°C and whole-cell currents were recorded using a MultiClamp 700B amplifier and analyzed off-line with Pclamp10.4 software (Molecular Devices, LLC, Sunnyvale, CA, USA). To record calcium currents in DRG and HEK293 cells, the external solution was composed (in mM) of 115 choline-Cl, 30 TEA-Cl, 2 CaCl₂, 10 glucose and 10 HEPES (pH 7.3 – 7.4 adjusted with TEA-OH; osmolality verified as 295 mOsm/kg). The internal solution was composed (in mM) of 125 CsCl, 10 HEPES (acid), 10 EGTA (ethylene glycol tetraacetic acid), 1 CaCl₂, 1 MgCl₂, 4 MgATP, and 0.3 MgGTP (pH 7.3 – 7.4 adjusted with CsOH; osmolality verified as 295 mOsm/kg). Since sodium ions were absent in the external solution, tetrodotoxin was not needed. Calcium currents were recorded at a holding potential of -100 mV and then depolarized to -30 mV for 100 ms to activate Ca_{v3.2} expressed in HEK cells. An interpulse interval of 10 seconds allowed the channel recovery from inactivation and achieved stable current recordings. All reagents were purchased from Sigma unless specified otherwise. Test compounds were usually applied via a rapid-solution-exchange system with 8 fine polyplastic tubings (100 mm OD) glued in a holder in parallel and located closely to the recorded cells. The current responses were normalized to the control, percent inhibition was

calculated, and sigmoidal dose-response curves were generated using XLFit (IDBS, Surrey, UK) or Prism (GraphPad Software, La Jolla, CA, US) for calculation.

Analgesic effect using Complete Freund adjuvant (CFA)-inflammatory pain mouse model. The reported protocol⁴ of Complete Freund adjuvant (CFA)-inflammatory pain in mice was followed for the study of inhibition of pains. Tested molecules and a reference, gabapentin (100 mg/kg, I.P.), was individually administered to mice (n = 6 - 8) that had CFA-induced inflammatory on their left hind paws. The maximum inhibition was rated at +++++, strong inhibition at +++, medium at ++, and weak at +, comparing with gabapentin.

Inhibition of seizure in pentylenetetrazol (PTZ)-induced fatality in mice. It has been reported that antiepileptics, such as carbamazepine, lamotrigine and gabapentin, are used for the treatment of neuropathic pain.¹ Overactivation of T-channel involves the generation of seizure activity.¹ Following the reported protocol² in the pentylenetetrazol (PTZ)-induced seizure model in mice (n = 6 per drug) was used to evaluate the inhibition of pains. Prolonged latencies to seizure and decrease of fatality rate were studied comparing to vehicle (2% DMSO in 0.5% HPC). Fatality latency and rate – the percentage (%) of mice in each treatment group that died within a 20-minutes cutoff of the observational period, were recorded and calculated. Average fatality latency and rate of each treatment group were used to evaluate the molecule's ability to either prevent or delay the onset of PTZ-induced seizures and death.

In vivo analgesic effects in spared nerve injury (SNI) pain models. To evaluate the therapeutic effects and safety of highly active compounds, the spared nerve injury (SNI) model, one of the reliable neuropathic pain models in mice or rats,³ was used to assess the effects of compounds **4** and **8**. Baseline thermal and mechanical thresholds of male rats (Sprague-Dawley rats, 275 – 325 g, Harlan) were taken prior to surgery. Baseline thresholds were measured on the rats' left hind paws. Drug-tested rats (n = 6 - 8 per drug) and without-drug-treated rats underwent the SNI surgery on their left hind leg. A group of rats (n = 6 – 8/group) underwent surgery without injury of the nerves as a sham group. Two weeks post-surgery, the thermal thresholds of each rat were reassessed. The SNI rats displayed approximate 55% reduction in thermal pain thresholds and 75 – 80% reduction in mechanical pain thresholds, confirming the presence of hyperalgesia and mechanical allodynia in the SNI rats. Each rat received either compounds **4**, **8** (each 30 mg/kg, 1 mL I.P.) or vehicle (2% DMSO in 0.5% HPC). Thermal and mechanical thresholds were measured at 1, 2, 3, 4, 5, 6, 7, and 8 hours post-treatment/dosing. The tested compounds were shown to elevate both thermal and mechanical thresholds compared to vehicle.

References.

1. Perez-Reyes, E. *Mol. Pharm.* **2010**, *77*, 136.
[https://doi: 10.1124/mol.109.062133](https://doi:10.1124/mol.109.062133)
2. Xie, X.; Lancaster, B.; Peakman, T.; Garthwaite, J. *Pflugers Archiv.: Eur. J. Phys.* **1995**, *430*, 437.
<https://doi:10.1007/BF00373920>
3. Decosterd, I.; Woolf, C. J. *Pain* **2000**, *87*, 149.
[https://doi.org/10.1016/S0304-3959\(00\)00276-1](https://doi.org/10.1016/S0304-3959(00)00276-1)

4. Choi, S.; Na, H.; Kim, J.; Lee, J.; Lee, S.; Kim, D.; Park, J.; Chen, C. C.; Campbell, K.; Shin, H. S. *Genes Brain Behav.* **2007**, *6*, 425.

<https://doi.org/10.1111/j.1601-183X.2006.00268.x>




Figure S3. ¹H NMR spectrum of compound **1**.




Figure S4. ¹³C NMR spectrum of compound **1**.





Figure S5. MS spectrum of compound 1.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₂₁ H ₂₅ N ₃ O ₃	FW	367.4415
---------	---	----	----------



2

MG06-011-DP-PROTON-DMSO




J:MG06-011-DP-PROTON-DMSO

Figure S6. ¹H NMR spectrum of compound 2.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₂₁ H ₂₅ N ₃ O ₃	FW	367.4415
---------	---	----	----------



MG06-011-DP-CARBON.ESP

2



I:\T-TYPE CALCIUM CHANNEL\NMR\1-2\MG06-011-DP-CARBON.ESP

Figure S7. ¹³C NMR spectrum of compound 2.

+Q1: 8 MCA scans from Sample 1 (TuneSampleID) of MT20140708214256.wiff (Turbo Spray)

Max. 1.5e6 cps.





Figure S8. MS spectrum of compound 2.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₁₈ H ₂₃ N ₃ O ₂	FW	353.4182
---------	---	----	----------



3

MG07-023-DP-H




J:\NOTEBOOK7\MG07-023-DP-H

Figure S9. ¹H NMR spectrum of compound 3.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₁₉ H ₂₃ N ₃ O ₂	FW	353.4182
---------	---	----	----------



3

MG07-023-C13



J:\NOTEBOOK7\MG07-023-C13

Figure S10. ¹³C NMR spectrum of compound **3**.




Figure S11. MS spectrum of compound **3**.




Figure S12. ¹H NMR spectrum of compound 4.




Figure S13. ^{13}C NMR spectrum of compound 4.







Figure S14. MS spectrum of compound 4.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/Formula C₂₂H₂₇N₃O₂ FW 365.4687


5

Figure S15. ¹H NMR spectrum of compound 5.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/Formula C₂₂H₂₇N₃O₂ FW 365.4687


SW-5-77DPCARBON

Figure S16. ¹³C NMR spectrum of compound 5.

**Figure S17.** MS spectrum of compound **5**.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₂₂H₂₇N₃O₃ FW 381.4681



MG05-050-FRAC5-6




Figure S18. ¹H NMR spectrum of compound 6.





Figure S19. MS spectrum of compound **6**.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₂₂H₂₆FN₃O₃ FW 399.4585



7

MG079-FRAC19-22





Figure S20. ¹H NMR spectrum of compound 7.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₂₂H₂₆FN₃O₃ FW 399.4585



7

MG05-079-FRA19-22




Figure S21. ¹³C NMR spectrum of compound 7.





Figure S22. MS spectrum of compound 7.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₉H₂₉N₃O₂ FW 331.4525



8

TN-11-35-NCE16





Figure S23. ¹H NMR spectrum of compound 8.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₉H₂₉N₃O₂ FW 331.4525



TN-11-35-DP-C13






Figure S24. ¹³C NMR spectrum of compound **8**.

Figure S25. MS spectrum of compound **8**.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₂₉H₃₁N₃O₄ FW 485.5741



9

MG05-086-FRAC11-12




J:\T-TYPE CALCIUM CHANNEL\NMR\1-9\MG05-086-FRAC11-12

Figure S26. ¹H NMR spectrum of compound 9.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₂₉ H ₃₁ N ₃ O ₄	FW	485.5741
---------	---	----	----------



9

MG05-086-C13

Figure S27. ¹³C NMR spectrum of compound 9.






Figure S28. MS spectrum of compound **9**.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/Formula C₂₉H₃₀FN₃O₄ FW 503.5646


10

MG05-060-FRAC7-9




J:\T-TYPE CALCIUM CHANNEL\NMR\1-10\MG05-060-FRAC7-9

Figure S29. ¹H NMR spectrum of compound 10.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/Formula C₂₉H₃₀FN₃O₄ FW 503.5646

10

MG05-060-13C

Figure S30. ¹³C NMR spectrum of compound 10.





Figure S31. MS spectrum of compound **10**.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₄H₁₇N₃O FW 243.3043



13

MG05-090-FRAC10-14





Figure S32. ¹H NMR spectrum of compound 13.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₄H₁₇N₃O FW 243.3043



13

MG05-090-DP-C13.ESP




Figure S33. ¹³C NMR spectrum of compound **13**.





Figure S34. MS spectrum of compound **13**.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₄H₂₁N₃O FW 247.3360



14





Figure S35. ¹H NMR spectrum of compound 14.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₄H₂₁N₃O FW 247.3360



14

MG05-093-C13



J:\NOTEBOOK5\MG05-093-C13

Figure S36. ¹³C NMR spectrum of compound 14.





Figure S37. MS spectrum of compound 14.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₇ H ₁₂ N ₂	FW	124.1836
---------	---	----	----------



SW-5-64FRAC10





Figure S38. ¹H NMR spectrum of compound 17.


+Q1: 6 MCA scans from Sample 1 (TuneSampleID) of MT20140312192659.wiff (Turbo Spray)

Max. 6.4e6 cps.

Figure S39. MS spectrum of compound **17**.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₈H₁₉N₃O FW 257.3309



18




Figure S40. ¹H NMR spectrum of compound 18.




Figure S41. MS spectrum of compound 18.





Figure S42. MS spectrum of compound 19.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₁H₁₈N₂O₂ FW 210.2728



20

MG07-026-DP




J:\NOTEBOOK7\MG07-026-DP

Figure S43. ¹H NMR spectrum of compound 20.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula	C ₁₁ H ₁₈ N ₂ O ₂	FW	210.2728
---------	---	----	----------



20

MG07-026-DP-C13




J:\NOTEBOOK7\MG07-026-DP-C13

Figure S44. ¹³C NMR spectrum of compound 20.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₂H₂₀N₂O₃ FW 240.2988



21

MG07-053-FRAC51-56




J:\NOTEBOOK7\MG07-053-FRAC51-56

Figure S45. ¹H NMR spectrum of compound 21.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₂H₂₀N₂O₃ FW 240.2988



21

MG07-053-C13




Figure S46. ¹³C NMR spectrum of compound 21.

+Q1: 21 MCA scans from Sample 1 (TuneSampleID) of MT20140723181728.wiff (Turbo Spray)

Max. 2.6e6 cps.





Figure S47. MS spectrum of compound 21.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₉H₂₄N₂O₄ FW 344.4049




22

MG06-046-FRAC16




J:\NOTEBOOK6\MG06-046-FRAC16

Figure S48. ¹H NMR spectrum of compound 22.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/Formula C₁₉H₂₄N₂O₄ FW 344.4049

22

MG06-046-FRA16-13C



J:\NOTEBOOK6\MG06-046-FRA16-13C

Figure S49. ¹³C NMR spectrum of compound 22.

+Q1: 15 MCA scans from Sample 1 (TuneSampleID) of MT20140418122515.wiff (Turbo Spray)

Max. 4.9e6 cps.





Figure S50. MS spectrum of compound 22.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₄H₁₆N₂O₂ FW 244.2890



23

MG05-040-CRUDE




J:\NOTEBOOK5\MG05-040-CRUDE

Figure S51. ¹H NMR spectrum of compound **23**.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₄H₁₆N₂O₂ FW 244.2890



23

MG05-040-FRAC30-36-C13




J:\NOTEBOOK5\MG05-040-FRAC30-36-C13

Figure S52. ¹³C NMR spectrum of compound 23.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₂₂H₂₃N₃O₃ FW 377.4363




24

MG05-081-FRAC6-7




J:\T-TYPE CALCIUM CHANNEL\NMR\1-24\MG05-081-FRAC6-7

Figure S53. ¹H NMR spectrum of compound 24.

This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/Formula C₂₂H₂₃N₃O₃ FW 377.4363

24

MG-05-081-FRA6-7-C13



J:\T-TYPE CALCIUM CHANNEL\NMR\1-24\MG-05-081-FRA6-7-C13

Figure S54. ¹³C NMR spectrum of compound 24.






Figure S55. MS spectrum of compound 24.

Figure S56. MS spectrum of compound **25**.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₂H₂₁N₂O FW 223.3146



27

MG06-094-CRUDE




J:\NOTEBOOK6\MG06-094-CRUDE

Figure S57. ¹H NMR spectrum of compound 27.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₂H₂₁N₃O FW 223.3146



27

MG06-094-DP-C13



J:\T-TYPE CALCIUM CHANNEL\NMR\1-27\MG06-094-DP-C13

Figure S58. ¹³C NMR spectrum of compound 27.

+Q1: 19 MCA scans from Sample 1 (TuneSampleID) of MT20141014190638.wiff (Turbo Spray)

Max. 1.2e7 cps.





Figure S59. MS spectrum of compound 27.


This report was created by ACD/NMR Processor Academic Edition. For more information go to www.acdlabs.com/nmrproc/

Formula C₁₂H₂₃N₃O FW 227.3464



28

MG06-097-H



J:\NOTEBOOK6\MG06-097-H

Figure S60. ¹H NMR spectrum of compound 28.




Figure S61. MS spectrum of compound 28.