

Supplementary Material

Electrochemically induced tandem Knoevenagel-Michael assembling of aldehydes with kojic acid: direct and efficient arylbis[3-hydroxy-6-(hydroxymethyl)-4-oxo-4H-pyran-2-yl]methanes formation

Michail N. Elinson,^{a*} Yuliya E. Ryzhkova,^a Anatoly N. Vereshchagin,^a Natalia A. Leonova,^a Alexandra P. Minaeva,^{a,b} and Mikhail P. Egorov^a

^a*N. D. Zelinsky Institute of Organic Chemistry, Leninsky prospect 47, 119991 Moscow, Russia*

Email: elinson@ioc.ac.ru



^b*D. I. Mendeleev University of Chemical Technology of Russia, Miusskaya square 9, 125047 Moscow, Russia*


Table of Contents



¹ H and ¹³ C NMR spectra of novel 2,2'-[(aryl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-ones] 2c,d,f-m with tetramethylsilane (TMS) as internal standard	S2
--	----


Chemical shifts (δ) are reported in parts per million (ppm) downfield of TMS**2,2'-[(4-Ethylphenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2c)**

© Zelinsky Institute of Organic Chemistry, Moscow; Bruker AM300 SF=300.13 MHz (1H) SI=32K SW=6002 O1=2401 PW=9.0 AQ=1.352 RD=0.00 NS=1 SR=1.69 TE=299K 13 February 2020 Opr: Daeva E.D.; Solv: DMSO-d6;




2,2'-[(4-*Tert*-butylphenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4*H*-pyran-4-one] (2d)

2,2'-[(2,5-Dimethoxyphenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2f)


2,2'-[(3-Fluorophenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2g)

2,2'-[(4-Fluorophenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2h)



© Zelinsky Institute of Organic Chemistry, Moscow; Bruker AM300 SF=300.13 MHz (1H) SI=32K SW=6002 OI=2401 PW=9.0 AQ=1.352 RD=0.00 NS=1 SR=1.69 TE=300K 28 January 2020 Opr: Daeva E.D.; Solv: DMSO-d6;



© Zelinsky Institute of Organic Chemistry, Moscow; Bruker AM300 SF=75.47 MHz (13C) SI=32K SW=18114 OI=8301 PW=14.3 AQ=0.900 RD=1.00 NS=502 SR=26.63 TE=300K 27 February 2020 Opr: Daeva E.D.; Solv: DMSO-d6;




2,2'-[(2-Chlorophenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2i)




2,2'-[(4-Chlorophenyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2j)

© Zelinsky Institute of Organic Chemistry, Moscow, Bruker AMB00 SF=300.13 MHz (1H) SI=32K SW=6002 O1=2401 PW=9.0 AQ=1.352 RD=0.00 NS=1 SR=1.69 TE=300K 28 January 2020 Opr: Daeva E.D.; Solv: DMSO-d6;




© Zelinsky Institute of Organic Chemistry, Moscow, Bruker AMB00 SF=75.47 MHz (13C) SI=32K SW=18114 O1=8301 PW=14.3 AQ=0.900 RD=1.00 NS=747 SR=26.93 TE=300K 28 February 2020 Opr: Daeva E.D.; Solv: DMSO-d6;




2,2'-[(1-Naphthyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2k).



© Zelinsky Institute of Organic Chemistry, Moscow, Bruker AM300 SF=300.13 MHz (1H) SI=32K SW=6002 OI=2401 PW=9.0 AQ=1.352 RD=0.00 NS=1 SR=1.69 TE=300K 20 March 2020 Opr. Daeva E.D.; Solv: DMSO-d6;



© Zelinsky Institute of Organic Chemistry, Moscow, Bruker AM300 SF=75.47 MHz (13C) SI=32K SW=18114 OI=8301 PW=14.3 AQ=0.901 RD=1.00 NS=563 SR=29.03 TE=299K 14 July 2020 Opr. Daeva E.D.; Solv: DMSO-d6;




2,2'-[(Pyridin-3-yl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2I).



2,2'-[(2-Furyl)methylene]bis[3-hydroxy-6-(hydroxymethyl)-4H-pyran-4-one] (2m).

© Zelinsky Institute of Organic Chemistry, Moscow, Bruker AM300 SF=300.13 MHz (1H) SI=32K SW=6002 OI=2401 PW=9.0 AQ=1.352 RD=0.00 NS=1 SR=1.69 TE=299K 10 July 2020 Opr: Daeva E.D., Solv: DMSO-d6;



© Zelinsky Institute of Organic Chemistry, Moscow, Bruker AM300 SF=75.47 MHz (13C) SI=32K SW=18114 OI=8301 PW=14.3 AQ=0.901 RD=1.00 NS=520 SR=30.55 TE=299K 16 July 2020 Opr: Daeva E.D., Solv: DMSO-d6;

