

Supplementary Material

Chemical synthesis of palmitoylated histone H4

Hironobu Hojo*^a and Isao Suetake*^b

^a *Institute for Protein Research, Osaka University, Suita, Osaka 5650871, Japan*

^b *Nakamura Gakuen Graduate School University, Fukuoka 8140198, Japan*

Email: hojo@protein.osaka-u.ac.jp, suetake@nakamura-u.ac.jp

Table of Contents

Trial synthetic routes of segment 2	S2
---	----




Fig. S1. Trial synthesis of segment **2** by the Boc method. Elution conditions: column, Cosmosil 5C18 AR-II (4.6 x 150 mm, Nacalai Tesque, Kyoto) at the flow rate of 1 mL min⁻¹; eluent, A, 0.1% TFA, B, acetonitrile containing 0.1% TFA; temperature, 50 °C; gradient, 20% B at 0 min to 40% B at 20 min linearly.




Fig. S2. Trial synthesis of the partial sequence (32-56) of segment **2** by the Fmoc-method. The introduction of Ile²⁹ to Lys³¹ was abandoned. The HPLC profile a) shows the analysis of crude peptide obtained with microwave irradiation during Fmoc group removal for Pro³² to Lys⁴⁴. Profile b) shows the analysis of crude peptide obtained without microwave irradiation during Fmoc group removal for Pro³² to Lys⁴⁴. Elution conditions: column, Cosmosil 5C18 AR-II (4.6 x 150 mm, Nacalai Tesque, Kyoto) at the flow rate of 1 mL min⁻¹; eluent, A, 0.1% TFA, B, acetonitrile containing 0.1% TFA; temperature, 50 °C; gradient, 20% B at 0 min to 60% B at 20 min linearly.