## **Supplementary Material**

## Unusual ionization of phosphine-boranes under RP-HPLC-HRMS conditions disclose a potential system for reduction of C=O bond

Sylwia Sowa,<sup>a</sup> Ewelina Buczak,<sup>b</sup> Paweł Woźnicki,<sup>a</sup> Olga Bąk,<sup>c</sup> Piotr Borowski,<sup>\*d</sup> Beata Herbaczyńska-Stankevič,<sup>a</sup> and Marek Stankevič<sup>\*a</sup>

 <sup>a</sup>Department of Organic Chemistry, Institute of Chemical Sciences, Faculty of Chemistry, Maria Curie-Skłodowska University in Lublin, 33 Gliniana st., 20-614 Lublin, Poland
<sup>b</sup>Minisynthesis Team, Łukasiewicz Research Network – Industrial Chemistry Institute, 8 Rydygiera st.,

01-793 Warsaw, Poland

<sup>c</sup>Department of Chromatographic Methods, Institute of Chemical Sciences, Faculty of Chemistry, Maria Curie-Skłodowska University in Lublin, 3 Marie Curie-Skłodowska sq., 20-031 Lublin, Poland

<sup>d</sup>Department of Theoretical Chemistry, Institute of Chemical Sciences, Faculty of Chemistry, Maria Curie-Skłodowska University in Lublin, 3 Marie Curie-Skłodowska sq., 20-031 Lublin, Poland

Email: marek.stankevic@mail.umcs.pl piotr.borowski@mail.umcs.pl

## **Table of Contents**

S1. <sup>1</sup> H NMR spectrum of (3–hydroxypropyl)methylphenylphosphine–borane <b>20</b> S3
S2. <sup>13</sup> C NMR spectrum of (3–hydroxypropyl)methylphenylphosphine–borane <b>20</b> S4
S3. <sup>31</sup> P NMR spectrum of (3–hydroxypropyl)methylphenylphosphine–borane <b>20</b> S5
S4. <sup>1</sup> H NMR spectrum of ((3 <i>R</i> )–3–hydroxybutyl)methylphenylphosphine–borane (mixture of
diastereomers) 21
S5. <sup>13</sup> C NMR spectrum of ((3 <i>R</i> )–3–hydroxybutyl)methylphenylphosphine–borane (mixture of
diastereomers) <b>21</b>
S6. <sup>31</sup> P NMR spectrum of ((3 <i>R</i> )–3–hydroxybutyl)methylphenylphosphine–borane (mixture of
diastereomers) 21
S7. <sup>1</sup> H NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanone <b>27</b> S9
S8. <sup>13</sup> C NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanone <b>27</b> S10
S9. <sup>31</sup> P NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanone <b>27</b> S11
S10. <sup>1</sup> H NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanol <b>28</b> S12
S11. <sup>13</sup> C NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanol <b>28</b> S13
S12. <sup>31</sup> P NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanol <b>28</b> S14
S13. <sup>1</sup> H NMR spectrum of ethyl ( <i>t</i> -butylphenylboranatophosphinyl)acetate <b>29</b>
S14. <sup>13</sup> C NMR spectrum of ethyl ( <i>t</i> -butylphenylboranatophosphinyl)acetate <b>29</b>
S15. <sup>31</sup> P NMR spectrum of ethyl ( <i>t</i> -butylphenylboranatophosphinyl)acetate <b>29</b> S17

S16. <sup>1</sup> H NMR spectrum of tri( <i>p</i> -anisyl)phosphine-borane <b>30</b>	
S17. <sup>13</sup> C NMR spectrum of tri( <i>p</i> -anisyl)phosphine-borane <b>30</b>	
S18. <sup>31</sup> P NMR spectrum of tri( <i>p</i> -anisyl)phosphine-borane <b>30</b>	



S1. <sup>1</sup>H NMR spectrum of (3–hydroxypropyl)methylphenylphosphine–borane 20 (500 MHz, CDCl<sub>3</sub>)



S2. <sup>13</sup>C NMR spectrum of (3–hydroxypropyl)methylphenylphosphine–borane 20 (126 MHz, CDCl<sub>3</sub>)





**S3.** <sup>31</sup>P NMR spectrum of (3–hydroxypropyl)methylphenylphosphine–borane **20** (202 MHz, CDCl<sub>3</sub>)

## Issue in honor of Professor Gyorgy Keglevich



**S4.** <sup>1</sup>H NMR spectrum of ((3*R*)–3–hydroxybutyl)methylphenylphosphine–borane (mixture of diastereomers) **21** (500 MHz, CDCl<sub>3</sub>).

©AUTHOR(S)



**S5.** <sup>13</sup>C NMR spectrum of ((3*R*)–3–hydroxybutyl)methylphenylphosphine–borane (mixture of diastereomers) **21** (126 MHz, CDCl<sub>3</sub>)

DN-276-1.003.esp

BH₃ ∮ ₽ ,OH Y} ) Any many prover the provide and the provide a state of the second and t When Mine 75 70 65 60 55 50 45 40 35 30 25 20 15 10 5 0 Chemical Shift (ppm) 95 90 85 80

S6. <sup>31</sup>P NMR spectrum of ((3*R*)–3–hydroxybutyl)methylphenylphosphine–borane (mixture of diastereomers) 21 (202 MHz, CDCl<sub>3</sub>)



S7. <sup>1</sup>H NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanone 27 (500 MHz, CDCl<sub>3</sub>)

BHE-0002F6.003.esp



**S8.** <sup>13</sup>C NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanone **27** (126 MHz, CDCl<sub>3</sub>)

BHE-0002F6.002.esp



**S9.** <sup>31</sup>P NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanone **27** (202 MHz, CDCl<sub>3</sub>)



**S10.** <sup>1</sup>H NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanol **28** (500 MHz, CDCl<sub>3</sub>)

BHE-0236F2SUCHY.003.esp



**S11.** <sup>13</sup>C NMR spectrum of 2–(diphenylboranatophosphinyl)–1–indanol **28** (126 MHz, CDCl<sub>3</sub>)





**S13.** <sup>1</sup>H NMR spectrum of ethyl (*t*-butylphenylboranatophosphinyl)acetate **29** (500 MHz, CDCl<sub>3</sub>)

BHE-X315-22-F1.003.esp



**S14.** <sup>13</sup>C NMR spectrum of ethyl (*t*-butylphenylboranatophosphinyl)acetate **29** (126 MHz, CDCl<sub>3</sub>)



**S15.** <sup>31</sup>P NMR spectrum of ethyl (*t*–butylphenylboranatophosphinyl)acetate **29** (202 MHz, CDCl<sub>3</sub>)



**S16.** <sup>1</sup>H NMR spectrum of tri(*p*-anisyl)phosphine-borane **30** (500 MHz, CDCl<sub>3</sub>)



**S17.** <sup>13</sup>C NMR spectrum of tri(*p*-anisyl)phosphine-borane **30** (126 MHz, CDCl<sub>3</sub>)

