Supplementary Material

Formal allene insertion into amides. Reaction of propargyl magnesium bromide with morpholine amides

Amir M. Al Mufti,[a,b] Marina V. Molchanova,[a,c] Ivan N. Myasnyanko,[a,d] Mikhail S. Baranov,[a,d] Viktoria A. Ikonnikova,[a,b] Andrey A. Mikhaylov[a,*]

[a] *Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry of the Russian Academy of Sciences, 16/10 Miklukho-Maklaya St., Moscow, 117997, Russia*
[b] *D.I. Mendeleev University of Chemical Technology of Russia, 9 Miusskaya sq., 125047 Moscow, Russia*
[c] *Lomonosov Moscow State University, Leninskie Gory 1 (3), Moscow, 119991, Russia*
[d] *Pirogov Russian National Research Medical University, 1 Ostrovityanova St., Moscow 117997, Russia*
Email: mikhaylov_andrey@yahoo.com

Table of Contents

General experimentation

All reactions were performed in round-bottom flasks fitted with rubber septa. Reactions sensitive to air and/or moisture were performed under a positive pressure of argon. Air- and moisture-sensitive liquids were transferred by syringe. Analytical thin-layer chromatography (TLC) was performed using aluminum plates pre-coated with silica gel (silica gel 60 F_{254}, Sorbfil). TLC plates were visualized by exposure to 254 nm ultraviolet light (UV) or were stained by submersion in acidic ethanolic solution of vanillin followed by brief heating (vanillin) or submersion in aqueous potassium permanganate solution followed by extensive washing with water ($KMnO_4$). Flash-column chromatography was carried out on silica gel (60 Å, 230–400 mesh, Merck). All solvents for chromatography and extractions were technical grade and distilled prior use.

All reagents were obtained from commercial suppliers and were used without further purification. Et_2O and THF were stored over sodium benzophenone ketyl and were distilled directly prior use.

Nuclear magnetic resonance spectra were recorded using Bruker Furier 300, Bruker Avance 800 instruments at indicated temperature. Data are represented as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet and/or multiple resonances), coupling constant (*J*) in Hertz, integration. Proton chemical shifts are expressed in parts per million (ppm, δ scale) and are referenced to residual protium in the NMR solvents ($CHCl_3$, δ 7.26 ppm). Carbon chemical shifts are expressed in parts per million (ppm, δ scale) and are referenced to the carbon resonances of the NMR solvents ($CDCl_3$, δ 77.16 ppm).

High-resolution mass spectra were recorded on a Bruker micrOTOF-Q II mass spectrometer using electrospray ionization (ESI–TOF). Melting points were determined on Kofler melting point apparatus and are uncorrected.

Copies of NMR spectra



3a

^{1}H NMR (300 MHz, $CDCl_3$)





3a

^{13}C NMR (75 MHz, $CDCl_3$)





4a

92:8 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)





4a

^{13}C NMR (75 MHz, $CDCl_3$)





3b

^{1}H NMR (300 MHz, $CDCl_3$)





3b

^{13}C NMR (75 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



4b

95:5 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)

ARKIVOC 2024, *ii*, S1-S58



4b

^{13}C NMR (75 MHz, $CDCl_3$)





3c

^{1}H NMR (300 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



3c

^{13}C NMR (75 MHz, $CDCl_3$)





4c

95:5 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



4c

^{13}C NMR (75 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



3d

^{1}H NMR (300 MHz, $CDCl_3$)





3d

^{13}C NMR (75 MHz, $CDCl_3$)

187.93
166.07
163.10
162.75
138.76
138.72
129.75
129.63
115.15
114.87
94.13
66.37
46.48
16.18

260 250 240 230 220 210 200 190 180 170 160 150 140 130 120 110 100 90 80 70 60 50 40 30 20 10 0 -10 -20 -30 -40 -50 -6

ARKIVOC 2024, *ii*, S1-S58



4d

93:7 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4d

^{13}C NMR (75 MHz, $CDCl_3$)





4e

90:10 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



4e

^{13}C NMR (75 MHz, $CDCl_3$)





3f

^{1}H NMR (300 MHz, $CDCl_3$)



3f

^{13}C NMR (75 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



4f

88:12 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4f

^{13}C NMR (75 MHz, $CDCl_3$)





4g

92:8 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)





4g

^{13}C NMR (75 MHz, $CDCl_3$)





4i

80:20 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4i

80:20 enol/keto forms

^{13}C NMR (75 MHz, $CDCl_3$)

202.70
194.83
194.52
181.35
158.97
158.37
134.71
132.96
130.86
130.16
124.10
120.93
120.71
111.72
111.59
101.87
58.93
55.63
55.42
30.34
26.14





3j

^{1}H NMR (300 MHz, $CDCl_3$)



3j

^{13}C NMR (75 MHz, $CDCl_3$)





3k

^{1}H NMR (300 MHz, $CDCl_3$)





3k

^{13}C NMR (75 MHz, $CDCl_3$)





4k

^{1}H NMR (800 MHz, $CDCl_3$)





4k

^{13}C NMR (201 MHz, $CDCl_3$)





4l

87:13 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4l

87:13 enol/keto forms

^{13}C NMR (75 MHz, $CDCl_3$)





3m

^{1}H NMR (800 MHz, $CDCl_3$)



3m

^{13}C NMR (201 MHz, $CDCl_3$)





4m

97:3 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4m

^{13}C NMR (75 MHz, $CDCl_3$)



3n

(ca. 90% purity)

^{1}H NMR (300 MHz, $CDCl_3$)



ARKIVOC 2024, *ii*, S1-S58



3n

(ca. 90% purity)

^{13}C NMR (75 MHz, CDCl$_3$)





4n

84:16 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)





4n

84:16 enol/keto forms

^{13}C NMR (75 MHz, $CDCl_3$)





3o

(ca. 85% purity)

^{1}H NMR (300 MHz, $CDCl_3$)





3o

(ca. 85% purity)

^{13}C NMR (75 MHz, $CDCl_3$)





4o

88:12 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4o

88:12 enol/keto forms

^{13}C NMR (75 MHz, $CDCl_3$)



4p

90:10 enol/keto forms

^{1}H NMR (300 MHz, $CDCl_3$)



4p

90:10 enol/keto forms

^{13}C NMR (75 MHz, $CDCl_3$)



3r

^{1}H NMR (300 MHz, $CDCl_3$)





3r

^{13}C NMR (75 MHz, $CDCl_3$)





5

^{1}H NMR (800 MHz, $CDCl_3$)



^{13}C NMR (201 MHz, $CDCl_3$)





6

^{1}H NMR (300 MHz, $CDCl_3$)





6

^{13}C NMR (75 MHz, $CDCl_3$)



7

^{1}H NMR (800 MHz, $CDCl_3$)



7

^{13}C NMR (201 MHz, $CDCl_3$)