

Friedel–Crafts Chemistry. Part 63. Syntheses of some *condensed N-heterocyclic* systems via combined Darzens and Friedel-Crafts approaches

Hassan A. K. Abd El-Aal* and Ali A. Khalaf

Chemistry Department, Faculty of Science, Assiut University, Assiut, 71516, Egypt

E-mails: hassankotb33@yahoo.com; hassankotb@aun.edu.eg


Received 07-13-2024

Accepted 08-23-2024

Published on line 08-30-2024

Abstract

An efficient protocol for the synthesis of novel fused *N*-heteropolycycles **10a–f** (tetracyclic-fused and bridged dibenzo-azepinone, -azocinones, -azoninone, benzo[*b*]pyrido[3,2-*f*]azocinone and benzo[*c*]pyrido[3,2-*g*]azoninones) via combined Darzens and Friedel-Crafts methodologies is described. The starting ketones **5a,b** were conveniently obtained via cyanoethylation of aminoxylens **1a,b** followed by hydrolysis of the resulting nitriles **3a,b** to the corresponding carboxylic acids **4a,b** and were finally cycloacylated by intramolecular Friedel-Crafts reaction. The key acetylated intermediates **7a,b** were obtained by Darzens condensation. Subsequently, two steps led to substituted heterocyclic acids **9a–f** which successfully underwent Friedel–Crafts cycloacylations mediated by $\text{AlCl}_3/\text{CH}_3\text{NO}_2$, TfOH or P_2O_5 catalysts. The method offers readily access to diverse drug-like **10a–f** scaffolds in moderate to good yields.



Keywords: Friedel-Crafts cyclization, Darzens reaction, medium-sized *N*-heterocycles, azocines, azonines, Brønsted and Lewis acid catalysts

Introduction

Polycyclic systems containing medium-sized *N*-heterocycles (e.g. benzofused and bridged-azepines and higher ring systems) are prominent scaffolds in numerous biologically active natural products^{1–3} and pharmaceutically relevant compounds,^{4–6} possessing a wide diversity of biological activities (Figure1).^{7–9} Given their structural complexity as well as their intriguing pharmacological activities, syntheses of these scaffolds have been subject to many synthetic endeavors.




Figure 1. Some of biologically active natural alkaloids and examples of condensed heterocycles prepared by Friedel–Crafts processes.

In literature, numerous comprehensive reviews surveying their occurrence, synthetic strategies and biological activities have been published.^{10,11} But up to our knowledge, even with these awesome accomplishments, there is a lack of efficient and generalized protocols for the synthesis of eight- and higher membered azaheterocycles. Only some examples of dopamine receptor antagonists have been isolated from several indole-alkaloid families, such as, Erythrina, Aspidosperma, Hasubanan and Iboga alkaloids.¹² The scarcity associated with the syntheses of condensed polycycles containing nine- and higher-membered amine containing ring systems can mainly be attributed to synthetic difficulties associated with the presence of a large number of conformations, low energetic transannular reactions¹³ as well as with unfavorable entropy barriers.¹⁴

It is worthy of mention here that some of the successfully applied approaches to synthesize condensed azocine systems included: cycloaddition reactions between enamines and ethyl propiolate followed by electrocyclic ring-opening reaction,¹⁵ cascade reactions of 1,2,4-triazines with enamines,¹⁶ reductive ring-expansion reactions of aromatic cyclic ketoximes with di-isobutylaluminumhydride,¹⁷ intramolecular condensation of ethyl *N*-substituted-cyanomethylantranilates using potassium *tert*-butoxide,¹⁸ thermal isomerization reactions based on the *tert*-amino effect of *o,o'*-functionalized biaryls originating from Suzuki reaction followed by the Knoevenagel condensation,¹⁹ microwave-assisted Suzuki-Miyaura cross-coupling followed by ring-closing metathesis of various acyclic and alicyclic aminobiaryl intermediates,²⁰ Mo-alkylidene complexes catalyzed ring closing metathesis of appropriate diene-systems,²¹ aza-Claisen rearrangement of

substituted allylamines followed by intramolecular Heck reaction,²² ring closing metathesis using Grubbs-catalyst derivatives of diene, enyne and amide intermediates formed by a reductive amination of 2-vinylindole carbaldehydes.²³ As a result, these pioneering syntheses proceeds from one of the following limitations: (i) limitations in substrate scope, (ii) utilizing expensive catalysts, (iii) low functional tolerance, (iv) syntheses proceeds through lengthy sequences with low overall yields.


In our previous work in this series, we developed new synthetic routes for the construction of a number of fused carbo- and heteropolycycles containing naphthalene, thiophene, quinoline and indole moieties via Friedel-Crafts cycloacylations of suitably synthesized precursors.²⁴⁻²⁷ Herein, we extend our efforts applying some parallel facile protocols of broad applicability for the construction of a wide range of polyfunctionalized heteropolycycles including; dibenzo-azepinones, -azocinones, -azoninones, benzo[*b*]pyrido[3,2-*f*]azocinones and benzo[*c*]pyrido[3,2-*g*]azoninones, via Lewis and Brønsted acid-mediated Friedel-Crafts cycloacylations of easily prepared precursors.

Results and Discussion

Synthetically, the desired key starting bicyclic-heterocarboxylic acids **9a-f** were readily generated starting with the synthesis of propanenitriles **3a,b** (Scheme 1). These propionitriles were readily obtained in good yields via cyanoethylation²⁸ of the correspondingly aromatic amines **1a,b** with acrylonitrile (**2**) in the presence of BnMe₃NOH (Triton B) under standard conditions. Basic hydrolysis of nitriles **3a,b** to the corresponding carboxylic acids **4a,b** using KOH in diethylene glycol, was followed by cyclization by H₂SO₄ to furnish quinolinone **5a** and benzo[*c*]azepinone **5b** in good yields. Treatment of these ketones with ethyl 2-bromopropionate in the presence of EtONa in ether following Darzens²⁹ condensation yielded glycidic ester **6a,b**.

Hydrolysis of the latter esters were achieved by EtONa in ethanol, followed by decarboxylations of the resulting salts by heating with glacial acetic acid. This was accompanied by a vigorous CO₂ evolution and led to the acetyl derivatives **7a,b** in good yields. The key intermediate acids **8a,b** were obtained in 82–87% yields by the oxidative demethylation (Haloform) reactions of the resulting methyl ketones **7a,b** in the presence of iodine and KOH in dioxane. The requisite highly electrophilic precursors *N*-arylcarboxylic acids **9a-f** were readily prepared in moderate yield via Ullmann³⁰ coupling reactions of acids **8a,b** with various aromatic halides in presence of K₂CO₃ in DMF.

The versatility of this approach was demonstrated by the attempted cycloacylations of precursors **9a-f** to novel tetracyclic ketones **10a-f** via extensive optimization of the cyclization conditions. Therefore, several factors were considered as we further optimized the conditions to improve the effectiveness of the transformation such as the screening of more suitable acidic promoters, variations of the reaction temperature, the type of solvent and the length of the reaction time. The chosen conditions and their results are summarized in Scheme 1 and Table 1. Ring closure progress of substrates **9a-f** has been performed by using AlCl₃/CH₃NO₂ or TfOH (trifluoromethanesulfonic acid) or P₂O₅ catalysts as promoting Friedel-Crafts cycloacylation reactions. The structure elucidations of all new compounds were carried out by spectral and elemental analysis.















Scheme 1. Reagents and conditions: (i) Acrylonitrile (**2**)/BnMe₃NOH (Triton B), dioxane, 80-90 °C, 5 h, (ii) KOH, DEG, 15h,Reflux, HCl, (iii) H₂SO₄ (85%), 90-100 °C, 6 h, (iv) CH₃CHBrCO₂Et/EtONa/ether, 0-5°C, 9 h, (v) *a*.EtONa/EtOH, 0-5°C, 5h, *b*. glacial AcOH, 70-80 °C, 1h, NaHCO₃, (vi) I₂/KOH, Dioxane, 60-70 °C, 2 h, AcOH, (vii) Aryl halide (PhBr or PhCH₂Cl or 2-Picolyl chloride), K₂CO₃/Pyridine/CuI, DMF, 110-20 °C, 10h, (viii) Cycloacylations of acids precursors mediated by AlCl₃/CH₃NO₂ or TfOH or P₂O₅ promoters (Table 1).

The striking observations from initial examination of cyclization conditions suitable for hetero-acids **9a-f** led to the conclusion that several potentials had to be addressed for this type of cyclization reactions. In such cases, Friedel-Crafts ring closure is complicated by the continued presence of the poor leaving -OH group ability, as well as by the electrophilic inhibitions³¹ of acidic promoters through their binding properties on the basic nitrogen (sp³-hybridized) present in substrate. In an attempt to circumvent this issue, it was speculated from the classical valence bond theory³² that the rate of the cyclization is strongly dependent on the catalyst strength, while the ease of the cyclization depends mainly on the binding activity of the acidic catalysts on the heteroatom present in the electron-rich substrate.

Presumably, the stronger the catalyst, the more it coordinates with the electron density of the heteroatom, leading ultimately to deactivation of its nucleophilicity and consequently, retardation of the cyclization process. Thus, mild strength catalysts favor the cyclization process in such cases. In fact, numerous literature studies on the mechanism of Friedel-Crafts acylations^{33,34} have been reported and included two distinct mechanistic pathways for electrophilic aromatic substitutions depending on the nature of the acylating agent and the strength of the acidic promoters. For example, the suggested mechanistic rationalization for the formation of tetracyclic ketone **10b** from sufficiently electron rich precursor **9b** is presented in Scheme 2.

Table 1. Condition screening for intramolecular acylation of heterocyclic acids **9a–f**


Entry	Substrate	Cyclization conditions	Time, h	Product	Yield (%)
1		Method I*	10		83
		Method II**	6		75
		Method III***	12		70
2		Method I	11		86
		Method II	8		77
		Method III	15		80
3		Method I	18		82
		Method II	8		84
		Method III	16		76
4		Method I	14		78
		Method II	8		70
		Method III	18		71
5		Method I			79
		Method II	8		83
		Method III	8		75
6		Method I	20		86
		Method II	10		72
		Method III	10		70

* Method I: acids **9a–f** (4 mmol) in CH₂Cl₂ (10 ml), AlCl₃ (10 mmol), MeNO₂ (80 mmol), room temperature.

** Method II: acids **9a–f** (4 mmol), TfOH (1.6 ml, 20 mmol), 1,2-DCE (15 ml), 60–70 °C.

*** Method III: acids **9a–f** (4 mmol), P₂O₅ (5 g), toluene (15 mL), reflux.

From a mechanistic point of view, acyl-cations, either a resonance stabilized σ -complex, or Wheland intermediate are generated by the treatment of heterocyclic acid **9b** with acidic promoters (Scheme 2). The catalyst-substrate σ -complex is formed using a Lewis acid, while Wheland intermediates form by the treatment of electron rich carboxylic acid with Brønsted acid, respectively. Subsequently, removal of H₂O molecules generates a free resonating acyl-carbocation σ -complex or an ion pair.³⁵ Electrophilic substitution reactions are taking place by loss of a ring proton to produce ketone **10b**.



Scheme 2. Proposed mechanisms for cyclizations of heterocyclic acid **9b**.

It has been suggested³⁶ that a strongly electrophilic Lewis or Brønsted acid provide transition states resembling a π -complex, while the less electrophilic catalysts provide transition states similar to a stabilized σ -complex. Increasingly, the electrophilicity of this complex is determined by a combination of the nature of the acylating agent and the strength of the acidic catalyst. However, the regiochemistry is determined by the nature of the transition state leading to the intermediate carbocation. These mechanistic investigations suggested that poor acid catalysts are unable to coordinate effectively with the heteroatoms. It seemed therefore that the low reactivities of the three selected catalysts we used are due to catalyst inhibition problems. Wherefore, more than stoichiometric or excess amounts of the oxophilic promoter necessary for the Friedel-Crafts cycloacylations of electron-rich arenes. Examination of the conditions suitable for the Friedel-Crafts cycloacylations proved more fruitful.

Concurrent with these studies, the literature precedents since the late 1930s showed a great effort, however exclusively on the use of the Friedel-Crafts processes for the construction of complex carbo- and heterocyclic scaffolds from highly reactive electron-deficient and electron-rich arenes (Figure 1).^{10,37-41}

The formation of cyclic products **10a-f** was proven based on the ¹H NMR, ¹³C NMR and IR spectroscopic techniques. A closer look at the acetyl-intermediates **7a,b** showed that such molecules are emphasizing inequivalent protons due to the presence of chiral center and cyclic diastereotopic protons.

For all of the acyclic precursors **9a-f** and the corresponding tetracyclic products **10a-f** involved, the generation of a stereogenic centre led to the fact that the adjacent methylene group protons become diastereotopic⁴² in nature. In addition to the chirality of the carbon and nitrogen atoms, these rings have a large number of conformations⁴³ which collaborates with the non-bonded interactions between atoms,⁴⁴ the considerable amount of Baeyer, Pitzer and Prelog strains,⁴⁵ transannular reactions in medium-rings conformers⁴⁶ as well as the shielding effect exerted by the magnetic anisotropic effect⁴⁷ generated by heteroatoms. These effects lead to protons of the methylene group adjacent to the stereogenic center or bonded to the heteroatom, become inequivalent and are present as unresolved signals in their NMR spectrums (Figure 3).




Figure 3. Diastereotopic protons containing tetracyclic structure **10b** and its acid precursor **9b**.

For example, a comparison of the spectroscopic data of heterocyclic acid **9b** and tetracyclic ketone **10b** does clearly show the disappearance of the characteristic COOH-stretch, found at 2610 cm^{-1} in the IR spectrum of acid **9b** and the appearance of the characteristic CO-stretch at 1700 cm^{-1} in the IR spectrum of ketone **10b**.

The inspection of the ^1H NMR spectrum of acid **9b** and ketone **10b** displays unresolved signals unlike the other multiplets of the spectrum. Thus, the ^1H NMR of the carboxylic acid **9b** displayed several distinct signals in which the aromatic protons appeared in the range of δ 6.43–7.27 ppm, two methyl groups appeared as singlets near δ 2.12 and 2.27 ppm, benzylic- CH_2 showed as characteristic singlet near δ 4.32 ppm and the diagnostic COOH signal at δ 10.85 ppm.

As expected, the remaining three characterized signals included two bridged diastereotopic CH_2 groups and the chiral CH proton. These appeared as a complex set of overlapped signals with different coupling constants. The upfield diastereotopic protons of C^3H_2 -group (pseudo-axial & -equatorial; H_a & H_b) appeared near δ 2.09 ppm with multiplicities of doublet of doublet of doublet of doublets with different coupling constants. From the other hand, the C^4H proton appeared as doublet of doublets near δ 4.07 ppm with coupling constants, 9.8 and 4.4 Hz. The third downfield signal for $\text{N}-\text{C}^2\text{H}_2$ protons appeared at δ 3.24 ppm with multiplicities of doublet of doublet of doublets and three coupling constants of 13.7, 6.9 and 1.8 Hz.

In comparison, the ^1H NMR of azocinone skeleton **10b** showed aromatic proton signals in the range of δ 6.43–7.86 ppm, two methyl groups appeared as singlets at δ 2.10 and 2.26 ppm, cyclic $\text{N}-\text{C}^6\text{H}_2$ protons appeared as doublet near δ 4.46 ppm with coupling constants of 13.8 Hz. The upfield C^{13}H_2 -protons appeared near δ 1.98 ppm with multiplicities of doublet of doublet of doublet of doublets with coupling constants 13.3, 7.9, 3.5 and 3.3 Hz. The chiral- C^{12}H proton appeared as doublet of doublets near δ 4.19 ppm with coupling constants 5.1 and 1.5 Hz.

However, the downfield $\text{N}-\text{C}^{14}\text{H}_2$ protons appeared at δ 3.44 ppm as doublet of doublet of doublets with coupling constants of 14.4, 7.9 and 3.5 Hz. Also, the ^1H NMR spectrum of ketone **10b** indicates the absence of the diagnostic carboxylic acid signal shown at δ 10.85 ppm. Further, changes in the ^{13}C NMR spectra proved that the cyclization process had occurred with the disappearance of the COOH-peak at 176 ppm in the ^{13}C NMR spectrum of acid **9b** and the appearance of a new carbonyl group at 197 ppm in the ^{13}C NMR spectrum of ketone **10b**.

Conclusions

In summary, we have developed a novel easy and concise catalytic protocol for the synthesis of tetracyclic heteroaromatic scaffolds **10a–f**, in particular, dibenzo-azepinone, -azocinones, azoninone, benzo[*b*]pyrido[3,2-*f*]azocinone and benzo[*c*]pyrido[3,2-*g*]azoninones via combined Darzens condensations and Friedel-Crafts

cycloacylations starting from simple readily available acyclic precursors. The operational simplicity and ready availability of all precursors should make this methodology a useful tool for the assembly of the structurally diverse fused *N*-heterocycles in very good overall yields.

Experimental Section

General. All chemicals used were of reagent grade and solvents were freshly distilled and dried by standard procedures before use. Melting points were taken on a digital Gallenkamp capillary melting point apparatus and are uncorrected. Infrared (IR) spectra were obtained on a Perkin-Elmer 1600 FT-IR spectrophotometer using KBr wafer and thin film techniques (ν cm^{-1}) and are in cm^{-1} ; The NMR spectra were recorded on JEOL LA 400 MHz FT-NMR (400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR) using CDCl_3 solvent with tetramethylsilane (Me_4Si , TMS) as internal standard. Chemical shifts (δ) and *J* values are reported in ppm and Hz, respectively. Elemental analyses were carried out by a GmbH Vario EL III, 2400, CHNOS-elemental analyzer. The progress of reactions were monitored by thin-layer chromatography (TLC) analysis on precoated silica plates (Silufol, UV-254 TLC, aluminum sheets) and plates were visualized with UV light (at 254 and/or 360 nm). Flash column chromatography, where necessary, was performed on silica gel (230–400 mesh).

General procedure for the synthesis of substituted aryl propanenitriles 3a,b. To an ice-cold solution of aromatic amines **1a** or **1b** (35.00 mmol) and acrylonitrile **2** (3.4 g, 42.00 mmol) in dioxane (25 mL) was added triton B (1.5 ml). The reaction mixture was heated in a water bath at 80–90 °C for 5 h. The mixture was then cooled and concentrated in *vacuo*. The resultant pasty was triturated with methanol (3×5 mL) and the resultant precipitate was filtered, washed with methanol and finally dried to give the crude product **3a,b**. Crystal appearance, yields, and spectral data of acids **3a,b** are given:

3-(2,5-Dimethylphenylamino)propanenitrile (3a). White plates; 79%; mp 59–61 °C (acetone); IR (KBr, ν , cm^{-1}): 3420 (N-H), 3045 (Ar-H), 2975 (C-H), 2250 (CN), 1600, 1585, 1480, 1445, 1375, 1245, 775. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.08 (3H, s, CH_3), 2.26 (3H, s, CH_3), 2.95 (2H, t, *J* 6.7 Hz, $\text{NH-C}^3\text{H}_2$), 3.23 (2H, t, *J* 6.7 Hz, $\text{-C}^2\text{H}_2$), 6.32 (1H, dd, *J* 1.2, 0.5 Hz), 6.64–6.84 (2H, dd, *J* 7.8, 1.2 Hz), 9.84 (s, 1H, NH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 17.7 (1C, -CH_3), 18.6 (1C, -CH_3), 21.3 (1C, $\text{C}^\alpha\text{H}_2$, C-2), 39.6 (1C, C^βH_2 , C-3), 114.4 (1C, Ar.,C-6), 118.0 (1C, -CN), 129.1 (1C, Ar.,C-4), 129.9 (1C, Ar.,C-3), 130.3 (1C, Ar.,C-2), 133.7 (1C, Ar.,C-5), 140.2 (1C, Ar.,C-1). Anal. Calcd. for $\text{C}_{11}\text{H}_{14}\text{N}_2$ (174); C, 75.86; H, 8.04; N, 16.09. Found; C, 75.77; H, 8.14; N, 16.15%.

3-(2,5-Dimethylbenzylamino)propanenitrile (3b). Pale yellow plates; 82%, mp 84–86 °C (benzene); IR (KBr, ν , cm^{-1}): 3490 (N-H), 3020 (Ar-H), 2990 (C-H), 2235 (CN), 1600, 1590, 1480, 1440, 1370, 1250, 774. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.12 (3H, s, CH_3), 2.32 (3H, s, CH_3), 2.83 (2H, t, *J* 6.8 Hz, $\text{NH-C}^3\text{H}_2$), 2.89 (2H, t, *J* 6.8 Hz, $\text{-C}^2\text{H}_2$), 3.67 (2H, s, PhCH_2), 6.77–6.99 (2H, dd, *J* 8.2, 2.5 Hz), 7.20 (1H, dd, *J* 2.5, 0.5 Hz), 9.92 (s, 1H, NH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 18.6 (1C, -CH_3), 20.0 (1C, -CH_3), 21.3 (1C, $\text{C}^\alpha\text{H}_2$, C-2), 44.2 (1C, $\text{C}^\delta\text{H}_2$, C-3), 46.7 (1C, C^βH_2 , C-3), 118.0 (1C, Ar.,C-6), 127.4 (1C, -CN), 128.9 (1C, Ar.,C-4), 129.1 (1C, Ar.,C-3), 129.8 (1C, Ar.,C-2), 134.8 (1C, Ar.,C-5), 135.3 (1C, Ar.,C-1). Anal. Calcd. for $\text{C}_{12}\text{H}_{16}\text{N}_2$ (188); C, 76.59; H, 8.51; N, 14.89. Found; C, 76.50; H, 8.60; N, 14.84 %.

General procedure for the synthesis of substituted bicyclic ketones 5a,b. These intermediate compounds were obtained in two reaction steps starting with propanenitriles **3a,b**. A summary of the steps is given in the following:

(i) To a solution of nitrile **3a** or **3b** (15.00 mmol) in diethylene glycol (30 mL) was added a solution of potassium hydroxide (5.6 g, 100.00 mmol) in H_2O (6 mL) and the resulting mixture was refluxed for 15 h. The reaction

mixture was then cooled and poured into ice-cold water (200 ml). The mixture was neutralized to pH 6-7 using HCl solution (5%). After standing for overnight, the resulted solid was collected, washed and dried to give crude acids **4a,b**. Further purification and yields are given in the following.

3-((2,5-dimethylphenyl)amino)propanoic acid (4a). Pale yellow crystals; 74 %; mp 141-43 °C (ethanol); IR (KBr, ν , cm^{-1}): 3380 (N-H), 3050 (Ar-H), 2975 (C-H), 2560 (OH), 1715 (C=O), 1600, 1590, 1480, 1440, 1380, 1255, 1120, 784. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.08 (3H, s, CH_3), 2.26 (3H, s, CH_3), 2.50 (2H, t, J 6.8 Hz, $\text{NH}-\text{C}^3\text{H}_2$), 3.22 (2H, t, J 6.8 Hz, $-\text{C}^2\text{H}_2$), 4.45 (1H, s, NH), 6.32 (1H, dd, J 1.2, 0.5 Hz), 6.64-6.84 (2H, dd, J 7.8, 0.5 Hz), 9.97 (s, 1H, COOH). Anal. Calcd. for $\text{C}_{11}\text{H}_{15}\text{NO}_2$ (193); C, 68.39; H, 7.77; N, 7.25. Found; C, 68.27; H, 7.90; N, 7.31%.

3-((2,5-dimethylbenzyl)amino)propanoic acid (4b). Cream crystals; 70 %; mp 147-49 °C (ethanol); IR (KBr, ν , cm^{-1}): 3420 (N-H), 3010 (Ar-H), 2988 (C-H), 2525 (OH), 1718 (C=O), 1600, 1590, 1480, 1440, 1365, 1230, 1128, 790. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.18 (6H, s, 2CH_3), 2.27 (3H, s, CH_3), 2.48 (2H, t, J 6.8 Hz, $\text{NH}-\text{C}^3\text{H}_2$), 2.85 (2H, t, J 6.8 Hz, $-\text{C}^2\text{H}_2$), 3.66 (2H, s, C^5H_2), 4.38 (1H, s, NH), 6.32 (1H, dd, J 1.2, 0.5 Hz), 6.60-6.84 (2H, dd, J 7.8, 0.5 Hz), 10.52 (s, 1H, COOH). Anal. Calcd. for $\text{C}_{12}\text{H}_{17}\text{NO}_2$ (207); C, 69.56; H, 8.21; N, 6.76. Found; C, 69.50; H, 8.34; N, 6.80%.

(ii) A mixture of carboxylic acid **4a** or **4b** (10.00 mmol) and H_2SO_4 (15 ml, 85%) was heated on a steam bath at 90-100 °C for 6 h. The resultant dark brown syrup was cooled to 0 °C and quenched by pouring into ice-cold water (150 mL) with vigorous stirring. Basification of the resulting solution with NaHCO_3 and the mixture was left to stand for 3h at room temperature. The oily residue was extracted with AcOEt (3×30 mL). The organic extracts was separated, washed with water, dried over MgSO_4 and concentrated to dryness to give the crude ketones **5a,b** as dark oil which crystallized on standing. Further purification and yields are given in the following.

2,3-Dihydro-5,8-dimethylquinolin-4(1H)-one (5a). Yellow crystals; 72%; mp 132–34°C (benzene); IR (KBr, ν , cm^{-1}): 3410 (N-H), 3040 (Ar-H), 2985 (C-H), 1700 (C=O), 1600, 1590, 1450, 1440, 1380, 1277, 1072, 766. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.19 (3H, s, CH_3), 2.33 (3H, s, CH_3), 2.62 (2H, ddd, J 14.8, 6.7, 2.4 Hz, $\text{NH}-\text{C}^2\text{H}_2$), 3.39 (2H, ddd, J 13.6, 6.7, 2.4 Hz, $-\text{C}^3\text{H}_2$), 6.68 (1H, d, J 8.1 Hz), 6.84 (1H, d, J 8.1 Hz), 9.74 (s, 1H, NH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 17.7 (1C, $-\text{CH}_3$), 20.0 (1C, $-\text{CH}_3$), 39.4 (1C, C^3H_2 , C-3), 41.4 (1C, C^2H_2 , C-2), 113.0 (1C, Ar.,C-4a), 128.9 (1C, Ar.,C-6), 129.2 (1C, Ar.,C-8), 130.3 (1C, Ar.,C-7), 136.8 (1C, Ar.,C-5), 137.2 (1C, Ar.,C-8a), 196.2 (1C, C=O, C-4). Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{NO}$ (175); C, 75.42; H, 7.42; N, 8.00. Found; C, 75.38; H, 7.50; N, 8.10 %.

1,2,3,4-Tetrahydro-6,9-dimethylbenzo[c]azepin-5-one (5b). Pale yellow needles; 75%; mp 114-16°C (ethanol); IR (KBr, ν , cm^{-1}): 3430 (N-H), 3062 (Ar-H), 2995 (C-H), 1695 (C=O), 1600, 1582, 1450, 1440, 1366, 1271, 1075, 690. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.25 (3H, s, CH_3), 2.40 (3H, s, CH_3), 2.76 (2H, ddd, J 15.6, 7.3, 3.7 Hz, $-\text{NH}-\text{C}^3\text{H}_2$), 3.08 (2H, ddd, J 9.9, 7.3, 3.7 Hz, $-\text{C}^4\text{H}_2$), 3.98 (2H, d, J 16.8 Hz, $-\text{C}^1\text{H}_2$), 7.02 (1H, d, J 8.2 Hz), 7.48 (1H, d, J 8.2 Hz), 10.06 (s, 1H, NH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 20.0 (2C, 2CH_3), 40.7 (1C, C^4H_2 , C-4), 41.6 (1C, C^3H_2 , C-3), 46.7 (1C, C^1H_2 , C-1), 126.4 (1C, Ar.,C-9), 128.8 (2C, Ar., C-7, C-8), 131.3 (1C, Ar.,C-6), 131.5 (1C, Ar.,C-9a), 136.3 (1C, Ar.,C-5a), 198.5 (1C, 1C, C=O, C-5). Anal. Calcd. for $\text{C}_{12}\text{H}_{15}\text{NO}$ (189); C, 76.19; H, 7.93; N, 7.40. Found; C, 76.15; H, 8.02; N, 7.36%.

General procedure for the synthesis of glycidic ester **6a,b**.

To an ice-cold stirred mixture of ketones **5a** or **5b** (10.02 mmol) and ethyl 2-bromopropanoate (10.08 mmol) in anhydrous ether (30 ml) was added dropwise over 1.5 h a solution of freshly prepared sodium ethoxide (1g, 15 mmol) in absolute ethanol (10 ml). After complete addition, the ice-bath was removed and the mixture was allowed to stir at room temperature for 7 h. Ice water (20 ml) was then added and the mixture was neutralized by cold HCl solution (5%). After standing for 2h in refrigerator, the ether solution was separated and the

mixture was extracted with ether (3×30 mL). The combined ether extracts was washed successively with water, saturated NaHCO₃ solution and finally with saturated NaCl solution. After drying over MgSO₄ and filtration, the solution was evaporated in *vacuo* to give crude glycidic ester **6a,b** as brownish viscous oil. These products were further purified by flash column chromatography (basic alumina, EtOAc/*n*-hexane, 1/1) gave pure esters as a brown oily products which crystallized on standing.

Ethyl 3,5',8'-trimethyl-2',3'-dihydro-1'H-spiro[oxirane-2,4'-quinoline]-3-carboxylate (6a). Brown solid; 68%; mp 64–66°C (ethanol); IR (KBr, ν , cm⁻¹): 3410 (N-H), 3010 (Ar-H), 2987 (C-H), 1745 (C=O), 1600, 1580, 1440, 1340, 1274, 1126, 794. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 1.31 (3H, t, *J* 7.1 Hz, CH₃CH₂O), 1.58 (3H, s, C³H₃), 1.93 (2H, ddd, *J* 14.5, 10.0, 2.0 Hz, C³H₂), 2.10 (3H, s, C⁵H₃), 2.34 (2H, ddd, *J* 14.2, 3.5, 2.0 Hz, C²H₂), 2.32 (3H, s, C⁸H₃), 3.12 (2H, ddd, *J* 14.9, 3.5, 2.0 Hz), 3.23 (ddd, *J* 14.9, 10.0, 2.0 Hz), 4.16 (2H, q, *J* 7.1 Hz, CH₃CH₂O), 6.46 (1H, d, *J* 8.0 Hz), 6.68 (1H, d, *J* 7.0 Hz), 9.84 (s, 1H, NH). Anal. Calcd. for C₁₆H₂₁NO₃ (275); C, 69.81; H, 7.63; N, 5.09. Found; C, 70.05; H, 7.62; N, 4.88%.

Ethyl 3',6,9-trimethyl-1,2,3,4-tetrahydrospiro[benzo[c]azepine-5,2'-oxirane]-3'-carboxylate (6b). Brown crystals; 72%; mp 80–83°C (ethanol); IR (KBr, ν , cm⁻¹): 3390 (N-H), 3030 (Ar-H), 2995 (C-H), 1740 (C=O), 1600, 1580, 1445, 1377, 1250, 1145, 690. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 1.33 (3H, t, *J* 7.1 Hz, CH₃CH₂O), 1.60 (3H, s, C³H₃), 2.09 (1H, ddd, *J* 14.8, 10.3, 2.8 Hz, C⁴Ha), 2.23 (6H, s, C⁶H₃ & C⁹H₃), 2.42 (1H, ddd, *J* 14.8, 2.9, 2.7 Hz, C⁴Hb), 2.81 (2H, ddd, *J* 10.3, 9.9, 2.9 Hz, -N-C³H₂), 3.82–4.04 (2H, d, *J* 15.1 Hz, C¹H₂), 4.01 (1H, d, *J* 15.1 Hz), 4.22 (2H, q, *J* 7.1 Hz, CH₃CH₂O), 6.86–6.94 (2H, d, *J* 8.2 Hz), 10.24 (s, 1H, NH). Anal. Calcd. for C₁₇H₂₃NO₃ (289); C, 70.58; H, 7.95; N, 4.84. Found; C, 70.40; H, 7.86; N, 5.02%.

General procedure for the synthesis of bicyclic ethanone-amines **7a,b**.

To an ice-cold stirred solution of sodium ethoxide (26.05 mmol, 1.78 g) in absolute ethanol (10 mL) was added water (1 mL) and the solution was stirred for 10 min. A solution of glycidic ester **6a** or **6b** (24.00 mmol) in absolute ethanol (20 mL) was added dropwise over a 10 min and the reaction was left to stir for 5 h at room temperature. The salt formed was filtered and washed with ice-cold ethanol (10 mL) to give the crude sodium salts which were used in the next transformation without further purification. The resulting crude sodium salts (15.02 mmol) was cooled in ice-bath and treated with glacial acetic acid (20 mL). A vigorous evolution of carbon dioxide occurred and after stirring at room temperature for around 20 min, the reaction had subsided. The reaction mixture was then heated on a steam-bath at 70–80 °C for 30 min or until the evolution of gases ceased. The reaction mixture was then cooled, diluted with water (150 mL), basified with NaHCO₃ and extracted with ether (3×30 mL). The combined organic extracts was washed with water (2×30 mL), with NaHCO₃ solution (10%, 30 mL), dried over MgSO₄. After filtration, the solution was concentrated under reduced pressure to afford crude ethanones **7a,b**.

1-(1,2,3,4-Tetrahydro-5,8-dimethylquinolin-4-yl)ethanone (7a). Yellow crystals; 80%; mp 120–22°C (benzene); IR (KBr, ν , cm⁻¹): 4050 (N-H), 3020 (Ar-H), 2980 (C-H), 1690 (C=O), 1600, 1580, 1440, 1380, 1270, 1100, 790. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.10 (3H, s, CH₃), 2.21 (3H, s, CH₃), 2.25 (3H, s, COCH₃), 2.40 (2H, dddd, *J* 13.5, 7.2, 6.9, 1.9 Hz, -NH-C²H₂), 3.25 (2H, ddd, *J* 14.3, 6.9, 1.9 Hz, -C³H₂), 3.65 (1H, dd, *J* 9.7, 4.6 Hz, -C⁴H), 6.41 (1H, d, *J* 8.0 Hz), 6.77 (1H, d, *J* 8.0 Hz), 9.80 (s, 1H, NH). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 17.7 (1C, -CH₃), 20.0 (1C, -CH₃), 29.0 (1C, -COCH₃), 29.8 (1C, CH, C-4), 41.6 (1C, C³H₂, C-3), 54.1 (1C, C²H₂, C-2), 128.0 (1C, Ar.,C-4a), 128.9 (1C, Ar.,C-6), 129.2 (1C, Ar.,C-7), 130.3 (1C, Ar.,C-8), 136.3 (1C, Ar.,C-5), 137.2 (1C, Ar.,C-8a), 208.4 (1C, C=O). Anal. Calcd. for C₁₃H₁₇NO (203); C, 76.84; H, 8.37; N, 6.89. Found; C, 76.88; H, 8.41; N, 6.82%.

1-(2,3,4,5-Tetrahydro-6,9-dimethyl-1H-benzo[c]azepin-5-yl)ethanone (7b). Yellow needles; 74%, m.p. 141–43°C (AcOEt); IR (KBr, ν , cm⁻¹): 3390 (N-H), 3070 (Ar-H), 2990 (C-H), 1715 (C=O), 1600, 1580, 1440, 1330, 1275, 1155, 790. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.08 (2H, dddd, *J* 13.4, 6.6, 4.8, 2.6 Hz, C⁴H₂), 2.15 (3H, s, CH₃), 2.21 (3H, s, CH₃), 2.27 (3H, s, COCH₃), 2.97 (2H, ddd, *J* 9.8, 6.6, 2.6 Hz, -NH-C²H₂), 3.82 (2H, d, *J* 14.5 Hz, C¹H₂),

3.93 (1H, dd, J 8.1, 1.4 Hz, C⁵H), 6.72 (1H, d, J 8.2 Hz), 6.79 (1H, d, J 8.2 Hz), 9.65 (s, 1H, NH). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 19.9 (2C, 2CH₃), 29.0 (1C, -COCH₃), 29.8 (1C, C⁴H₂, C-4), 41.6 (1C, CH, C-5), 46.7 (1C, C³H₂, C-3), 54.1 (1C, C¹H₂, C-1), 126.4 (1C, Ar.,C-9), 128.8 (2C, Ar., C-7, C-8), 132.9 (1C, Ar.,C-6), 136.3 (2C, Ar.,C-5a, C-9a), 210.0 (1C, C=O). Anal. Calcd. for C₁₄H₁₉NO (217); C, 77.42; H, 8.75; N, 6.45. Found; C, 77.52; H, 8.68; N, 6.42%.

General procedure for the synthesis of heterocyclic acids **8a,b**.

To a stirred solution of ethanone **7a** or **7b** (6.04 mmol) in dioxane (20 mL) was added KOH solution (3.5 g, 60.01 mmol) in water (10 mL). A solution of iodine (25 mmol) prepared from 6.4 g I₂ and 12.4 g KI in H₂O (30 mL) was added dropwise with stirring over 10 min at room temperature. The resulting yellow solution was left to stir at room temperature until fading the yellow color. The resultant solution was then heated on water-bath for 2 h at 60-70 °C. The resulting solution was evaporated to dryness in vacuo, cooled and the residue dissolved in hot H₂O (50 mL). After filtration, the solution was cooled and neutralized with ice-cold AcOH solution (10%) to pH 6-7. After standing for overnight in refrigerator, the resulting light tan solid was collected, washed with water and dried to give the crude acids **8a,b**.

1,2,3,4-Tetrahydro-5,8-dimethylquinoline-4-carboxylic acid (8a). Yellow crystals; 87%; mp 110-12 °C (ethanol); IR (KBr, ν , cm⁻¹): 3420 (N-H), 3060 (Ar-H), 2984 (C-H), 2640 (OH), 1718 (C=O), 1600, 1585, 1470, 1440, 1363, 1185, 793. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.01 (2H, dddd, J 13.3, 7.2, 6.9, 1.9 Hz, C²H₂), 2.10 (3H, s, CH₃), 2.26 (3H, s, CH₃), 3.19 (2H, ddd, J 13.9, 6.9, 1.9 Hz, -NH-C²H₂), 4.02 (1H, dd, J 9.7, 4.6 Hz, -C⁴H), 4.60 (1H, s, NH), 6.43 (1H, d, J 8.0 Hz), 6.77 (1H, d, J 8.0 Hz), 10.62 (1H, s, COOH). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 17.6 (1C, CH₃), 20.0 (1C, CH₃), 29.0 (1C, C³H₂, C-3), 41.6 (1C, C²H₂, C-2), 51.2 (1C, C⁴H, C-4), 128.0 (1C, Ar.,C-6), 128.9 (1C, Ar.,C-4a), 129.2 (1C, Ar.,C-7), 130.3 (1C, Ar.,C-8), 136.3 (1C, Ar.,C-5), 137.2 (1C, Ar.,C-8a), 174.5 (1C, COOH). Anal. Calcd. for C₁₂H₁₅NO₂ (205); C, 70.24; H, 7.31; N, 6.82. Found; C, 70.32; H, 7.33; N, 6.75%.

2,3,4,5-Tetrahydro-6,9-dimethyl-1H-benzo[c]azepine-5-carboxylic acid (8b). brownish solid; 82%; mp 154-56 °C (benzene); IR (KBr, ν , cm⁻¹): 3390 (N-H), 3010 (Ar-H), 2975 (C-H), 2590 (OH), 1715 (C=O), 1600, 1590, 1470, 1440, 1377, 1158, 765. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.00 (2H, dddd, J 13.3, 6.6, 4.7, 2.6 Hz, C⁴H₂), 2.15 (3H, s, CH₃), 2.23 (3H, s, CH₃), 3.00 (2H, ddd, J 8.7, 6.6, 2.6 Hz, -NH-C³H₂), 3.83 (2H, d, J 14.5 Hz, C¹H₂), 4.09 (1H, dd, J 8.1, 1.4 Hz, -C⁵H), 4.50 (1H, s, NH), 6.67 (1H, d, J 8.2 Hz), 6.76 (1H, d, J 8.2 Hz), 10.40 (1H, s, COOH). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 20.0 (2C, 2CH₃), 29.0 (1C, C⁴H₂, C-4), 41.6 (1C, C⁵H, C-5), 46.7 (1C, C³H₂, C-3), 51.2 (1C, C¹H₂, C-1), 126.4 (1C, Ar.,C-6), 128.9 (2C, Ar.,C-7, C-8), 132.9 (1C, Ar.,C-9), 136.3 (2C, Ar.,C-5a, C-9a), 176.6 (1C, COOH). Anal. Calcd. for C₁₃H₁₇NO₂ (219); C, 71.23; H, 7.76; N, 6.39. Found; C, 71.20; H, 7.82; N, 6.34%.

General procedure for N-arylation of heterocyclic acids **9a,b**.

To a mixture of acid **8a** or **8b** (8 mmol), K₂CO₃ (20 mmol), pyridine (1 mL) and CuI (0.3 g) in DMF (30 mL) was added slowly a solution of *PhBr* or *PhCH₂Cl* or *2-Picolyl chloride* (10 mmol) in DMF (20 mL). The reaction mixture was heated with continuous stirring for 10 h at 110-20 °C. The mixture was maintained at this temperature and monitored by TLC (30% AcOEt/hexane) until indicated that the reaction was complete. The mixture was then cooled and treated with NaOH solution (30 mL, 5%) and decolorizing carbon (1g). The mixture was then boiled for 20 min, cooled and filtered. The resulting filtrate was cooled to 0 °C and acidified using HCl solution (30 mL, 20%). After standing at refrigerator for overnight, the precipitate was filtered, washed and dried to afford the crude product **9a-d**. Further purification, yields and spectral data are given in the following:

1,2,3,4-Tetrahydro-5,8-dimethyl-1-phenylquinoline-4-carboxylic acid (9a). Pale yellow needles; 84%; mp 162-4 °C (ethanol); IR (KBr, ν , cm⁻¹): 3390 (OH), 3035 (Ar-H), 2950 (C-H), 2520, 1720 (C=O), 1600, 1585, 1470, 1440,

1370, 1144, 777. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.10 (2H, dddd, J 13.1, 6.9, 6.6, 2.5 Hz, $-\text{C}^3\text{H}_2$), 2.16 (3H, s, CH_3), 2.29 (3H, s, CH_3), 3.48 (2H, ddd, J 16.7, 6.6, 2.5 Hz, $-\text{NH}-\text{C}^2\text{H}_2$), 3.89 (1H, dd, J 10.1, 3.8 Hz, $-\text{C}^4\text{H}$), 6.43 (1H, d, J 7.9 Hz), 6.52 (2H, dtd, J 8.2, 1.2, 0.5 Hz), 6.80 (1H, d, J 7.9 Hz), 6.98 (1H, tt, J 8.1, 1.2 Hz), 7.22 (2H, dddd, J 8.2, 8.1, 1.3, 0.5 Hz), 10.61 (1H, s, COOH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 17.3 (1C, CH_3), 20.5 (1C, CH_3), 29.0 (1C, C^3H_2 , C-3), 50.1 (1C, CH, C-4), 51.2 (1C, C^2H_2 , C-2), 124.5 (2C, Ar., C-6, C-4'), 127.8 (1C, Ar., C-2'), 128.0 (1C, Ar., C-6'), 128.2 (2C, Ar., C-7, C-8), 128.9 (1C, Ar., C-3'), 130.3 (1C, Ar., C-5'), 136.3 (1C, Ar., C-4a), 137.1 (1C, Ar., C-5), 137.7 (1C, Ar., C-8a), 148.7 (1C, Ar., C-1'), 174.5 (1C, COOH). Anal. Calcd. for $\text{C}_{18}\text{H}_{19}\text{NO}_2$ (281); C, 76.86; H, 6.76; N, 4.98. Found; C, 76.90; H, 6.82; N, 4.90%.

1-Benzyl-1,2,3,4-tetrahydro-5,8-dimethylquinoline-4-carboxylic acid (9b). Yellow crystals; 75%; mp 155–57 °C (acetone); IR (KBr, ν , cm^{-1}): 3345 (OH), 3040 (Ar-H), 2966 (C-H), 2610, 1720 (C=O), 1610, 1580, 1480, 1445, 1364, 1195, 768; ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.09 (2H, dddd, J 13.5, 7.1, 6.9, 1.8 Hz, C^3H_2), 2.12 (3H, s, CH_3), 2.27 (3H, s, CH_3), 3.24 (2H, ddd, J 13.7, 6.9, 1.8 Hz, $-\text{NH}-\text{C}^2\text{H}_2$), 4.07 (1H, dd, J 9.8, 4.4 Hz, $-\text{C}^4\text{H}$), 4.32 (2H, s, PhCH_2), 6.43 (1H, d, J 8.0 Hz), 6.67 (1H, d, J 8.0 Hz), 7.19 (1H, tt, J 7.7, 1.3 Hz), 7.27 (2H, dddd, J 7.9, 7.7, 1.8, 0.5 Hz), 7.27 (2H, dddd, J 7.9, 1.3, 1.0, 0.5 Hz), 10.85 (1H, s, COOH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 17.7 (1C, CH_3), 20.0 (1C, CH_3), 29.0 (1C, C^3H_2 , C-3), 50.1 (1C, CH, C-4), 51.2 (1C, C^2H_2 , C-2), 54.7 (1C, benzylic- CH_2), 127.8 (3C, Ar., C-4a, C-6, C-8), 128.0 (1C, Ar., C-7), 128.4 (2C, Ar., C-3', C-5'), 128.9 (1C, Ar., C-4'), 130.3 (1C, Ar., C-2'), 136.3 (1C, Ar., C-6'), 136.6 (1C, Ar., C-5), 137.1 (1C, Ar., C-1'), 137.7 (1C, Ar., C-8a), 176.9 (1C, COOH). Anal. Calcd. for $\text{C}_{19}\text{H}_{21}\text{NO}_2$ (295); C, 77.28; H, 7.11; N, 4.74. Found; C, 77.35; H, 7.04; N, 4.76%.

1,2,3,4-Tetrahydro-5,8-dimethyl-1-((pyridin-2-yl)methyl)quinoline-4-carboxylic acid (9c). Pale yellow crystals; 72%; mp 129–31 °C (acetone); IR (KBr, ν , cm^{-1}): 3360 (OH), 3065 (Ar-H), 2985 (C-H), 2690, 1719 (C=O), 1605, 1590, 1485, 1440, 1384, 1155, 768; ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.14 (2H, dddd, J 13.5, 7.1, 6.9, 1.8 Hz, $-\text{C}^3\text{H}_2$), 2.10 (3H, s, CH_3), 2.27 (3H, s, CH_3), 3.17 (2H, ddd, J 13.8, 6.9, 1.8 Hz, $-\text{NH}-\text{C}^2\text{H}_2$), 4.07 (1H, dd, J 9.8, 4.4 Hz, C^4H), 4.47 (2H, s, PyCH_2), 6.43 (1H, d, J 8.0 Hz), 6.67 (1H, d, J 8.0 Hz), 7.20 (1H, ddd, J 7.4, 4.5, 1.2 Hz), 7.26 (1H, ddd, J 7.6, 1.2, 0.5 Hz), 7.63 (1H, ddd, J 7.6, 7.4, 1.9 Hz), 8.51 (1H, ddd, J 4.5, 1.9, 0.5 Hz), 11.20 (1H, s, COOH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 17.5 (1C, CH_3), 20.4 (1C, CH_3), 29.0 (1C, C^3H_2 , C-3), 50.1 (1C, CH, C-4), 51.2 (1C, benzylic- CH_2), 54.1 (1C, C^2H_2 , C-2), 122.7 (1C, Ar., C-6), 125.1 (1C, Ar., C-4'), 128.0 (1C, Ar., C-4a), 128.9 (1C, Ar., C-6'), 130.3 (1C, Ar., C-8), 136.3 (1C, Ar., C-7), 136.7 (1C, Ar., C-5), 137.1 (1C, Ar., C-5'), 137.7 (1C, Ar., C-8a), 149.1 (1C, Ar., C-3'), 157.8 (1C, Ar., C-1'), 177.4 (1C, COOH). Anal. Calcd. for $\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_2$ (296); C, 72.97; H, 6.75; N, 9.45. Found; C, 73.04; H, 6.82; N, 8.77%.

2,3,4,5-Tetrahydro-6,9-dimethyl-2-phenyl-1H-benzo[c]azepine-5-carboxylic acid (9d). Cream crystals; 80%; mp 141–43 °C (benzene); IR (KBr, ν , cm^{-1}): 3380 (OH), 3030 (Ar-H), 2982 (C-H), 2650, 1720 (C=O), 1600, 1585, 1470, 1445, 1385, 1164, 775. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.04 (2H, dddd, J 13.3, 6.6, 4.7, 2.6 Hz, $-\text{C}^4\text{H}_2$), 2.21 (3H, s, CH_3), 2.23 (3H, s, CH_3), 3.33 (2H, ddd, J 15.7, 6.6, 2.6 Hz, $-\text{NH}-\text{C}^3\text{H}_2$), 4.13 (1H, dd, J 8.1, 1.4 Hz, $-\text{C}^5\text{H}$), 4.49 (2H, d, J 15.6 Hz, $-\text{C}^1\text{H}_2$), 6.76 (1H, d, J 8.2 Hz), 6.87 (1H, d, J 8.2 Hz), 6.88 (1H, tt, J 8.1, 1.2 Hz), 6.92 (2H, dtd, J 8.3, 1.2, 0.5 Hz), 7.24 (2H, dddd, J 8.3, 8.1, 1.4, 0.5 Hz), 10.8 (s, 1H, COOH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 20.4 (2C, 2 CH_3), 29.0 (1C, C^4H_2 , C-4), 50.1 (1C, CH, C-5), 51.2 (1C, C^1H_2 , C-1), 53.5 (1C, C^3H_2 , C-3), 124.5 (2C, Ar., C-2', C-6'), 126.4 (1C, Ar., C-4'), 127.8 (1C, Ar., C-8), 128.2 (2C, Ar., C-3', C-5'), 128.8 (2C, Ar., C-6, C-9), 132.9 (1C, Ar., C-7), 136.4 (2C, Ar., C-5a, C-9a), 148.3 (1C, Ar., C-1'), 175.2 (1C, COOH). Anal. Calcd. for $\text{C}_{19}\text{H}_{21}\text{NO}_2$ (295); C, 77.28; H, 7.11; N, 4.74. Found; C, 77.22; H, 7.06; N, 4.81%.

2-Benzyl-2,3,4,5-tetrahydro-6,9-dimethyl-1H-benzo[c]azepine-5-carboxylic acid (9e). White plates; 82%; mp 162–64 °C (benzene); IR (KBr, ν , cm^{-1}): 3390 (OH), 3045 (Ar-H), 2974 (C-H), 2684, 1725 (C=O), 1590, 1575, 1470, 1440, 1381, 1145, 770. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.05 (2H, dddd, J 13.7, 6.6, 4.7, 2.6 Hz, $-\text{C}^4\text{H}_2$), 2.16 (3H, s, CH_3), 2.21 (3H, s, CH_3), 2.93 (2H, ddd, J 11.0, 6.6, 2.6 Hz, $-\text{NH}-\text{C}^3\text{H}_2$), 3.69 (2H, s, PhCH_2), 3.89 (2H, d, J 12.1 Hz, $-\text{C}^1\text{H}_2$), 4.12 (1H, dd, J 8.1, 1.4 Hz, $-\text{C}^5\text{H}$), 6.70 (1H, d, J 8.2 Hz), 6.87 (1H, d, J 8.2 Hz), 7.17 (1H, tt, J 7.7,

1.4 Hz), 7.24 (2H, dddd, J 7.7, 1.4, 0.9, 0.5 Hz), 7.30 (2H, tdd, J 7.7, 1.8, 0.5 Hz), 10.53 (1H, s, COOH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 20.5 (2C, 2CH_3), 29.0 (1C, C^4H_2 , C-4), 50.1 (1C, CH, C-5), 51.2 (1C, C^1H_2 , C-1), 53.5 (1C, C^3H_2 , C-3), 61.9 (1C, benzylic- CH_2), 126.4 (1C, Ar., C-4'), 127.8 (3C, Ar., C-3', C-5', C-8), 128.4 (2C, Ar., C-2', C-6'), 128.9 (2C, Ar., C-6, C-9), 132.9 (1C, Ar., C-7), 136.3 (2C, Ar., C-1', C-5a), 136.6 (1C, Ar., C-9a), 178.8 (1C, COOH). Anal. Calcd. for $\text{C}_{20}\text{H}_{23}\text{NO}_2$ (309); C, 77.66; H, 7.44; N, 4.53. Found; C, 77.74; H, 7.46; N, 4.45%.

2,3,4,5-Tetrahydro-6,9-dimethyl-2-((pyridin-2-yl)methyl)-1H-benzo[c]azepine-5-carboxylic acid (9f). Pale yellow needles; 73%; mp 172 °C *dec.* (acetone); IR (KBr, ν , cm^{-1}): 3410 (OH), 3080 (Ar-H), 2960 (C-H), 2510, 1715 (C=O), 1600, 1570, 1455, 1330, 1245, 1158, 779. ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.07 (2H, dddd, J 13.7, 6.6, 4.7, 2.6 Hz, $-\text{C}^4\text{H}_2$), 2.18 (3H, s, CH_3), 2.20 (3H, s, CH_3), 2.88 (2H, ddd, J 9.8, 6.6, 2.6 Hz, $-\text{NH}-\text{C}^3\text{H}_2$), 3.89 (2H, d, J 13.1 Hz, $-\text{C}^1\text{H}_2$), 3.94 (2H, s, PyCH_2), 4.12 (1H, dd, J 8.1, 1.4 Hz, $-\text{C}^5\text{H}$), 6.74 (1H, d, J 8.2 Hz), 6.87 (1H, d, J 8.2 Hz), 7.11 (1H, ddd, J 7.4, 4.6, 1.2 Hz), 7.25 (1H, ddd, J 7.7, 1.2, 0.5 Hz), 7.60 (1H, ddd, J 7.7, 7.4, 1.9 Hz), 8.51 (1H, ddd, J 4.6, 1.9, 0.5 Hz), 10.41 (1H, s, COOH). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 20.7 (2C, 2CH_3), 29.0 (1C, C^4H_2 , C-4), 50.1 (1C, CH, C-5), 51.2 (1C, C^1H_2 , C-1), 53.5 (1C, C^3H_2 , C-3), 54.1 (1C, benzylic- CH_2), 122.7 (1C, Ar., C-4'), 125.1 (1C, Ar., C-6'), 126.4 (1C, Ar., C-6), 128.8 (2C, Ar., C-7, C-8), 132.9 (1C, Ar., C-9), 136.3 (2C, Ar., C-5a, C-5'), 136.7 (1C, Ar., C-9a), 149.1 (1C, Ar., C-3'), 157.8 (1C, Ar., C-1'), 180.6 (1C, COOH). Anal. Calcd. for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_2$ (310); C, 73.54; H, 7.09; N, 9.03. Found; C, 73.51; H, 7.14; N, 8.96%.

General procedure for ring closure of heterocyclic acids 9a-f.

Method I. A solution of acid **9a-f** (4.00 mmol) in DCM (10 mL) was added dropwise to AlCl_3 (10.00 mmol) in CH_3NO_2 (80.00 mmol) over 10 min at ambient temperature. The mixture was stirred for the time given in Table 1. Afterwards, the mixture was quenched with ice-cold HCl solution (20 mL, 10%) and finally extracted with ether (3×20 mL). The combined organic layer was washed with H_2O and Na_2CO_3 solution (20 mL, 10%). After drying over MgSO_4 and filtration, the solvent was evaporated under reduced pressure to give the crude products **10a-f**.

Method II. TfOH (1.6 mL, 20.00 mmol) was added dropwise to a cooled (0 °C) solution of acid **9a-f** (4.00 mmol) in 1,2-Dichloroethane (15 mL) and the mixture was stirred at the required temperature for the time given in Table 1. Thereafter, the mixture was quenched by aqueous NaHCO_3 solution (30 mL, 50%) at 0 °C and the product was extracted with EtOAc (3×30 mL). The combined organic extracts were washed with H_2O , dried over anhydrous MgSO_4 and evaporated in *vacuo* to give the crude products **10a-f**.

Method III. A mixture of acid **9a-f** (4.00 mmol) and P_2O_5 (5 g) in toluene (15 mL) was refluxed for the time given in Table 1. Completion of the reaction was monitored by TLC-analysis (15% AcOEt /hexane). Afterwards, the mixture was cooled to room temperature and aqueous NaHCO_3 solution (30 mL, 50%) was added at 0 °C. The product was then extracted with EtOAc (3×30 mL). The combined organic extracts were washed with H_2O , dried over anhydrous MgSO_4 filtered, and concentrated under reduced pressure to afford the desired product **10a-f**.

In all procedures, the crude residue was subjected to flash chromatography (basic alumina, 25%, EtOAc /hexane) to afford the pure tetracyclic products **10a-f**. Further purification and yields of the products from the given methods are illustrating in the following.

6,9-Dimethyl-5,10-ethanodibenzo[b,f]azepin-11(10H)-one (10a). Yield 0.87 g (83%, method I), 0.79 g (75%, method II), 0.73 g (70%, method III), Pale yellow crystals; mp 140-143 °C (acetone); IR (KBr, ν , cm^{-1}): 3030 (Ar-H), 2955 (C-H), 1690 (C=O), 1600, 1590, 1440, 1377, 1275, 1180, 795; ^1H NMR (400 MHz, CDCl_3 , δ , ppm): 2.04 (2H, dddd, J 13.8, 7.9, 4.1, 3.1 Hz, C^{12}H_2), 2.16 (3H, s, CH_3), 2.28 (3H, s, CH_3), 3.99 (2H, ddd, J 12.8, 7.9, 4.1 Hz, C^{13}H_2), 4.28 (1H, dd, J 4.6, 1.6 Hz, C^5H_2), 6.48 (1H, d, J 8.0 Hz), 6.77 (1H, d, J 8.0 Hz), 7.09 (1H, ddd, J 8.4, 1.3, 0.5 Hz), 7.26 (1H, ddd, J 7.9, 7.4, 1.3 Hz), 7.36 (1H, ddd, J 7.9, 1.3, 0.5 Hz), 7.62 (1H, ddd, J 8.4, 7.4, 1.3 Hz). ^{13}C NMR (100 MHz, CDCl_3 , δ , ppm): 17.1 (1C, $-\text{CH}_3$), 20.6 (1C, $-\text{CH}_3$), 29.0 (1C, bridged- CH_2 , C-12), 50.1 (1C, CH, C-

10), 54.1 (1C, *bridged*-CH₂, C-13), 116.0 (1C, Ar., C-2), 121.9 (1C, Ar., C-8), 127.2 (1C, Ar., C-4), 128.0 (1C, Ar., C-11a), 128.2 (1C, Ar., C-9a), 128.4 (1C, Ar., C-7), 128.9 (1C, Ar., C-6), 130.3 (1C, Ar., C-1), 136.3 (1C, Ar., C-9), 137.1 (1C, Ar., C-3), 137.7 (1C, Ar., C-4a), 141.5 (1C, Ar., C-5a), 197.7 (1C, C=O, C-11). Anal. Calcd. for C₁₈H₁₇NO (263); C, 82.13; H, 6.46; N, 5.32. Found; C, 82.20; H, 6.49; N, 5.28%.

1,4-Dimethyl-6*H*-5,12-ethanodibenzo[*b,f*]azocin-11(12*H*)-one (10b). Yield 0.95 g (86%, method I), 0.85 g (77%, method II), 0.88 g (80%, method III), Yellow solid; mp 154-57 °C (acetone); IR (KBr, ν , cm⁻¹): 3040 (Ar-H), 2977 (C-H), 1700 (C=O), 1600, 1585, 1480, 1440, 1381, 1285, 1158, 786. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 1.98 (2H, dddd, *J* 13.3, 7.9, 3.5, 3.3 Hz, C¹³H₂), 2.1 (3H, s, CH₃), 2.26 (3H, s, CH₃), 3.44 (2H, ddd, *J* 14.4, 7.9, 3.5 Hz, C¹⁴H₂), 4.19 (1H, dd, *J* 5.1, 1.5 Hz, C¹²H), 4.46 (2H, d, *J* 13.8 Hz, C⁶H₂), 6.43 (1H, d, *J* 7.9 Hz), 6.67 (1H, d, *J* 7.9 Hz), 7.29 (1H, ddd, *J* 7.9, 1.4, 0.5 Hz), 7.37 (1H, ddd, *J* 7.9, 7.5, 1.4 Hz), 7.49 (1H, ddd, *J* 7.9, 7.5, 1.4 Hz), 7.86 (1H, ddd, *J* 7.9, 1.4, 0.5 Hz). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 17.7 (1C, -CH₃), 20.4 (1C, -CH₃), 29.0 (1C, *bridged*-CH₂, C-13), 50.1 (1C, CH, C-12), 53.5 (1C, *bridged*-CH₂, C-14), 54.1 (1C, CH₂, C-6), 126.4 (1C, Ar., C-2), 126.7 (1C, Ar., C-4), 128.0 (1C, Ar., C-12a), 128.5 (2C, Ar., C-3, C-7), 128.9 (1C, Ar., C-9), 129.8 (1C, Ar., C-10), 130.3 (1C, Ar., C-6a), 133.0 (1C, Ar., C-10a), 136.3 (1C, Ar., C-1), 137.1 (1C, Ar., C-8), 137.7 (1C, Ar., C-4a), 199.3 (1C, C=O, C-11). Anal. Calcd. for C₁₉H₁₉NO (277); C, 82.31; H, 6.85; N, 5.05. Found; C, 82.30; H, 6.92; N, 4.97%.

7,10-Dimethyl-6,12-dihydro-5*H*-6,11-ethanobenzo[*b*]pyrido[3,2-*f*]azocin-5-one (10c). Yield 0.91 g (82%, method I), 0.93 g (84%, method II), 0.84 g (76%, method III), White crystals; mp 170-173 °C (ethanol); IR (KBr, ν , cm⁻¹): 3070 (Ar-H), 2995 (C-H), 1688 (C=O), 1600, 1590, 1470, 1440, 1361, 1274, 1136, 765. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.09 (2H, dddd, *J* 13.3, 7.9, 3.5, 3.3 Hz, C¹⁴H₂), 2.10 (3H, s, CH₃), 2.26 (3H, s, CH₃), 3.59 (2H, ddd, *J* 14.3, 7.9, 3.5 Hz, C¹³H₂), 4.33 (1H, dd, *J* 5.1, 1.5 Hz, C⁶H), 4.63 (2H, d, *J* 13.3 Hz, C¹²H₂), 6.43 (1H, d, *J* 7.9 Hz), 6.67 (1H, d, *J* 7.9 Hz), 7.26 (1H, dd, *J* 7.8, 4.6 Hz), 7.88 (1H, dd, *J* 7.8, 1.9 Hz), 8.78 (1H, dd, *J* 4.6, 1.9 Hz). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 17.2 (1C, -CH₃), 21.1 (1C, -CH₃), 29.0 (1C, *bridged*-CH₂, C-14), 50.1 (1C, CH, C-6), 54.1 (2C, *bridged*-CH₂, C-12, C-13), 123.4 (1C, Ar., C-8), 128.0 (1C, Ar., C-3), 128.3 (1C, Ar., C-6a), 128.9 (1C, Ar., C-10), 130.3 (1C, Ar., C-9), 135.4 (1C, Ar., C-4a), 136.3 (1C, Ar., C-7), 137.1 (1C, Ar., C-4), 137.7 (1C, Ar., C-10a), 149.1 (1C, Ar., C-2), 155.8 (1C, Ar., C-12a), 199.0 (1C, C=O, C-5). Anal. Calcd. for C₁₈H₁₈N₂O (278); C, 77.69; H, 6.47; N, 10.07. Found; C, 77.76; H, 6.43; N, 10.02%.

7,10-Dimethyl-6*H*-5,11-ethanodibenzo[*b,f*]azocin-12(11*H*)-one (10d). Yield 0.86 g (78%, method I), 0.77 g (70%, method II), 0.78 g (71%, method III), Yellow crystals, mp 160-63 °C (benzene); IR (KBr, ν , cm⁻¹): 3020 (Ar-H), 2978 (C-H), 1690 (C=O), 1600, 1585, 1500, 1440, 1329, 1245, 1171, 790. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.15 (2H, dddd, *J* 13.7, 6.9, 4.3, 1.9 Hz, C¹³H₂), 2.21 (3H, s, CH₃), 2.25 (3H, s, CH₃), 3.61 (2H, ddd, *J* 13.2, 6.9, 1.9 Hz, C¹⁴H₂), 4.47 (1H, dd, *J* 6.9, 1.7 Hz, C⁶H), 4.79 (2H, d, *J* 13.9 Hz, C¹¹H₂), 6.67 (1H, d, *J* 8.2 Hz), 6.73 (1H, d, *J* 8.2 Hz), 6.85 (1H, ddd, *J* 8.1, 1.2, 0.5 Hz), 7.29 (1H, ddd, *J* 7.9, 7.4, 1.2 Hz), 7.62 (1H, ddd, *J* 8.1, 7.4, 1.4 Hz), 7.70 (1H, ddd, *J* 7.9, 1.4, 0.5 Hz). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 20.2 (2C, 2CH₃), 29.0 (1C, *bridged*-CH₂, C-13), 50.1 (1C, CH, C-11), 53.5 (1C, *bridged*-CH₂, C-14), 54.1 (1C, CH₂, C-6), 116.0 (1C, Ar., C-4), 121.9 (1C, Ar., C-12a), 126.4 (1C, Ar., C-2), 127.2 (1C, Ar., C-9), 128.2 (1C, Ar., C-8), 128.4 (1C, Ar., C-1), 128.8 (2C, Ar., C-7, C-10), 132.9 (1C, Ar., C-6a), 136.3 (2C, Ar., C-3, C-10a), 141.5 (1C, Ar., C-4a), 202.6 (1C, C=O, C-12). Anal. Calcd. for C₁₉H₁₉NO (277); C, 82.31; H, 6.86; N, 5.05. Found; C, 82.24; H, 6.80; N, 5.11%.

8,11-Dimethyl-7,12-dihydro-6,12-ethanodibenzo[*c,g*]azonin-13(5*H*)-one (10e). Yield 0.91 g (79%, method I), 0.95 g (83%, method II), 0.87 g (75%, method III), Yellow needles, mp 150-52 °C (AcOEt); IR (KBr, ν , cm⁻¹): 3010 (Ar-H), 2965 (C-H), 1700 (C=O), 1600, 1585, 1470, 1440, 1362, 1245, 1149, 792. ¹H NMR (400 MHz, CDCl₃, δ , ppm): 2.17 (3H, s, CH₃), 2.23 (3H, s, CH₃), 2.42 (2H, dddd, *J* 13.8, 6.7, 3.6, 2.7 Hz, C¹⁴H₂), 2.91 (2H, ddd, *J* 11.3, 6.7, 2.7 Hz, C¹⁵H₂), 3.97 (2H, d, *J* 15.7 Hz, C⁵H₂), 4.06 (2H, d, *J* 14.1 Hz, C⁷H₂), 4.23 (1H, dd, *J* 5.8, 1.5 Hz, C¹²H), 6.74 (1H, d, *J* 8.2 Hz), 6.88 (1H, d, *J* 8.2 Hz), 7.37 (1H, ddd, *J* 7.9, 7.5, 1.9 Hz), 7.48 (1H, ddd, *J* 8.1, 7.5, 1.3 Hz), 7.50 (1H, ddd, *J* 8.1, 1.9, 0.6 Hz), 7.83 (1H, ddd, *J* 7.9, 1.3, 0.6 Hz). ¹³C NMR (100 MHz, CDCl₃, δ , ppm): 20.0 (2C,

2CH₃), 29.0 (1C, *bridged*-CH₂, C-14), 50.1 (1C, CH, C-12), 53.5 (1C, CH₂, C-7), 54.1 (1C, *bridged*-CH₂, C-15), 55.2 (1C, CH₂, C-5), 126.5 (2C, Ar., C-1, C-9), 126.7 (1C, Ar., C-4), 128.3 (2C, Ar., C-2, C-10), 128.9 (2C, Ar., C-3, C-11), 129.8 (1C, Ar., C-4a), 132.8 (2C, Ar., C-8, C-13a), 136.4 (2C, Ar., C-7a, C-11a), 200.2 (1C, C=O, C-13). Anal. Calcd. for C₂₀H₂₁NO (291); C, 82.47; H, 7.21; N, 4.81. Found; C, 82.44; H, 7.30; N, 4.76%.

7,10-Dimethyl-11,13-dihydro-6,12-ethanobenzo[*c*]pyrido[3,2-*g*]azonin-5(6*H*)-one (10f). Yield 1.0 g (86%, method I), 0.84 g (72%, method II), 0.81 g (70%, method III), White crystals; mp 142-44 °C (acetone); IR (KBr, ν, cm⁻¹): 3050 (Ar-H), 2978 (C-H), 1690 (C=O), 1600, 1480, 1440, 1395, 1238, 1182, 765. ¹H NMR (400 MHz, CDCl₃, δ, ppm): 2.25 (3H, s, CH₃), 2.27 (3H, s, CH₃), 2.42 (2H, dddd, *J* 13.8, 6.7, 3.6, 2.7 Hz, C¹⁵H₂), 2.96 (2H, ddd, *J* 11.3, 6.7, 2.7 Hz, C¹⁴H₂), 4.03 (2H, d, *J* 14.1 Hz, C¹¹H₂), 4.25 (1H, dd, *J* 5.8, 1.5 Hz, C⁶H), 4.36 (2H, d, *J* 15.7 Hz, C¹³H₂), 6.74 (1H, d, *J* 8.5 Hz), 6.88 (1H, d, *J* 8.5 Hz), 7.20 (1H, dd, *J* 7.8, 4.5 Hz), 7.89 (1H, dd, *J* 7.8, 1.9 Hz), 8.89 (1H, dd, *J* 4.5, 1.9 Hz). ¹³C NMR (100 MHz, CDCl₃, δ, ppm): 20.1 (2C, 2CH₃), 29.0 (1C, *bridged*-CH₂, C-15), 50.1 (1C, CH, C-6), 53.5 (1C, CH₂, C-13), 54.1 (2C, *bridged*-CH₂, C-11, C-14), 123.4 (1C, Ar., C-3), 126.4 (1C, Ar., C-9), 128.3 (1C, Ar., C-8), 128.9 (2C, Ar., C-7, C-10), 132.9 (1C, Ar., C-4a), 135.4 (1C, Ar., C-10a), 136.3 (2C, Ar., C-4, C-6a), 149.1 (1C, Ar., C-2), 155.8 (1C, Ar., C-13a), 199.7 (1C, C=O, C-5). Anal. Calcd. for C₁₉H₂₀N₂O (292); C, 78.08; H, 6.84; N, 9.58. Found; C, 78.11; H, 6.79; N, 9.64%.

Acknowledgements

The authors offer their appreciation and gratitude to Assiut University for funding this work as research number AUN2024F.SCI.L0002 and for all the facilities received while performing and writing this work by the Chemistry department, Faculty of Science Assiut University, Assiut, Egypt.

Supplementary Material

Supplementary data associated with this article is available in the Supplementary Material.

References

1. Buckingham, J., Baggaley, K. H., Roberts, A. D., Szabó, L. F. *Dictionary of Alkaloids* 2nd Ed.; CRC Press: USA, 2010.
<https://doi.org/10.1201/EBK1420077698>
2. Tewari, A. K.; Singh, V. P.; Yadav, P.; Gupta, G.; Singh, A.; Goel, R. K.; Shinde, P.; Mohan, C. G., *Bioorg. Chem.* **2014**, 56, 8.
<https://doi.org/10.1016/j.bioorg.2014.05.004>
3. Doxsee, K. M., in *Comprehensive Heterocyclic Chemistry II* Katritzky, A. R.; Rees, C. W.; Scriven, E. F. V., Eds.; Pergamon, Oxford, 1996, Vol. 9, p 527 and references cited therein.
4. Enzensperger, C.; Kilian, S.; Ackermann, M.; Koch, A.; Kelch, K.; Lehmann, J. *Bioorg. Med. Chem. Lett.* **2007**, 17, 1399.
<https://doi.org/10.1016/j.bmcl.2006.11.093>
5. Robaa, D.; Enzensperger, C.; AbulAzm, S. E.; Hefnawy, M. M.; Subbagh, H. I. E.; Wani, T. A.; Lehmann, J. *J. Med. Chem.* **2011**, 54, 7422.

- <https://doi.org/10.1021/jm200676f>
6. Villar, H.; Frings, M.; Bolm, C., *Chem. Soc. Rev.* **2007**, 36, 55.
<https://doi.org/10.1039/B508899M>
7. Kalshetti, M. G.; Argade, N. P., *J. Org. Chem.* **2017**, 82, 11126.
<https://doi.org/10.1021/acs.joc.7b02122>
8. Liu, R.; Huang, X.-P.; Yeliseev, A.; Xi, J.; Roth, B. L. *Anesthesiology* **2014**, 120, 714.
<https://doi.org/10.1097/ALN.000000000000076>
9. Wesnes, K. A.; Edgar, C.; Dean, A. D.; Wroe, S. J. *Epilepsy Behav.* **2009**, 14, 522.
<https://doi.org/10.1016/j.yebeh.2008.11.012>
10. Oniciu, D. C. in *Comprehensive Heterocyclic Chemistry III* Katritzky, A. R.; Ramsden, C. A.; Scriven, E. F. V.; Taylor, R. J. K. Eds.; Pergamon Press: New York, 2008; Vol. 14, pp 1–47 and references cited therein.
11. Yao, T.; Li, J.; Jiang, C.; Zhao, C. *Chem. Catalysis*, **2022**, 2, 2929.
<https://doi.org/10.1016/j.checat.2022.07.021>
12. Hudicky, T. in *Alkaloids: Chemical and Biological Perspectives* S.W. Pelletier Ed., Vol. 5, Wiley-Interscience, New York, 1987.
13. Illuminati, G.; Mandolini, L., *Acc. Chem. Res.* **1981**, 14, 95.
<https://doi.org/10.1021/ar00064a001>
14. Paquette, L. A.; Hartung, R. E.; Hofferberth, J. E.; Vilotijevic, I.; Yang, J. *J. Org. Chem.* **2004**, 69, 2454.
<https://doi.org/10.1021/jo0358675>
15. Gil, L.; de Freitas Gil, R. P.; dos Santos, D. C.; Marazano, C. *Tetrahedron Lett.* **2000**, 41, 6067.
[https://doi.org/10.1016/S0040-4039\(00\)01008-X](https://doi.org/10.1016/S0040-4039(00)01008-X)
16. Raw, S. A.; Taylor, R. J. K., *J. Am. Chem. Soc.* **2004**, 126, 12260.
<https://doi.org/10.1021/ja045780g>
17. Cho, H.; Iwama, Y.; Sugimoto, K.; Mori, S.; Tokuyama, H. *J. Org. Chem.* **2010**, 75, 627.
<https://doi.org/10.1021/jo902177p>
18. Kisel, V. M.; Kovtunenkov, V. A.; Turov, A. V.; Tyltin, A. K.; Babichev, F. S., *Chem. Heterocycl. Compd.* **1991**, 27, 92.
<https://doi.org/10.1007/BF00633227>
19. Polonka-Bálint, Á.; Saraceno, C.; Ludányi, K.; Bényei, A. Mátyus, P. *Synlett* **2008**, 2846.
<https://doi.org/10.1055/s-0028-1083537>
20. Sharma, A.; Appukkuttan, P.; Van der Eycken, E. *Chem. Commun.* **2012**, 48, 1623.
<https://doi.org/10.1039/C1CC15238F>
21. Schrock, R. R.; Hoveyda, A. H., *Angew. Chem. Int. Ed.* **2003**, 42, 4592.
<https://doi.org/10.1002/anie.200300576>
22. Majumdar, K. C.; Nandi, R. K.; Samanta, S.; Chattopadhyay, B. *Synthesis* **2010**, 985.
<https://doi.org/10.1055/s-0029-1218630>
23. Nagata, T.; Nakagawa, M.; Nishida, A. *J. Am. Chem. Soc.* **2003**, 125, 7484.
<https://doi.org/10.1021/ja034464j>
24. Khalaf, A. A.; Roberts, R. M. *J. Org. Chem.*, **1970**, 35, 3717.
<https://doi.org/10.1021/jo00836a029>
25. Abd El-Aal, H. A. K., *Chem. Heterocycl. Compd.* **2020**, 56, 1353.
<https://doi.org/10.1007/s10593-020-02822-1>
26. Abd El-Aal, H. A. K. *Arkivoc* **2023**, (vii) 202311986
<https://doi.org/10.24820/ark.5550190.p011.986>

27. Abd El-Aal, H. A. K. *Aust. J. Chem.* **2023**, 76, 760.
<https://doi.org/10.1071/CH23030>
28. Bruson, H. A.; Riener, T. W. *J. Am. Chem. Soc.* **1943**, 65, 23.
<https://doi.org/10.1021/ja01241a007>
29. Newman, M. S.; Magerlein, B. J., *Org. React.* **1949**, Vol 5, 413.
30. Cano, R.; Ramón, D. J.; Yus, M. *J. Org. Chem.* **2011**, 76, 654.
<https://doi.org/10.1021/jo1022052>
31. Carey, F. A.; Sundberg, R. J. *Advanced Organic Chemistry: Part B: Reactions and Synthesis* Kluwer Academic/Plenum Publishers, New York, NY, 4th Ed, **2001**, page 693.
32. Anderson, H.J.; Loader, C.E.; Xu, R.X.; Le, N.; Gogan, N.J.; McDonald, R.; Edwards, L.G. *Can. J. Chem.* **1985**, 63, 896.
<https://doi.org/10.1139/v85-149>
33. Olah, G.A.; Tashiro, M.; Kobayashi, S. *J. Am. Chem. Soc.*, **1970**, 92, 6371.
<https://doi.org/10.1021/ja00724a063>
34. Santiago, C.; Houk, K.N.; Perrin, C.L. *J. Am. Chem. Soc.* **1979**, 101, 1337.
<https://doi.org/10.1021/ja00499a077>
35. March, J. *Advanced Organic Chemistry* John Wiley and Sons, New York, 4th, Ed, 1999-2000.
36. Huffman, J.W.; Smith, V.J.; Padgett, L. W. *Tetrahedron* **2008**, 64, 2104.
<https://doi.org/10.1016/j.tet.2007.12.043>
37. Katritzky, A. R.; Xu, Y.-J.; He, H.-Y., *J. Chem. Soc., Perkin Trans. 1*, **2002**, 592. and references cited therein.
38. Paquette, L. A.; Tae, J.; Arrington, M.; Sadoun, A. H. *J. Am. Chem. Soc.* **2000**, 122, 2742.
<https://doi.org/10.1021/ja994053w>
39. Anand, R. V.; Baktharaman, S.; Singh, V. K. *J. Org. Chem.* **2003**, 68, 3356.
<https://doi.org/10.1021/jo034049+>
40. Maier, M. E. *Angew. Chem. Int. Ed.* **2000**, 39, 2073.
[https://doi.org/10.1002/1521-3773\(20000616\)39:12<2073::AID-ANIE2073>3.0.CO;2-0](https://doi.org/10.1002/1521-3773(20000616)39:12<2073::AID-ANIE2073>3.0.CO;2-0)
41. Fürstner, A.; Müller, T. *Synlett* **1997**, 1010.
<https://doi.org/10.1055/s-1997-930>
42. Becker, E. *High Resolution NMR Theory and Chemical Application* 3rd Ed., Academic Press, 2000, pp 139.
<https://doi.org/10.1016/B978-012084662-7/50050-8>
43. Hanack, M., *Conformational Theory*, Academic Press, New York, 1965.
44. Eliel, E. L.; Wilen, S. H., *Stereochemistry of Organic Compounds*, Wiley, New York, 1994.
<https://doi.org/10.1002/9780470147306>
45. Haufe, G.; G. Mann, *Chemistry of Alicyclic compounds*, Elsevier, Amsterdam, 1989.
46. Majeski, K. M.; Vinkovic, M.; Skare, D.; Marchand, A. P. *Arkivoc* **2002**,(iv), 30.
<https://doi.org/10.3998/ark.5550190.0003.404>
47. Anet, F. A. L. *Conformational Analysis, Scope and Present limitations* Academic Press, New York, 1971.

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)