

Synthesis and antimicrobial activity of novel pyrido[2,3-*d*]pyrimidin-4(1*H*)-ones, and 2-pyrazoline derivatives

Ahmed Abdou O. Abeed,^{a *} Mohamed Mohany,^b Sinisa Djurasevic,^c Salim S. Al-Rejaie,^b and Talaat I. El-Emary^a

^a Department of Chemistry, Faculty of Science, Assiut University, Assiut 71516, Egypt

^b Department of Pharmacology and Toxicology, College of Pharmacy, King Saud University, P.O. Box 55760, Riyadh 11451, Saudi Arabia

^c Faculty of Biology, University of Belgrade, 11158 Belgrade 118, Serbia

Email: ahmed.abeed76@aun.edu.eg

Received mm-dd-yyyy


Accepted mm-dd-yyyy

Published on line mm-dd-yyyy

Dates to be inserted by editorial office

Abstract

The synthesis of chalcone **1** allowed the production of a novel group of compounds by linking indole with thiophene moieties. The new series of substituted pyridines **2-8**, 2-pyrazolines **9-14**, and azepines **15, 16** were synthesized using chalcone **1** as a crucial intermediate. 2-Aminopyridines **2, 3**, and **4** were produced by reacting compound **1** with cyanoacetamide, ethyl cyanoacetate, and malononitrile, respectively. The α,β -unsaturated ketone **1** underwent conversion to 2-thioxodihydropyridopyrimidinone **5** and 2-hydrazinyl pyridopyrimidinone **6**. These compounds served as the initial substrate for the synthesis of compounds **7** and **8**. 1-Thiocarbamoyl-2-pyrazoline **9** was synthesized by the cycloaddition reaction between thiosemicarbazide and chalcone. 1-(4-oxo-5*H*-thiazolyl)-2-pyrazoline **10** and 1-(4-phenylthiazolyl)-2-pyrazoline **11** were synthesized by treating thiocarbamoyl derivative **9** with chloroacetic acid or phenacyl bromide. Condensation of thiazolyl pyrazoline **10** with aromatic aldehydes yielded the arylidenes **12-14**. The α,β -unsaturated ketone **1** underwent a reaction with bifunctional agents such as *o*-phenylenediamine or *o*-aminophenol, resulting in the formation of azepines **15** and **16**. The novel compounds were structurally determined with spectral data including FTIR, ¹H NMR, ¹³C NMR, and MS spectroscopy as well as elemental analysis. In addition, the new substances were tested for antibacterial activity versus an array of *Gram* positive and *Gram*-negative bacteria, as well as four fungal species. Our findings revealed that 2-pyrazolines **9-14** had remarkable effects when compared to the other compounds.



Keywords: Pyridines, 2-pyrazolines, azepines, indole, thiophene, antimicrobial potency


Introduction

In 1846, Anderson discovered pyridine, an aromatic heterocyclic compound. Subsequently, Korner (1869) and Dewar (1871) determined the pyridine structure. It is a constituent of almost 7000 existing pharmacological compounds having medicinal characteristics. Pyridine has a vital role in the field of chemistry [1,2]. *N,N*-dimethyl-*N*-(4-methylpyridyl)-*N*-alkyl-ammonium chloride **A** is an example a pyridine analog that shows potency and sensitivity for fungal and bacterial strains [3]. Furthermore, clinical and theoretical research on pyrazolopyridine-5-carboxylic acids **B**, showed antibacterial action toward a drug-resistant *S. epidermidis* [4]. Studies have shown that thienopyridines are effective against the two kinds of bacteria [5]. Research findings suggest that pyridine derivatives exhibit significant effectiveness in combating *Herpes simplex* virus type 1 (HSV-1) [6], as well as exhibiting anticancer [7-10], anti-diabetic [11,12], and anti-inflammatory properties [13-15].

2-Pyrazoline is a saturated pyrazole distinguished by the presence of two next atoms of nitrogen and an endocyclic double bond within its ring framework. Due to its superior stability concerning the other two forms, 2-pyrazoline is the most well-researched among the three forms of pyrazolines, namely 1-pyrazoline, 2-pyrazoline, and 3-pyrazoline. Research has demonstrated that 2-pyrazolines **C** exhibits potent anticancer effects on cervical cancer cells from humans (HeLa) and breast cancer cells from humans (MCF7) [16]. In addition, nicotinoyl pyrazolines **D** showed great effectiveness on cancer cells when employed as inhibitors of tubulin assembly [17] (**Figure 1**). Several 2-pyrazoline structures exhibited notable biological properties, such as considerable antiproliferative activity [18-20], antibacterial and antifungal effects [21-27], anti-hyperglycemic activity [28], as well as anti-inflammatory and analgesic properties [29-32]. Moreover, it is noteworthy that 3-indolecarboxaldehyde and 2-acetyl thiophene are widely used as fundamental components for synthesizing a wide variety of heterocyclic compounds that have significant biological functionalities [33-46].

Based on the foregoing and a review of our work [31], we decided to produce a new category of pyridine, 2-pyrazoline, and azepine derivatives by reacting chalcone **1** composed of indole and thiophene moieties with different active chemical reagents. The newly synthesized compounds were evaluated for their antimicrobial activity.

Some of reported pyridine and 2-pyrazoline derivatives with biological importance



The novel synthesized pyridine, pyridopyrimidine and 2-pyrazoline analogs





Figure 1. The reported and new synthesized biologically active pyridine, pyridopyrimidine, and 2-pyrazolines.

Results and Discussion


The key starting material, thienyl indoyl propenone **1**, was generated through the Claisen-Schmidt procedure by reacting 2-acetylthiophene and indole-3-carboxaldehyde [47]. The chalcone **1** was interacted with each of cyanoacetamide, ethyl cyanoacetate, and malononitrile, in the presence of ammonium acetate in hot ethanol, to provide 2-amino-3-carboxamidepyridine **2**, ethyl 2-aminothienyl indolyl pyridine-3-carboxylate **3**, and 2-amino-3-cyanothienyl indolyl pyridine **4**, respectively (**Scheme 1**).



Scheme 1. The synthesis of 2-aminopyridines (**2-4**).


The structures of products **2-4** were verified utilizing spectral data (FTIR, NMR, and mass) and elemental analysis. Thus, Fourier Transform Infrared (FTIR) spectrum of product **2** revealed absorbance peaks at ν 3446, 3239, and 3098 cm^{-1} due to pyridine-NH₂ and indole-NH, as well as a peak at ν 1628 cm^{-1} belonging to the carbonyl group. The ¹H NMR spectrum indicated that signals from two amino groups come together with aromatic proton signals at δ 7.01-7.84 ppm. Similarly, the existence of carbonyl carbon was verified by signals in the ¹³C NMR spectrum with δ = 166.7 ppm. The FTIR analysis for **3** revealed bands of absorption at ν 445, 3374, and 3333 cm^{-1} , which are associated with the amino group and indole-NH. Further, absorbance at ν 1711 cm^{-1} corresponds to the carbonyl group of the ester. A signal at δ 176.9 ppm was detected in the ¹³C NMR of **3**, indicating the presence of the ester carbonyl group attributable to the carbonyl group. The ¹H NMR displayed peaks at δ 11.00 and 13.25 ppm, due to the NH.

Chalcone **1** and 6-amino-2-thiouracil were refluxed in acetic acid to yield pyridopyrimidinone **5**. The FTIR spectrum of the latter compound exhibited peaks at ν 3390, 3307, and 3176 cm^{-1} belonging to pyrimidine-NH and indole-NH, as well as absorption bands at ν 1659 cm^{-1} of the pyrimidine. The ¹³C NMR of **5** indicated two signals at δ 165.2 and 176.0 ppm correlating to C=O and C=S of the pyrimidine ring, respectively. To examine the synthetic potential of compound **5**, it was heated with hydrazine hydrate to produce 2-hydrazinyl pyridopyrimidinone **6** that was then employed as a reactant with active methylene reagents like acetylacetone and ethyl acetoacetate, yielding 2-pyrazolyl pyridopyrimidines **7** and **8**. The product structures **6-8** were validated using elemental and spectroscopic data.



Scheme 2. The synthesis of pyrido[2,3-*d*]pyrimidines (**5-8**).

When chalcone **1** was heated with thiosemicarbazide, it afforded 1-thiocarbamoyl-2-pyrazoline **9**. FTIR Spectroscopy of **9** showed the existence of an amino functional group, revealing two peaks at ν 3447 and 3312 cm^{-1} . The ¹H NMR spectrum had three doublets of doublets at δ 3.68-3.76, 3.90-3.98, and 6.29-6.35 ppm, owing to the H_A, H_B, and H_X, indicating the formation of a 2-pyrazoline nucleus. The ¹³C NMR spectrum was consistent with FTIR and ¹H NMR, with three distinct signals at δ 49.1 for pyrazoline-CH₂, 66.8 for pyrazoline-CH, and 177.0 ppm due to the C=S group. When α -halocarbonyl reagents such as chloroacetic acid or phenacyl bromide reacted with thiocarbamoyl derivative **9**, the products 1-(4-oxo-5*H*-thiazol-2-yl)-2-pyrazoline **10** or 1-(4-phenylthiazol-2-yl)-2-pyrazoline **11** were formed. The reactivity of the methylene group at the thiazole moiety in product **10** was assessed by treating it using various aromatic aldehydes like benzaldehyde, *p*-chlorobenzaldehyde, and *p*-bromobenzaldehyde to produce the arylidenes **12-14** (Scheme 3). The products **9-14** were fully verified through elemental analysis together with spectral data such as the FTIR, NMR, with MS spectrometry. Chalcone **1** underwent treatment with *o*-phenylene diamine or *o*-aminophenol to obtain benzodiazepine **15** and benzoxazepine **16**, respectively. The formation of benzodiazepine **15** was verified through FTIR, ¹H- and ¹³C NMR. The FTIR diagram of **15** indicated the lack of the chalcone carbonyl group as well as the existence of an absorption bands at ν 3333 and 3172 cm^{-1} belonging to diazepine-NH and indole-NH. The ¹H NMR spectrum of **15** verified the formation of the benzodiazepine nucleus through the appearance of three doublets of doublets at δ 3.13-3.20 for H_A, 3.81-3.88 for H_B, and 5.00-5.56 for H_X. In addition, the ¹³C NMR diagram confirmed the absence of a carbonyl group of ketone **1** and also the presence of two signals at 18.5 and 46.2 for CH₂ and CH of the dihydroazepine ring.



Scheme 3. The synthesis of 2-pyrazolines (**9-11**) and azepines (**15** and **16**).

Antimicrobial screening

Antibacterial activity. All compounds **2-16** were evaluated for antibacterial effects against both *Gram*-positive (*Bacillus cereus*, *Staphylococcus aureus*) and *Gram*-negative (*Pseudomonas aeruginosa*, *Escherichia coli*). *Chloramphenicol* was selected as the reference (**Table 1**). The minimum inhibitory concentrations (*MICs*) were calculated. Each of the compounds displayed considerable antibacterial effects. The compounds, 2-aminopyridines **2-4** and pyrido[2,3-*d*] pyrimidin-4(1*H*)-ones **5-8** displayed moderate antibacterial properties.

2-Pyrazoline derivatives **9-14** exhibit significant activity against tested against other antibacterial species, with *MICs* ranging from 4-30 µg/mL. When compared to pyridines **2-8**, the azepine analogs had significantly higher antibacterial efficacy. Compounds **10** and **11**, containing 2-pyrazoline and thiazole moieties, showed high potency, and this was increased after inserting the aryl group, giving the arylidene **12-14**. 1-(5-*p*-Bromobenzyliden-4-oxothiazol-2-yl)-3-(thien-2-yl)-5-(1*H*-indol-3-yl)-2-pyrazoline (**14**) displayed the most potent antibacterial activity against *B. cereus*, *S. aureus*, *P. aeruginosa*, and *E. coli*, with *MICs* of 4, 15, 14, and 18 µg/ml, respectively. *B. cereus* was the most sensitive bacterial species against 1-(5-*p*-bromobenzyliden-4-oxothiazol-2-yl)-3-(thien-2-yl)-5-(1*H*-indol-3-yl)-2-pyrazoline (**14**) possesses a *MIC* of 4 µg/mL.

Antifungal activity. Compounds **2-16** were tested for antifungal action against the following species: *Geotrichum candidum*, *Candida albicans*, *Trichophyton rubrum*, and *Aspergillus flavus*. *Clotrimazole* was selected as the reference drug (**Table 2**). Similar to antibacterial actions, arylidene derivatives **12-14** exhibited high antifungal potency with *MICs* that ranged from 30-50 µg/mL. Compound **14** showed unique *MICs* of 4, 15, 14, and 18 µg/mL against fungal species; *G. candidum*, *C. albicans*, *T. rubrum*, and *A. flavus*, respectively. Compounds **15** and **16** having azepine nucleus generated acceptable results, with *MICs* ranging from 15 to 38 µg/mL. *G. candidum* was one of the most inhibited fungi species from 4-(thienyl-2-yl)-2-(indol-3-yl)-2,3-dihydrobenzo [*b*][1,4]oxazepine (**16**) has a *MIC* of 15 µg/mL.

Table 1. The antibacterial activity of the synthesized compounds, *MIC* ($\mu\text{g/mL}$)

Compound No.	<i>Gram positive</i>		<i>Gram negative</i>	
	<i>Bacillus cereus</i>	<i>Staphylococcus aureus</i>	<i>Pseudomonas Aruginose</i>	<i>Escherichia Coli</i>
2	90	100	90	97
3	50	80	92	95
4	40	70	89	88
5	45	72	90	91
6	42	71	80	75
7	41	78	75	70
8	25	35	50	64
9	10	25	24	25
10	21	20	30	2
11	10	22	25	28
12	11	18	19	16
13	12	17	18	19
14	4	15	14	18
15	18	19	20	25
16	20	20	26	21
Reference*	10	10	10	10

*Chloramphenicol was used as antibacterial standard. (-) no activity.

Table 2. The antifungal activity of the synthesized compounds, MIC ($\mu\text{g/mL}$).

Compound No.	Pathogenic Fungi			
	<i>Geotrichum candidum</i>	<i>Candida albicans</i>	<i>Trichophyton rubrum</i>	<i>Aspergillus flavus</i>
2	-	110	100	98
3	115	105	95	90
4	100	98	-	87
5	94	90	82	77
6	88	89	76	-
7	74	-	-	-
8	70	72	64	-
9	67	68	60	70
10	64	60	57	61
11	50	55	52	57
12	42	50	50	50
13	38	48	49	46
14	30	42	40	40
15	25	37	38	37
16	15	30	30	20
Reference*	20	10	10	10

* Clotrimazole was used as antifungal standard.

(-) no activity.

Conclusions

This work presents a novel group of pyridines, pyridopyrimidines, 2-pyrazolines, and azepines structured on thiophene and indole nuclei. The chemical reactivity of the starting compound thienyl indolyl 2-propenone **1** was studied by its reactivity with various reagents. 2-Aminopyridines **2**, **3**, and **4** were obtained using active methylene compounds. In addition, 2-thioxopyridopyrimidinone **5** was achieved by the treatment of 6-amino-2-thiouracil with **1**. By reacting **5** with a variety of reagents, 2-substituted pyridopyrimidinones **6-8** were generated. Furthermore, 2-pyrazolines **9-14** and azepines **16**, and **17** derive from the synthesized starting compound **1**. Spectroscopic data (FTIR, NMR, and MS spectrometry) with the analysis of elements elucidated the compounds structures. The new compounds **2-16** have been evaluated for antimicrobial activity. The findings showed that 2-pyrazolines **9-14** have greater antibacterial activity than pyridine derivatives **2-4**. Furthermore, arylidenes with bromine atom **14** are more active than those with chlorine atom **13**. Azepines **15** and **16** (with seven membered rings) are more effective than compounds **2-4**, which has six members.

Experimental Section

General. In this work, the analytical-quality substances were used. The melting points were determined via the melting point instrument, APP Digital ST 15. FTIR spectra were obtained with the Shimadzu-408 infrared

spectrophotometer and are shown in cm^{-1} mode. NMR spectra were gathered using a Bruker AV-400 spectrometer. TMS was used as a standard. Mass spectrometry was conducted with a Varian MAT 312 apparatus operating in EI mode at a scan energy of 70 eV. System GmbH vario EL V2.3 1998 CHNS Mode was used to analyze elements.

General approach for the synthesis of 2-aminopyridines 2-4. A mixture of **1** (0.50 g, 2 mmol), active methylene reagents (2 mmol), and AcONH_4 (0.25 g, 3 mmol) was heated at reflux in AcOH (10 mL) for 6 h. Upon cooling, the separated product was filtered off, rinsed using water, and purified with ethanol.

2-Amino-3-carboxamide-6-(thien-2-yl)-4-(indole-3-yl)pyridine (2). Yield: 77%, white crystals: mp 180-183 °C. IR (ν/cm^{-1}): 3446, 3239, 3098 (br. NH_2 and indole-NH), 3052 (Ar-H), 1628 (CO); ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ = 7.01-7.84 (m, 7 H, 3 thienyl-H and 2NH_2), 7.97-8.54 (m, 6H, 5 indole-H and pyridine-H), 12.13 (s, 1H, indole-NH); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 112.9, 113.1, 115.6, 120.9, 121.6, 123.2, 125.6, 128.6, 129.1, 129.3, 132.5, 133.7, 134.4, 138.0, 138.7, 146.7, 166.7 (CO). ESIMS m/z (%) 334.2 [M^+] (72). Anal. Calcd. For $\text{C}_{18}\text{H}_{14}\text{N}_4\text{OS}$ (334.39): C, 64.65; H, 4.22; N, 16.75; S, 9.59. Found: C, 64.58; H, 4.16; N, 16.67; S, 9.48.

Ethyl 2-amino-6-(thien-2-yl)-4-(indol-3-yl)pyridine-3-carboxylate (3). Yield: 72%, orange crystals: mp 193-195°C. IR (ν/cm^{-1}): 3445, 3374, 3333 (NH_2 and indole-NH), 3026 (Ar-H), 1711 (CO); ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 1.31-1.38 (t, J 24.4 Hz, 3H, CH_3), 4.24-4.29 (q, J 14.0, 7.2 Hz, 2H, CH_2), 7.23-8.22 (m, 9 H, 3 thienyl-H, 5 indole-H and pyridine-H), 8.55 (s, 2H, NH_2 vanished with D_2O), 12.57 (s, 1H, indole-NH); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ = 14.5 (CH_3), 61.8 (CH_2), 92.8, 110.4, 112.9, 113.1, 115.6, 118.9, 120.8, 121.6, 123.2, 125.7, 128.6, 132.4 133.0, 138.0, 138.6, 146.9, 176.9 (CO). ESIMS m/z (%) 363.2 [M^+] (87). Anal. Calcd. For $\text{C}_{20}\text{H}_{17}\text{N}_3\text{O}_2\text{S}$ (363.43): C, 66.10; H, 4.71; N, 11.56; S, 8.82. Found: C, 66.03; H, 4.64; N, 11.46; S, 8.70.

2-Amino-3-cyano-6-(thien-2-yl)-4-(indole-3-yl)pyridine (4). Yield: 68%, yellow crystals: mp 211-213 °C. IR (ν/cm^{-1}): 3350, 3210, 3121 (NH_2 and indole-NH), 3047 (Ar-H), 2217 (CN); ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 7.24-8.52 (m, 9 H, 3 thienyl-H, 5 indole-H and pyridine-H), 8.66 (s, 2H, NH_2 vanished by D_2O), 12.35 (s, 1H, indole-NH); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 98.2, 110.4, 111.4, 113.5, 116.4, 118.8, 119.2, 122.0, 123.0, 125.7, 128.6, 130.6 133.7, 136.6, 138.6, 142.7, 153.0. ESIMS m/z (%) 316.2 [M^+] (69). Anal. Calcd. for $\text{C}_{18}\text{H}_{12}\text{N}_4\text{S}$ (316.38): C, 68.33; H, 3.82; N, 17.71; S, 10.13. Found: C, 68.27; H, 3.76; N, 17.60; S, 10.03.

7-(Thien-2-yl)-5-(1*H*-indol-3-yl)-2-thioxo-2,3-dihydropyrido[2,3-*d*]pyrimidin-4(1*H*)-one (5). 6-Amino-2-thiouracil (0.28 g, 2 mmol) was mixed with chalcone **1** (0.50 g, 2 mmol) in glacial AcOH (10 mL) then refluxed for 15 h. Once cooled, the liquid was added to broken ice and the resulting solid filtered off and purified from acetic acid giving a brown powder. Yield 74%: mp 221-223 °C. IR (ν/cm^{-1}): 3390, 3307, 3176 (2 NH and indole-NH), 3026 (Ar-H), 1659 (CO); ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 6.65-8.39 (m, 9 H, 3 thienyl-H, 5 indole-H and pyridine-H), 11.00 (s, 1H, NH vanished by D_2O), 12.70 (s, 1H, indole-NH), 13.25 (s, 1H, NH vanished by D_2O); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 100.9, 110.9, 112.1, 116.1, 117.4, 118.0, 120.6, 121.3, 129.7, 130.6, 132.6, 135.5, 135.9, 146.3, 150.6, 155.6, 165.2 (C=O), 176.0 (C=S). ESIMS m/z (%) 376.3 (M^+ , 68%). Anal. Calcd. for $\text{C}_{19}\text{H}_{12}\text{N}_4\text{OS}_2$ (376.45): C, 60.62; H, 3.21; N, 14.88; S, 17.04. Found: C, 60.54; H, 3.13; N, 14.80; S, 16.94.

2-Hydrazinyl-7-(thien-2-yl)-5-(1*H*-indol-3-yl)pyrido[2,3-*d*]pyrimidin-4(3*H*)-one (6). A solution of 2-thioxopyridopyrimidine (0.75 g, 2 mmol) in ethanol (10 mL) was treated with hydrazine hydrate (80%, 4 mL) and kept at reflux for 20 h. Upon cooling, the precipitate was collected and purified from DMF. Yield 68%, white powder: mp 301-302 °C. IR (ν/cm^{-1}): 3410 and 3235 (NH and NH_2), 3180 (indole-NH), 1665 (CO); ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 6.88-8.15 (m, 9 H, 3 thienyl-H, 5 indole-H and pyridine-H), 8.21 (s, 2H, NH_2 vanished by D_2O), 12.40 (s, 1H, indole-NH vanished with D_2O), 13.00 (s, 1H, NH vanished by D_2O); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$): δ 101.8, 102.6, 106.2, 108.2, 127.7, 128.2, 129.4, 129.7, 130.9, 131.2, 133.8, 134.2, 150.6, 151.9, 158.3, 159.2,

167.5, 168.7, 168.9, 177.0. ESIMS m/z (%) 374.3 (M^+ , 59%). Anal. Calcd. for $C_{19}H_{14}N_6OS$ (374.42): C, 60.95; H, 3.77; N, 22.45; S, 8.56. Found: C, 60.89; H, 3.70; N, 22.40; S, 8.46.

General process for synthesizing 2-pyrazoloyl pyrido[2,3-d]pyrimidines (7 and 8).

2-hydrazinyl

pyridopyrimidine **6** (0.74 g, 2 mmol) was mixed with acetylacetone or ethyl acetoacetate (2 mmol) in ethanol and the mixture refluxed for 8 h. Upon cooling, the formed solid was collected, dried and recrystallized with 1,4-dioxane/ethanol.

2-(3,5-Dimethyl-1H-pyrazol-1-yl)-7-(thien-2-yl)-5-(1H-indol-3-yl)pyrido[2,3-d]pyrimidin-4(3H)-one (7). Yield: 82%, white crystals: mp 204–206 °C. IR (ν/cm^{-1}): 3385 (NH), 3148 (indole-NH), 2982 (Aliph-H), 1656 (CO); 1H NMR (400 MHz, DMSO- d_6): δ 1.70 (s, 3H, CH₃), 2.20 (s, 3H, CH₃), 6.60 (s, 1H, pyrazole-H₄), 7.24–8.22 (m, 9 H, 3 thienyl-H, 5 indole-H and pyridine-H), 12.32 (s, 1H, indole-NH), 13.12 (s, 1H, NH vanished by D₂O) ppm; ^{13}C NMR (100 MHz, DMSO- d_6): δ 14.1 (CH₃), 15.3 (CH₃), 108.3, 111.8, 113.9, 114.8, 116.6, 117.7, 118.9, 120.4, 122.3, 125.9, 127.4, 129.2, 130.8, 132.6, 134.4, 135.7, 146.4, 151.2, 152.9, 154.1, 155.8, 166.2 ppm. ESIMS m/z (%) 438.4 (M^+ , 62%). Anal. Calcd for $C_{24}H_{18}N_6OS$ (438.50): C, 65.74; H, 4.14; N, 19.17; S, 7.31. Found: C, 65.68; H, 4.08; N, 19.11; S, 7.21.

2-(3-Methyl-5-oxo-2-pyrazolin-1-yl)-7-(thien-2-yl)-5-(1H-indol-3-yl)pyrido[2,3-d]pyrimidin-4(3H)-one (8). Yield: 76%, white plates: mp 199–201 °C. IR (ν/cm^{-1}): 3305 (NH), 3151 (indole-NH), 3058 (Ar-H), 1675 (CO), 1645 (pyrazolone-CO); 1H NMR (400 MHz, DMSO- d_6): δ 2.28 (s, 3H, CH₃), 2.70 (s, 2H, pyrazoline-CH₂), 7.12–8.34 (m, 9 H, 3 thienyl-H, 5 indole-H and pyridine-H), 12.61 (s, 1H, indole-NH), 13.00 (s, 1H, NH vanished by D₂O) ppm; ^{13}C NMR (100 MHz, DMSO- d_6): δ = 17.5 (CH₃), 42.6 (CH₂), 107.9, 112.1, 114.2, 115.3, 117.8, 119.2, 121.0, 122.8, 126.1, 127.8, 129.0, 130.7, 132.5, 135.8, 145.8, 150.4, 152.8, 154.2, 155.9, 165.2, 167.4 ppm. ESIMS m/z (%) 440.3 (M^+ , 81%). Anal. Calcd for $C_{23}H_{16}N_6O_2S$ (440.48): C, 62.72; H, 3.66; N, 19.08; S, 7.28. Found: C, 62.66; H, 3.60; N, 19.00; S, 7.20.

1-Thiocarbamoyl-3-(thien-2-yl)-5-(1H-indol-3-yl)-2-pyrazoline (9). A solution of NaOH (4 mmol) in H₂O (1 mL) was introduced to chalcone **1** (0.50 g, 2 mmol) and thiosemicarbazide (2 mmol) dissolved in ethanol (10 mL). The mixture was subjected to reflux for 10 h, following which the product was transferred to crushed ice. The solid mass underwent filtration, drying, and crystallization with dioxane. Yield: 68%, yellow crystals: mp 231–232 °C. IR (ν/cm^{-1}): 3447, 3312 (NH₂), 3180 (NH), 3039 (Ar-H); 1H NMR (400 MHz, DMSO- d_6): δ = 3.68–3.76 (dd, J = 24.0, 10.0 Hz, H_A), 3.90–3.98 (dd, J = 20.0, 14.0 Hz, H_B), 6.29–6.35 (dd, J 14.0, 10.0 Hz, H_X), 7.12–8.28 (m, 8 H, 3 thienyl-H and 5 indole-H), 8.34 (s, 2H, NH₂), 11.59 (s, 1H, indole-NH) ppm; ^{13}C NMR (100 MHz, DMSO- d_6): δ = 49.1 (pyrazoline-CH₂), 66.8 (pyrazoline-CH), 111.6, 112.9, 113.1, 115.6, 120.9, 121.1, 122.5, 123.9, 124.4, 125.7, 128.6, 138.0, 141.3, 177.0 ppm. ESIMS m/z (%) 326.2 (M^+ , 81%). Anal. Calcd. for $C_{16}H_{14}N_4S_2$ (326.44): C, 58.87; H, 4.32; N, 17.16; S, 19.65. Found: C, 58.80; H, 4.25; N, 17.04; S, 19.53.

1-(4-Oxo-5H-thiazol-2-yl)-3-(thien-2-yl)-5-(1H-indol-3-yl)-2-pyrazoline (10). An equimolar mixture containing 1-thiocarbamoyl-2-pyrazoline **9** (0.65 g, 2 mmol), chloroacetic acid (0.18 g, 2 mmol), with anhydrous NaOAc (0.40 g) in glacial AcOH (20 mL) was refluxed for 10 h. The formed solid while cooling was crystallized with DMF. Yield: 68%, white crystals: mp 177–178 °C. IR (ν/cm^{-1}): 3247 (NH), 3051 (Ar-H), 1703 (CO); 1H NMR (400 MHz, DMSO- d_6): δ = 3.50–3.57 (dd, J = 20.0, 8.0 Hz, H_A), 3.89 (s, 2H, CH₂), 4.22–4.29 (dd, J = 16.0, 12.0 Hz, H_B), 6.18–6.23 (dd, J 12.0, 8.0 Hz, H_X), 7.17–8.57 (m, 8 H, 3 thienyl-H and 5 indole-H), 11.68 (s, 1H, indole-NH) ppm; ^{13}C NMR (100 MHz, DMSO- d_6): δ 48.2 (pyrazoline-CH₂), 60.4 (thiazole-CH₂), 66.5 (pyrazoline-CH), 112.4, 121.2, 122.5, 123.2, 124.6, 125.0, 128.5, 132.1, 134.8, 137.6, 141.1, 142.4, 148.3, 167.8 (CO) ppm. ESIMS m/z (%) 366.3 (M^+ , 72%). Anal. Calcd. for $C_{18}H_{14}N_4OS_2$ (366.46): C, 58.99; H, 3.85; N, 15.29; S, 17.50. Found: C, 58.92; H, 3.78; N, 15.20; S, 17.40.

1-(4-Phenylthiazol-2-yl)-3-(thien-2-yl)-5-(1*H*-indol-3-yl)-2-pyrazoline (11). Phenacyl bromide (0.39 g, 2 mmol) was mixed with 1-thiocarbamoyl-2-pyrzoline **9** (0.65 g, 2 mmol) in ethanol (25 mL) and refluxed for 6 h. The solid obtained upon cooling filtered off and crystallized in 1,4-dioxane. Yield: 64%, brown powder: mp 190-192 °C. IR (ν/cm^{-1}): 3151 (NH), 3051 (Ar-H); ^1H NMR (400 MHz, DMSO- d_6): δ = 3.91-3.98 (dd, J = 18.0, 10.0 Hz, H_A), 4.16-4.23 (dd, J 16.0, 12.0 Hz, H_B), 6.12-6.18 (dd, J 12.0, 10.0 Hz, H_X), 7.13-8.15 (m, 14 H, 5 Ar-H, 3-thienyl-H, 5-indole-H, and thiazole-H), 9.00 (s, 1H, thiazole-H), 11.70 (s, 1H, indole-NH) ppm; ^{13}C NMR (100 MHz, DMSO- d_6): δ = 48.2 (pyrazoline-CH₂), 60.1 (thiazole-CH₂), 66.5 (pyrazoline-CH), 101.8, 102.5, 106.3, 108.2, 127.8, 128.0, 128.7, 129.6, 129.9, 130.9, 131.1, 133.9, 134.2, 150.7, 151.8, 158.0, 159.1 ppm. ESIMS m/z (%) 429.1 (M⁺, 73%). Anal. Calcd. for C₂₄H₁₈N₄S₂ (426.56): C, 67.58; H, 4.25; N, 13.13; S, 15.03. Found: C, 67.51; H, 4.18; N, 13.02; S, 15.13.

General route for the synthesis of 1-arylidenethiazolyl-2-pyrazolines (12-14). The aromatic aldehydes (2 mmol), anhydrous AcONa (0.40 g), and compound **10** (0.73 g, 2 mmol) were refluxed in AcOH (10 mL) for 6 h. The solution was cooled before being added to broken ice. The solid was obtained through filtration, drying, and crystallization with dioxane, resulting in the formation of products **12-14**.

1-(5-Benzyliden-4-oxothiazol-2-yl)-3-(thien-2-yl)-5-(1*H*-indol-3-yl)-2-pyrazoline (12). Yield: 80%, orange powder: mp 230-231 °C. IR (ν/cm^{-1}): 3130 (NH), 3025 (Ar-H), 1711 (CO); ^1H NMR (400 MHz, DMSO- d_6): δ 3.31-3.39 (dd, J = 24.0, 10.0 Hz, H_A), 3.90-3.98 (dd, J 20.0, 12.0 Hz, H_B), 6.11-6.16 (dd, J 12.0, 10.0 Hz, H_X), 6.84-8.65 (m, 14 H, 5 Ar-H, 3 thienyl-H, 5 indole-H and =CH-), 11.69 (s, 1H, indole-NH) ppm; ^{13}C NMR (100 MHz, DMSO- d_6): δ 44.2 (pyrazoline-CH₂), 66.8 (pyrazoline-CH), 111.6, 112.4, 116.1, 116.5, 118.6, 119.3, 120.9, 121.1, 123.1, 125.8, 125.9, 127.8, 129.8, 130.0, 132.0, 136.6, 137.6, 152.4, 155.8, 160.3, 168.6 (CO) ppm. ESIMS m/z (%) 454.4 (M⁺, 62%). Anal. Calcd. for C₂₅H₁₈N₄OS₂ (454.57): C, 66.06; H, 3.99; N, 12.33; S, 14.11. Found: C, 66.00; H, 3.97; N, 12.25; S, 14.04.

1-(5-p-Chlorobenzyliden-4-oxothiazol-2-yl)-3-(thien-2-yl)-5-(1*H*-indol-3-yl)-2-pyrazoline (13). Yield: 69%, pale yellow crystals: mp 200-201 °C. IR (ν/cm^{-1}): 3130 (NH), 3068 (Ar-H), 1710 (CO); ^1H NMR (400 MHz, DMSO- d_6): δ 3.32-3.40 (dd, J 24.0, 10.0 Hz, H_A), 3.91-3.99 (dd, J 0.0, 12.0 Hz, H_B), 6.12-6.17 (dd, J 2.0, 10.0 Hz, H_X), 6.90-8.60 (m, 13 H, 4 Ar-H, 3 thienyl-H, 5 indole-H and =CH-), 11.71 (s, 1H, indole-NH) ppm; ESIMS m/z (%) 488.9 (M⁺, 73%) and 491.0 (M⁺+1, 24%). Anal. Calcd. for C₂₅H₁₇ClN₄OS₂ (489.01): C, 61.40; H, 3.50; Cl, 7.25; N, 11.46; S, 13.11. Found: C, 61.33; H, 3.42; Cl, 7.14; N, 11.40; S, 13.01.

1-(5-p-Bromobenzyliden-4-oxothiazol-2-yl)-3-(thien-2-yl)-5-(1*H*-indol-3-yl)-2-pyrazoline (14). Yield: 77%, red crystals: mp 220-221 °C. IR (ν/cm^{-1}): 3145 (NH), 3012 (Ar-H), 1722 (CO); ^1H NMR (400 MHz, DMSO- d_6): δ 3.46-3.50 (dd, J 15.0, 7.0 Hz, H_A), 3.92-3.95 (dd, J 10.0, 8.5 Hz, H_B), 6.34-6.36 (dd, J 8.5, 7.0 Hz, H_X), 6.90-8.60 (m, 13 H, 4 Ar-H, 3 thienyl-H, 5 indole-H and =CH-), 11.71 (s, 1H, indole-NH) ppm; ESIMS m/z (%) 533.2 (M⁺, 54%) and 534.3 (M⁺+1, 53%). Anal. Calcd. for C₂₅H₁₇BrN₄OS₂ (533.46): C, 56.29; H, 3.21; Br, 14.98; N, 10.50; S, 12.02. Found: C, 56.23; H, 3.14; Br, 14.87; N, 10.40; S, 12.18.

General method for the synthesis of azepines (15 and 16). A mixture of **1** (0.50 g, 2 mmol) and either o-phenylenediamine or o-aminophenol (2 mmol) in ethanol (10 mL) along with triethylamine (3-4 drops) was refluxed for 10 h. The solution was subjected to cooling at 0 °C and permitted to stand overnight. The solid product was filtered off, rinsed with water, and then purified in 1,4-dioxane.

4-(Thieny-2-yl)-2-(indol-3-yl)-2,3-dihydro-1*H*-benzo[*b*][1,4]diazepine (15). Yellow crystals: mp 223-225 °C. IR (ν/cm^{-1}): 3333 (NH), 3172 (indole-NH), 3074 (Ar-H) cm⁻¹. ^1H NMR (400 MHz, DMSO- d_6): δ = 3.13-3.20 (dd, J 16.0, 12.0 Hz, H_A), 3.81-3.88 (dd, J 15.2, 13.2 Hz, H_B), 4.25 (s, 1H, NH), 5.00-5.56 (dd, J 13.2, 12.0 Hz, H_Z), 7.24-8.56 (m, 12H, thienyl-H, indole-H, and benzodiazepine-H), 12.48 (s, 1H, indole-NH) ppm; ^{13}C NMR (100 MHz,

DMSO-*d*₆): δ = 18.5 (CH₂), 46.2 (CH), 60.2, 68.6, 116.2, 118.4, 120.9, 121.6, 122.04, 132.87, 133.6, 137.5, 137.8, 145.5, 150.4, 153.3, 155.2, 155.9, 158.0. ESIMS *m/z* (%) 343.2 [M⁺] (77). Anal.Calcd. for C₂₁H₁₇N₃S (343.44): C, 73.44; H, 4.99; N, 12.23; S, 9.34. Found: C, 73.39; H, 4.93; N, 12.13; S, 9.23.

4-(Thieny-2-yl)-2-(indol-3-yl)-2,3-dihydrobenzo[*b*][1,4]oxazepine (16). White solid: mp 230-232 °C. IR (*v*/cm⁻¹): 3148 (indole-NH), 3008 (Ar-H) cm⁻¹. ¹H NMR (400 MHz, DMSO-*d*₆): δ = 3.30-3.40 (dd, *J* = 24.0, 16.0 Hz, H_A), 4.10-4.19 (dd, *J* 20.0, 18.0 Hz, H_B), 5.70-5.78 (dd, *J* 18.0, 16.0 Hz, Hz), 7.20-8.60 (m, 12H, thienyl-H, indole-H, and benzodioxazepine-H), 12.32 (s, 1H. indole-NH) ppm; ¹³C NMR (100 MHz, DMSO-*d*₆): δ = 20.3 (CH₂), 48.9 (CH), 62.0, 70.2, 116.4, 118.5, 120.8, 121.5, 122.4, 123.8, 125.9, 132.7, 133.5, 137.4, 137.3, 145.0, 150.2, 153.0, 155.6, 155.8, 158.2. ESIMS *m/z* (%) 344.1 [M⁺] (70). Anal.Calcd. for C₂₁H₁₆N₂OS (344.43): C, 73.23; H, 4.68; N, 8.13; S, 9.31. Found: C, 73.16; H, 4.60; N, 8.02; S, 9.24.

Biological activity

In vitro antimicrobial assay procedure. The studied compounds **2-16** were dissolved in DMSO resulting in a 5% solution. The resulting solution was applied to saturate filter paper discs (Whatman No. 3 and 5 mm in diameter). The discs were set on the surface of solidified nutrient agar dishes seeded with the bacteria that were examined (*Gram positive; Bacillus cereus and Staphylococcus aureus, and Gram negative; Pseudomonas aruginose or Escherichia coli*) or Czapek Dox agar dishes seeded with the fungi that were investigated (*Geotrichum candidum, Candida albicans, Trichophyton rubrum, and Aspergillus flavus*). The diameter of inhibiting zones (mm) was measured at the end of the incubation period (24-48 h) at 37 °C for bacteria and 28 °C with fungi (4-7 days) [48]. Discs soaked with DMSO were employed as controls. *Chloramphenicol* and *Clotrimazole* were chosen as references. The biologically active compounds were subsequently diluted with DMSO to produce an array of concentrations with the goal to ascertain each compound's *MIC*. The *MIC* values were calculated as µg/mL **Tables 1** and **2** provide the antibacterial and antifungal activity data.

Acknowledgement

The authors extend their appreciation to the Researchers Supporting Project (RSPD2024R758), King Saud University, Riyadh, Saudi Arabia.

Funding Information

This work was funded by the Researchers Supporting Project (RSPD2024R758), King Saud University, Riyadh, Saudi Arabia.

Supplementary Material

The FTIR spectra, ¹H NMR and ¹³C NMR of the new compounds can be found in the supplementary material file.

References

1. Lou, X.; Yang, Y. Y. W. *J. Am. Chem. Soc.* **2021**, *28*, 11976-11981.
<https://doi.org/10.1021/jacs.1c07006>
2. Henry, G. D. *Tetrahedron* **2004**, *60*, 6043.
<https://doi.org/10.1016/j.tet.2004.04.043>
3. Brycki, B.; Małecka, I.; Koziróg, A.; Otlewska, A. *Molecules* **2017**, *22*, 130.
<https://doi.org/10.3390/molecules22010130>
4. Leal, B.; Afonso, I. F.; Rodrigues, C. R.; Abreu, P. A.; Garrett, R.; Pinheiro, L. C. S.; Azevedo, A. R.; Borges, J. C.; Vegi, P. F.; Santos, C. C. *Bioorg. Med. Chem.* **2008**, *16*, 8196-8204.
<https://doi.org/10.1016/j.bmc.2008.07.035>
5. Zav'yalova, V. K.; Zubarev, A. A.; Shestopalov, A. M. *Russ. Chem. Bull.* **2009**, *58*, 1939-1944.
<https://doi.org/10.1007/s11172-009-0265-2>
6. Bernardino, A. M. R.; da Silva Pinheiro, L. C.; Rodrigues, C. R.; Loureiro, N. I.; Castro, H. C.; Lanfredi-Rangel, A.; Sabatini-Lopes, J.; Borges, J. C.; Carvalho, J. M.; Romeiro, G. A. *Bioorg. Med. Chem.* **2006**, *14*, 5765-5777.
<https://doi.org/10.1016/j.bmc.2006.03.013>
7. Hayakawa, M.; Kaizawa, H.; Kawaguchi, K.; Ishikawa, N.; Koizumi, T.; Ohishi, T.; Yamano, M.; Okada, M.; Ohta, M.; Tsukamoto, S. *Bioorg. Med. Chem.* **2007**, *15*, 403-412.
<https://doi.org/10.1016/j.bmc.2006.09.047>
8. Truong, D. T.; Ho, K.; Nhi, H. T. Y.; Nguyen, V. H.; Dang, T. T.; Nguyen, M. T. *Sci. Rep.* **2024**, *28*, 12218.
<https://doi.org/10.1038/s41598-024-62743-3>
9. Janković, Đ. D.; Šestić, T. L.; Bekić, S. S.; Savić, M. P.; Ćelić, A. S.; Scholda, J.; Kopp, F.; Marinović, M. A.; Petri, E. T.; Ajduković, J. J. *J. Steroid Biochem. Mol. Biol.* **2024**, *16*, 106545.
<https://doi.org/10.1016/j.jsbmb.2024.106545>
10. Li, A.; Huang, K.; Pan, W.; Wu, Y.; Liang, Y.; Zhang, Z.; Wu, D.; Ma, L.; Gou, Y. *J. Med. Chem.* **2024**, *67*, 9091-9103.
<https://doi.org/10.1021/acs.jmedchem.4c00257>
11. Adib, M.; Peytam, F.; Rahamanian-Jazi, M.; Mohammadi Khanaposhtani, M.; Mahernia, S.; Bijanzadeh, H. R.; Jahani, M.; Imanparast, S.; Faramarzi, M. A.; Mahdavi, M. *New J. Chem.* **2018**, *42*, 17268-17278.
<https://doi.org/10.1039/C8NJ02495B>
12. Suresh, L.; Kumar, P. S. V.; Onkar, P.; Srinivas, L.; Pydisetty, Y.; Chandramouli, G. V. P. *Res. Chem. Intermed.* **2017**, *43*, 5433-5451.
<https://doi.org/10.1007/s11164-017-2938-z>
13. Podila, N.; Penddinti, N. K. Rudrapal, M. Rakshit, G.; Konidala, S. K.; Pulusu, V. S.; Bhandare, R. R.; Shaik, A. B. *Helijon* **2024**, *10*, e29390.
<https://doi.org/10.1016/j.helijon.2024.e29390>
14. Bharate, S. B.; Mahajan, T. R.; Gole, Y. R.; Nambiar, M.; Matan, T. T.; Kulkarni-Almeida, A.; Balachandran, S.; Junjappa, H.; Balakrishnan, A.; Vishwakarma, R. A. *Bioorganic Med. Chem.* **2008**, *16*, 7167-7176.
<https://doi.org/10.1016/j.bmc.2008.06.042>
15. Hamblin, J. N.; Angell, T. D. R.; Ballantine, S. P.; Cook, C. M.; Cooper, A. W. J.; Dawson, J.; Delves, C. J.; Jones, P. S.; Lindvall, M.; Lucas, F. S. *Bioorganic Med. Chem. Lett.* **2008**, *18*, 4237-4241.
<https://doi.org/10.1016/j.bmcl.2008.05.052>
16. Tok, F.; Erdoğan, Ö.; Çevik, Ö.; Koçyiğit-Kaymakçıoğlu, B. *Acta Chim. Slov.* **2022**, *69*, 293-303.
<https://doi.org/10.17344/acsi.2021.7119>
17. Chen, K.; Zhang, Y. L.; Fan, J.; Ma, X.; Qin, Y. J.; Zhu, H. L. *Eur. J. Med. Chem.* **2018**, *156*, 722-737.

- <https://doi.org/10.1016/j.ejmech.2018.07.044>
18. Sever, B.; Altintop, M. D.; Radwan, M. O.; Özdemir, A.; Otsuka, M.; Fujita, M.; Ciftci, H. I. *Eur. J. Med. Chem.* **2019**, *182*, 111648.
<https://doi.org/10.1016/j.ejmech.2019.111648>
19. Gomha, S. M.; Abdelaziz, M. R.; Kheder, N. A.; Abdel-Aziz, H. M.; Alterary, S.; Mabkhot, Y. N. *Chem. Cent. J.* **2017**, *11*, 105.
<https://doi.org/10.1186/s13065-017-0335-8>
20. Abdel-Wahab, B. F.; Abdel-Aziz, H. A.; Ahmed, E. M. *Eur. J. Med. Chem.* **2009**, *44*, 2632-2635.
<https://doi.org/10.1016/j.ejmech.2008.09.029>
21. Pola, S.; Banoth, K. K.; Sankaranarayanan, M.; Ummani, R.; Garlapati, A. *Med. Chem. Res.* **2020**, *29*, 1819-1835.
<https://doi.org/10.1007/s00044-020-02602-8>
22. Dawane, B. S.; Konda, S. G.; Mandawad, G. G.; Shaikh, B. M. *Eur. J. Med. Chem.* **2021**, *45*, 387-392.
<https://doi.org/10.1016/j.ejmech.2009.10.015>
23. Hassan, S. Y. *Molecules* **2013**, *18*, 2683-2711.
<https://doi.org/10.3390/molecules18032683>
24. Dhonnar, S. L.; Jagdale, B. S.; Adole, V. A.; Sadgir, N. V. *Mol. Divers.* **2023**, *27*, 2441-2452.
<https://doi.org/10.1007/s11030-022-10562-x>
25. Karad, S. C.; Purohit, V. B.; Thakor, P.; Thakkar, V. R.; Raval, D. K. *Eur. J. Med. Chem.* **2016**, *112*, 270-279.
<https://doi.org/10.1016/j.ejmech.2016.02.016>
26. Ramírez-Prada, J.; Robledo, S. M.; Vélez, I. D.; Crespo, M. D. P.; Quiroga, J.; Abonia, R.; Montoya, A.; Svetaz, L.; Zacchino, S.; Insuasty, B. *Eur. J. Med. Chem.* **2017**, *131*, 237-254.
<https://doi.org/10.1016/j.ejmech.2017.03.016>
27. Deng, H.; Yu, Z. Y.; Shi, G. Y.; Chen, M. J.; Tao, K.; Hou, T. P. *Chem. Biol. Drug Des.* **2012**, *79*, 279-789.
<https://doi.org/10.1111/j.1747-0285.2011.01308.x>
28. Ovais, S., Pushpalatha, H., Reddy, G. B., Rathore, P., Bashir, R., Yaseen, S. Javed, K. *Eur. J. Med. Chem.* **2014**, *80*, 209-217.
<https://doi.org/10.1016/j.ejmech.2014.04.046>
29. Mantzanidou, M., Pontiki, E., Hadjipavlou-Litina, D. *Molecules* **2021**, *26*, 3439.
<https://doi.org/10.3390/molecules26113439>
30. Vasudha, D.; Jagadeesh, A.; Mali, S. N.; Bhandare, R. R.; Shaik, A. B. *Chem. Phys. Impact* **2024**, *8*, 100500.
<https://doi.org/10.1016/j.chphi.2024.100500>
31. Abeed, A. A. O.; Abdel Jaleel, G. A.; Youssef, M. S. K. *Curr. Org. Synth.* **2019**, *16*, 921-930.
<https://doi.org/10.2174/1570179416666190703115133>
32. Asif, M.; Almehmadi, M. M.; Alsaiari, A.; Allahyani, M. *Curr. Org. Synth.* **2023**, *21*, 858-888.
<https://doi.org/10.2174/0115701794260444230925095804>
33. Ziarani, G. M.; Hasani, S.; Mohajer, F.; Varma, R. S.; Rafiee, F. *Top Curr. Chem.* **2022**, *380*, 24.
<https://doi.org/10.1007/s41061-022-00379-5>
34. Venugopala, K. N.; Habeebuddin, M.; Aldhubiab, B. E.; Asif, A. H. *Molecules* **2021**, *26*, 5235.
<https://doi.org/10.3390/molecules26175235>
35. Böttcher, C.; Chapman, A.; Fellermeier, F.; Choudhary, M.; Scheel, D.; Glawischnig, E. *Plant Physiol.* **2014**, *165*, 841-853.
<https://doi.org/10.1104/pp.114.235630>
36. Gallego-Yerga, L.; Chiliquinga, A. J.; Peláez, R. *Int. J. Mol. Sci.* **2023**, *24*, 11093.

<https://doi.org/10.3390/ijms241311093>

37. Chen, H.; Deng, S.; Albadari, N.; Yun, M. K.; Zhang, S.; Li, Y.; Ma, D.; Parke, D. N.; Yang, L.; Seagroves, T. N.; White, S. W.; Miller, D. D.; Li, W. *J. Med. Chem.* **2024**, *64*, 12049-12074.
<https://doi.org/10.1021/acs.jmedchem.1c00715>
38. Song, M.; Hua ,Y.; Liu Y.; Xiao, X.; Yu, H.; Deng, X. *Molecules* **2023**, *28*, 6325.
<https://doi.org/10.3390/molecules28176325>
39. Aboshouk , D. R.; Youssef, M. A.; Bekheit ,M. S.; Hamed A. R.; Grgis, A. S. *RSC Adv.* **2024**, *14*, 5690-5728.
<https://doi.org/10.1039/D3RA08962B>
40. Yadav, R.; Meena, D.; Singh, K.; Tyagi, R.; Yadav, Y.; Sagar, R. *RSC Adv.* **2023**, *13*, 21890-21925.
<https://doi.org/10.1039/D3RA03862A>
41. Rupa, S. A.; Moni, M. R.; Patwary, M. A. M.; Mahmud, M. M.; Haque, M. A.; Uddin J, Abedin, S. M. T. *Molecules*, **2022**, *27*, 1656.
<https://doi.org/10.3390/molecules27051656>
42. Hassneen, H. M.; Abdallah, T. A. *Molecules*, **2003**, *8*, 333-341.
<https://doi.org/10.3390/80300333>
43. Mabkhot , Y. N.; Alatibi, F.; El-Sayed. N. N.; Al-Showiman, S.; Kheder, N. A.; Wadood, A.; Rauf A.; Bawazeer S.; Hadda, T. B. *Molecules* **2016**, *21*, 222.
<https://doi.org/10.3390/molecules21020222>
44. Mete, E.; Gul, H. I.; Kazaz, C. *Molecules* **2007**, *12*, 2579-2588.
<https://doi.org/10.3390/12122579>
45. Akulov, A. A.; Varaksin, M. V.; Charushin, V. N.; Chupakhin, O. N. *ACS Omega* **2019**, *10*, 825-834.
<https://doi.org/10.1021/acsomega.8b02916>
46. Meleddu, R.; Petrikaite, V.; Distinto, S.; Arridu, A.; Angius, R.; Serusi, L.; Škarnulytė, L.; Endriulaitytė, U.; Paškevičiu, Te. M.; Cottiglia, F.; Gaspari, M.; Taverna, D.; Deplano, S.; Fois, B.; Maccioni, E. *ACS Med. Chem. Lett.* **2018**, *10*, 571.
<https://doi.org/10.1021/acsmedchemlett.8b00596>
47. Tsukerman, S. V.; Nikitchenko, V. M.; Bugai, A. I.; Lavrushine, V. F. *Chem. Heterocycl. Comp.* **1969**, *5*, 268-272.
<https://doi.org/10.1007/BF00475844>
48. Kwon-Chung, K. J.; Bennett J. E. *Medical Mycology*, Lea & Febiger, Philadelphia, PA **1992**, 866.

ACCEPTED COPY. DO NOT DELETE THIS MESSAGE.

This manuscript has been corrected by the author, checked by the Scientific Editor, and contains all necessary changes.

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)

By checking the following box, the senior author acknowledges that they and the other authors have read, understood, and complied with the Instructions to Authors (ItA).

