

Supplementary Material

A facile and highly efficient one-step N-alkylation of thiazolidine-2,4-dione


Virgilio D. Ebajo, Jr.^{a*} and Glenn V. Alea^a

^a*Department of Chemistry, De La Salle University, 2401 Taft Avenue, Manila 0922 Philippines*
Email: virgilio.ebajo@dlsu.edu.ph


Table of Contents

¹ H-NMR data of 2	S2
¹ H-NMR data of 3	S3
¹ H-NMR data of 4	S4
¹ H-NMR data of 5	S5
¹ H-NMR data of 7	S6


¹H-NMR spectrum (300 MHz, DMSO-d₆) of 3-butylthiazolidine-2,4-dione (**2**).




^1H -NMR spectrum (300 MHz, DMSO- d_6) of 3-hexylthiazolidine-2,4-dione (**3**).



^1H -NMR spectrum (300 MHz, DMSO-d_6) of 3-octylthiazolidine-2,4-dione (**4**).



^1H -NMR spectrum (300 MHz, DMSO- d_6) of 3-decylthiazolidine-2,4-dione (**5**).



^1H -NMR spectrum (300 MHz, DMSO-d_6) of 3,3'-(propane-1,3-diyl)bis(thiazolidine-2,4-dione) (**7**).

