

Electrospray ionisation mass spectrometric studies of *N*-substituted 10-(aminosulfonyl)bornyl acrylate derivatives

Lulama P. Mciteka,^{a,b} Kevin A. Lobb,^a Marietjie A. Stander,^c and Perry T. Kaye^{a*}

^aDepartment of Chemistry, Rhodes University, Makhanda, South Africa

^bDepartment of Chemistry, University of the Western Cape, Bellville, Cape Town

^cCentral Analytical Facilities, Stellenbosch University, Stellenbosch, South Africa

Email: P.Kaye@ru.ac.za

Received mm-dd-yyyy


Accepted mm-dd-yyyy

Published on line mm-dd-yyyy

Dates to be inserted by editorial office

Abstract

Fragmentation patterns in the positive-ion electrospray ionization (ESI) mass spectra of representative *N*-substituted 10-(aminosulfonyl)bornyl acrylate derivatives have been explored using high-resolution methods. Evidence is presented for the involvement of common fragmentation pathways.



Keywords: Mass spectrometry, bornyl derivatives, sulfonamides, electrospray ionisation, fragmentation pathways


Introduction

The naturally occurring terpenoid borneol has found use in traditional medicine, and a number of bornyl esters have been shown to exhibit a range of pharmacological activities.¹ Bambagiotti *et al.*² have identified diagnostic fragment ions in the electron impact (EI) mass spectra of various bornyl esters, and Selva *et al.*³ have used negative ion chemical ionization (CI) mass spectrometry (MS) to explore the fragmentation of a variety of bornyl esters as well. We have previously studied⁴ fragmentation patterns in the EI mass spectra of selected chiral silyl enol ethers, including a bornyl ester derivative. Examples of applications of MS methods focusing on the analysis of bornyl systems in natural products include a gas chromatography-EIMS study⁵ of *Alpinia mutica* fruit oils and efficient differentiation of *C. camphora* chemotypes⁶ using Desorption Atmospheric Pressure Chemical Ionization (DAPCI-MS).

We have previously reported⁷ the synthesis of a series of optically pure camphor-10-sulfonamide-derived bornyl alcohol derivatives (**3**) (Scheme 1) as chiral auxiliaries, and the use of the subsequently derived acryloyl esters (**4**) as synthons in diastereoselective Baylis-Hillman reactions. To our knowledge, no Electrospray Ionisation (ESI) MS studies of such compounds have been undertaken and, in this communication, we now report our investigation of fragmentation patterns in the positive-ion ESI MS spectra of the representative bornyl acrylate derivatives **4a-c** and **5a**.

Results and Discussion

Access to the acryloyl esters **4a-c** was achieved in three steps as outlined in Scheme 1: i) formation of the camphor-10-sulfonamides **2a-c** from D-camphor-10-sulfonyl chloride **1**; ii) diastereoselective reduction to the epimeric bornyl alcohol derivatives **3a-c**, with clear predominance of the 2-*exo*-hydroxy analogues; and iii) reaction with acryloyl chloride to afford the esters **4a-c**, accompanied, in some cases, by the hydrochlorinated derivative (*i.e.*, **5a** and **5b**). Access to the acryloyl ester **4a** has also been achieved *via* direct generation of the 3-chloropropanoyl ester **5a**, followed by base-catalysed dehydrochlorination – as shown in Scheme 1. We now report on the results of a detailed exploration of the mass spectrometric fragmentation pathways exhibited by the representative compounds **4a-c** and **5a**.



Scheme 1. Preparation of bornyl acrylate derivatives.⁷ *Reagents and conditions:* i) DMAP, CH₃CN, 0 °C; ii) NaBH₄, EtOH-H₂O, 0 °C; iii) Al₂O₃, CH₂=CHCOCl; iv) Al₂O₃, ClCH₂CH₂COCl; v) Et₃N, N₂.

Positive-ion ESI mass spectra of the representative system, 2-*exo*-acryloyloxy-*N*-benzylbornane-10-sulfonamide **4a** (Figure 1), reveal a “base peak” at m/z 306.1528 and peaks at m/z 378.1743, 395.2002 and 400.1557 corresponding, respectively, to the pseudo-molecular ion **4a-H**⁺, the ammonio-molecular ion **4a-NH**₄⁺ and the sodio-molecular ion **4a-Na**⁺.






Figure 1. Positive-ion ESI MS spectrum of compound **4a**.

LC-MS/MS analyses indicate that the pseudo-molecular ion **4a-H⁺** and the ammonio-molecular ion **4a-NH₄⁺** exhibit parallel, but independent fragmentation pathways, i.e., independent in the sense that loss of NH₃ from **4a-NH₄⁺** to form **4a-H⁺** is not evident (Scheme 2).



Scheme 2. Significant ESI MS fragmentation series of the pseudo-molecular ion **4a-H⁺** and the ammonio-molecular ion **4a-NH₄⁺**; italicised data reflect the calculated *m/z* values.


In their ESI MS research on sulfonamides and tetracyclines, Tso and Aga⁸ attribute formation of an ammonio-molecular ion [M+NH₄⁺] to “gas-phase chemical ionisation induced by ammonium ions”. Subsequent loss of NH₃ then leads to a pseudo-molecular ion. In our study, however, the MS/MS spectrum of the ammonio-molecular ion **4a-NH₄⁺** reveals no evidence of the formation of the pseudo-molecular ion at *m/z* 378. On the other hand, the species corresponding to the base peak at *m/z* 306.1536 originates from both **4a-NH₄⁺** and **4a-H⁺**, and MS/MS analysis of this species (*m/z* 306) reveals further fragmentation to the species responsible for the peaks at *m/z* 135, 107 and 91 (Scheme 3).



Scheme 3. ESI MS fragmentation pathways of the ammonio-molecular ion **4a-NH₄⁺**; italicised data reflect the calculated *m/z* values.


Attention was initially focused on formation of the fragments corresponding to six significant peaks in the positive-ion ESI MS spectra of the ammonio-molecular ion **4a-NH₄⁺** (*m/z* 395), is illustrated in Scheme 3. Mechanistic rationalisations of the formation of these fragments are provided in Schemes 4 and 5.

As illustrated in Scheme 4, cleavage of the bornyl-oxygen bond and concomitant transfer of the 3-*endo* proton to a sulfonyl oxygen atom leads to fragment **4aI**, which is responsible for the base peak at *m/z* 306, and loss of ammonium acrylate from the ammonio-molecular ion **4a-NH₄⁺** (or, in the case of the pseudo-molecular ion **4a-H⁺**, loss of acrylic acid). Subsequent extrusion of SO₂ then leads to fragment **4aII** (*m/z* 242). The loss of SO₂ in the positive-ion mass spectra of sulfonamides has been reported by Sun *et al.*⁹ and by Niessen and Correa.¹⁰ The odd *m/z* value of the fragment with *m/z* 225 is consistent with an even-electron cation containing two nitrogen atoms and with the putative structure **4aIII**. Formation of this fragment is attributed to the tandem loss of three neutral molecules: i) ethylene, *via* intramolecular rearrangement; ii) sulfur dioxide, *via* precedented^{9,10} extrusion from a sulfonamide moiety; and iii) benzene from the benzylamine moiety. The presence of the same fragment (**4aIII**) in the MS/MS spectrum of the pseudo-molecular ion **4a-H⁺** suggests the possibility of a collision-induced dissociation³ cascade initiated by impact of NH₃ with the pseudo-molecular ion **4aI**.⁸




Scheme 4. Formation of the “base peak” fragment **4al** and the fragment ions **4all** and **4alll** from the ammonio-molecular ion **4a-NH₄⁺**; italicised data reflect the calculated *m/z* values.

The proposed pathways from the base-peak fragment **4al** (*m/z* 306) to the three, even-electron cationic fragments, **4alV**, **4aV** and **4aVI**, which are outlined in Scheme 5, are consistent with MS/MS data for **4al**. Loss of benzylaminosulfonic acid from **4al** leads, following sequential rearrangement of a primary carbocation (*via* a hydride shift) and a secondary carbocation (*via* Wagner-Meerwein rearrangement) to the even more stable tertiary allylic carbocation **4alV** (*m/z* 135). Tandem loss of benzylaminosulfonic acid and ethylene (*via* a *retro*-Diels Alder reaction) accounts for the resonance-stabilised secondary allylic cation **4aV** (*m/z* 107). Finally, cleavage of the *N*-benzylic bond in **4al** (fission b) affords a neutral camphor-10-sulfonamide tautomer and the observed aromatic tropylium cation **4aVI** at *m/z* 91.



Scheme 5. Generation of the even-electron cationic fragments **4aIV**, **4aV** and **4aVI** from the base-peak fragment **4a-I**; italicised data reflect the calculated m/z values.

The mass spectrum of the hydrochlorinated analogue **5a** of compound **4a** reveals the corresponding **M-H⁺** and **M-NH₄⁺** peaks at m/z 414 and 431, respectively. Fragments in common with those observed for compound **4a**, are evidenced by the ions at m/z 306, 242 and 225. The **M-H⁺** and **M-NH₄⁺** peaks for the *N*-(3-chlorophenyl) derivative **4b** are evident at m/z 398 and 415, respectively. Fragmentation of the latter, ammonio-molecular ion **4b-NH₄⁺** affords a fragment ($m/z = 326.0982$) which clearly contains a Cl^{35} atom, with the Cl^{37} isotopologue having $m/z = 328.0959$; this fragment is responsible for the base peak and corresponds to the base-peak fragments at m/z 306 in the spectra of the analogues **4a** and **5a**. The mass spectrum of 2-*exo*-(acryloyloxy)-*N*-(*t*-butyl)bornane-10-sulfonamide **4c** reveals clear pseudo-, ammonio- and sodio-molecular ion peaks at m/z 344.1892, 361.2157 and 366.1711, respectively. Formation of the sulfonamide **4cVII** (m/z 288.1269) is attributed to loss of isobutylene and NH_3 from the ammonio-molecular ion **4c-NH₄⁺** via “*fission b*” in Scheme 6, although similar loss of isobutylene alone from the pseudo-molecular ion would have the same effect. The base peak (m/z 216.1059) corresponds to the sulfonamide fragment **4cVIII** formed via similar elimination of isobutylene from the intermediate **4cI** (m/z 272.1681). While the latter fragment could equally well arise from the loss of acrylic acid from the pseudo-molecular ion, the MS/MS data obtained for the formation of analogous *N*-benzyl fragment **4aVIII** clearly implicates the corresponding ammonio-molecular ion **4c-NH₄⁺**.



Scheme 6. Selected fragmentation pathways from the ammonio-molecular ion **4c-NH₄⁺**; italicised data reflect the calculated *m/z* values.

Conclusions

High-resolution ESI MS and LC-MS/MS studies have provided insights into competing parallel, common and compound-specific fragmentation pathways in a series of representative *N*-substituted 10-(aminosulfonyl)-bornyl acrylate derivatives. Fragmentation pathways have been proposed with the aid of the MS/MS data and provide evidence of the involvement, *inter alia*, of *retro*-Diels-Alder, bicyclic terpenoid carbocation rearrangement and SO₂-extrusion processes.

Experimental Section

General. The synthesis and characterisation of compounds **4a-c** and **5a** have been reported previously.⁷ In an alternative approach to obtain a purer sample of the parent system **4a**, the diastereomeric alcohols **3a** were acylated with 3-chloropropanoyl chloride, followed by Et₃N-mediated dehydrochlorination and chromatographic (HPLC) purification as detailed below.

Neutral Al₂O₃ (0.244 g, 2.4 mmol) was added to the mixture of the diastereomeric alcohols **3a** (0.50 g, 1.6 mmol), followed by 3-chloropropanoyl chloride (0.38 g, 3 mmol). The resulting dispersion was shaken, sealed and kept unstirred at r.t. for 72 h. The mixture was then taken up in CHCl₃ (3 x 1 mL), filtered and the filtrate dried over anhydrous MgSO₄. The solvent was removed *in vacuo* to give an oil which was then treated with triethylamine (0.46 g, 3.1 mmol). The resulting mixture was stored under N₂ and stirred at 25 °C for 1 h and the taken up in EtOAc (5 mL); the organic solution was washed with brine and dried over anhydrous MgSO₄. Solvent was removed *in vacuo* to give an oil which was chromatographed [HPLC; elution with hexane-EtOAc (8: 2)] to afford four fractions.

Fraction 1. 2-Exo-acryloyloxy-N-(benzyl)bornane-10-sulfonamide 4a as white crystals (0.39 g, 49 %); δ_{H} (400 MHz; CDCl_3) 0.82 and 0.98 (6H, 2 x s, 8- and 9-Me), 1.23-2.03 (7H, series of multiplets, 3-, 5- and 6- CH_2 and 4-H), 2.81 and 3.44 (2H, 2 x d, 10- CH_2), 4.25 (2H, d, PhCH_2), 4.47 (1H, br s, NH), 5.04 (1H, d, 2-H) and 5.79 (1H, d, J 10.4 Hz, 3'-H), 6.09 (1H, dd, J 10.2 and 17.4 Hz, 2'-H) and 6.33 (1H, d, 3'-H_z) and 7.33 (5H, overlapping signals, ArH); δ_{C} (100 MHz; CDCl_3) 19.9 and 20.3 (C-8 and C-9), 27.1, 29.8 and 39.5 (C-3, C-5 and C-6), 44.4 (C-4), 47.2 (C-11), 49.2 and 49.4 (C-1 and C-7), 51.9 (C-10), 77.9 (C-2), 128.9 (C-2'), 130.3 (C-3'), 127.97, 128.0, 128.8 and 137.0 (ArC) and 164.8 (C=O); HRMS: m/z Calc. for $\text{C}_{20}\text{H}_{28}\text{NO}_4\text{S}$ (MH^+) 378.1734, Found 378.1743, and for $\text{C}_{20}\text{H}_{31}\text{N}_2\text{O}_4\text{S}$ (MNH_4^+) 395.2001, Found 395.2002.

Positive-ion high-resolution electrospray ionisation MS and LC-MS/MS data were recorded on a Waters Synapt G2 spectrometer (Capillary: 3 kV; Cone: 15 V; Calibration: sodium formate; Lock spray: leucine enkephalin) by the Central Analytical Laboratory at the University of Stellenbosch, South Africa. Relevant mass spectrometric data and instrumental conditions are available in the supporting information.

Acknowledgements

The authors thank the National Research Foundation (NRF), Rhodes University and the University of the Western Cape for generous bursary and/or financial support. This work is based on research supported by the National Research Foundation. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors, and, therefore, the NRF do not accept any liability in regard thereto. We also thank Prof Denzil Beukes (University of the Western Cape) for access to HPLC facilities.

Supplementary Material

ESI MS data for the representative compounds, and NMR data for compound **4a** can be found in the Supplementary Material accompanying this paper.

References

1. Silva, A. T. M.; Pereira, V. V.; de Almeida, L. T. G.; Ruiz, A. L. T. G.; de Carvalho, J. E.; Dias, D. F.; de Moreira, M. E. C.; Silva, R. R.; Duarte, L. P. *Rev Virtual Quim.* **2016**, *8*, 1020-1031.
<https://doi.org/10.5935/1984-6835.20160073>
2. Massimo, B. A.; Coran, S. A.; Giannellini, V.; Moneti, G.; Vincieri, F. F.; Selva, A.; Traldi, P. *Biomed. Mass Spectrom.* **1981**, *8*:343-346.
<https://doi.org/10.1002/bms.1200080805>
3. Selva, A.; Traldi, P.; Massimo, B. A. *Org. Mass Spectrom.* **1984**, *19*, 230-232.
<https://doi.org/10.1002/oms.1210190506>
4. Evans, M. D.; Kaye, P. T.; Learmonth, R. A. *J. Chem Res.* **1994**, *8*, 314-315.
5. Sirat, H. M.; Khalid, N. F. M.; Jani, N. A.; Basar, N. J. *Essent. Oil Res.* **2009**, *21*, 457-458.
<https://doi.org/10.1080/10412905.2009.9700217>

6. Guo, X.; Cui, M.; Deng, M.; Liu, X.; Huang, X.; Zhang, X.; Luo, L. *Sci Rep*, **2017**, 7. doi:10.1038/srep46579(2017).
<https://doi.org/10.1038/srep46579>
7. Mciteka, L. P.; Lobb, K. A.; Kaye, P. T. *Arkivoc* **2016**, (v), 151-163.
<https://doi.org/10.3998/ark.5550190.p009.737>
8. Tso, J.; Aga, D. S. *Anal. Chem.* **2011**, 83, 269-277.
<https://doi.org/10.1021/ac102391k>
9. Sun, M.; Dai, W.; Liu, D. Q. *J. Mass Spectrom.* **2008**, 43, 383-393.
<https://doi.org/10.1002/jms.1335>
10. Niessen, W. M. A.; Correa, C.; Ricardo, A. *Interpretation of MS/MS Mass Spectra of Drugs and Pesticides*, 1st Edn.; John Wiley & Sons; 2017.
<https://doi.org/10.1002/9781119294269>

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)