

Phenolic compounds from the rhizomes of *Zingiber pellitum*

Nguyen Phuong Hanh^a, Nguyen Thi Thu Minh^b, Nguyen Thu Uyen^b, Do Hoang Giang^b, Nguyen Quoc Binh^c, and Nguyen Tien Dat^{b,*}

^a Institute of Ecology and Biological Resources, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet Road, Cau Giay, Hanoi, Vietnam

^b Centre for High Technology Research and Development, VAST, 18-Hoang Quoc Viet, Cau Giay, Hanoi, Vietnam

^c Vietnam National Museum of Nature, VAST, 18-Hoang Quoc Viet, Cau Giay, Ha Noi, Vietnam

Email: ntdat@chtd.vast.vn

Received 12-01-2024

Accepted 02-10-2025


Published on line 02-19-2025

Abstract

Six new phenolic compounds have been isolated from the rhizome of *Zingiber pellitum*, including two new diarylheptanoids, zingiberpyrans A and B, rhamnocitrin-3-*O*-rhamnoside, meranzin hydrate, vanillin and (+)- α -viniferin. Their structures were elucidated by analyzing NMR, HRESIMS, and CD spectral evidences combining with the published data. Rhamnocitrin-3-*O*-rhamnoside (**3**) exhibited the strongest inhibition of nitric oxide production in LPS-stimulated RAW264.7 cells (IC₅₀ 49.6 μ M) while zingiberpyran A (**1**) was less active (IC₅₀ 71.0 μ M).



Zingiber pellitum Gagnep.



Keywords: *Zingiber pellitum*, diarylheptanoids, zingiberpyrans A, zingiberpyrans B, anti-inflammation

Introduction

Zingiber pellitum Gapnep. is an endemic species of Vietnam and is found in several regions of this country¹. It is an herbaceous plant, reaching a height of 0.8-1.2 meters, with densely long hair covering the entire plant and a tuberous rhizome. This species is distinguished by its unique inflorescence type, producing terminal flowers on leafy stems¹. Like other *Zingiber* species, *Z. pellitum* has a long history of use in traditional medicine to treat ailments such as cold, fever, and cough². A number of compounds have been isolated from different *Zingiber* species, such as diarylheptanoids, gingerols, flavonoids and terpenoids, exhibiting a wide spectrum of bioactivities, such as anti-inflammatory, antioxidant, anti-microbial and anti-cancer². However, previous reports on the chemical composition and biological activity of *Z. pellitum* predominantly focused on essential oil composition³⁻⁵. The present study, therefore, represents the first to report the isolation of six phenolic compounds (Figure 1), including two new diarylheptanoids, from *Z. pellitum* rhizomes, with their chemical structures determined through spectral analysis. Furthermore, the antioxidant and anti-inflammatory activities of the isolated compounds were also reported in this research.




Figure 1. Structures of the compounds isolated from *Z. pellitum* rhizomes.

Results and Discussion

By using various chromatographic techniques, the methanol extract of *Z. pellitum* rhizomes was separated to obtain two new diarylheptanoids **1** and **2**, and four known compounds, rhamnocitrin-3-*O*-rhamnoside (**3**),⁶ meranzin hydrate (**4**),⁷ vanillin (**5**),⁸ and (+)- α -viniferin (**6**).⁹

Compound **1** was obtained as an amorphous pale-yellow powder with the molecular formula $C_{20}H_{24}O_7$, which was confirmed by the sodium adduct at m/z 399.1379 $[M + Na]^+$ from the HRESIMS (calcd. for $C_{20}H_{24}NaO_7$, 399.1420). The 1H -NMR spectrum of **1** provided the information of 7 aromatic protons, therein 4 protons at δ_H 7.03 (2H, d, $J = 8.4$ Hz, H-2''/6'') and 6.70 (2H, d, $J = 8.4$ Hz, H-3''/5'') suggesting the presence of an AA'BB' spin system, and 3 proton signals at δ_H 6.81 (1H, d, $J = 8.4$ Hz, H-5'), 6.94 (1H, d, $J = 1.8, 7.8$ Hz, H-6'), 7.10 (1H, d, $J = 1.8$ Hz, H-2') were considered to belong to an ABX system (Table 1). The ^{13}C -NMR and HSQC demonstrated the presence of 20 carbon signals, regarding 12 signals belonging to two sets of aromatic systems, five oxymethines at δ_C 69.2, 73.7, 73.9, 75.8, and 79.4, one methoxy group at δ_C 56.5 and two methylene sp^3 carbons δ_C 35.4 and 39.8 (Table 2). In the COSY spectrum, the H-H spin-spin interactions were observed between H-2 (δ_H 4.39) \leftrightarrow H-3 (δ_H 3.53) \leftrightarrow H-4 (δ_H 4.17) \leftrightarrow H-5 (δ_H 1.77, 2.02) \leftrightarrow H-6 (δ_H 3.77) \leftrightarrow H-7 (δ_H 3.61) \leftrightarrow H-8 (δ_H 2.67, 2.83), which allowed for the identification of a continuous bond chain from C-2 to C-8 (Figure 2). The HMBC experiment showed the couplings from H-2 (δ_H 4.39) to C-1' (δ_C 133.5), C-2' (δ_C 112.7), C-6' (δ_C 122.0) and C-4 (δ_C 69.3) and interaction from H-8 (δ_H 2.67, 2.83) to C-1'' (δ_C 131.1), C-2''/6'' (δ_C 131.4) và C-6 (δ_C 73.9) allowing the identification of compound **1** as a diarylheptanoid with two benzene rings at C-2 and C-8 positions. Besides, HMBC couplings from H-2 (δ_H 4.39) to C-6 (δ_C 73.9) confirmed the presence of a cyclic pyran ring (Figure 2). The location of the methoxy group was assigned at C-3' based on the HMBC correlation from the methoxy signal (δ_H 3.90) to C-3' (δ_C 147.5).

Table 1. 1H NMR data (mult., J in Hz) of **1** and **2**

No	1 (in CD_3OD)	2 (in CD_3OD)	2 (in $DMSO-d_6$)
2	4.40 (1H, d, 10.2)	5.09 (1H, d, 6.0)	4.91 (1H, d, 4.8)
3	3.55 (1H, dd, 10.2, 3.0)	3.31 (1H, overlapped)	3.09 (1H, s)
4	4.17 (1H, q-like, 3.0)	4.68 (1H, q-like, 3.6)	4.44 (1H, q-like, 3.0)
5	2.05 (1H, ddd, 13.8, 12.0, 2.4)	2.04 (1H, ddd, 13.8, 7.8, 6.0)	1.84 (1H, m)
	1.79 (1H, ddd, 13.8, 3.6, 2.4)	1.82 (1H, m)	1.78 (1H, m)
6	3.78 (1H, dt, 12.0, 3.0)	3.32 (1H, overlapped)	3.20 (1H, s)
7	3.61 (1H, td, 7.2, 3.6)	3.46 (1H, m, H-7),	3.35 (1H, m, H-7),
8	2.85 (1H, dd, 13.2, 6.6)	2.60 (1H, dd, 13.8, 4.)	2.50 (1H, dd, 13.8, 4.2)
	2.69 (1H, dd, 13.2, 7.2)	2.53 (1H, dd, 13.8, 9.0)	2.39 (1H, dd, 13.8, 9.0)
2'	7.10 (1H, d, 1.8)	6.97 (1H, d, 1.8),	6.89 (1H, d, 1.2),
5'	6.81 (1H, d, 8.4)	6.82 (1H, d, 8.4)	6.75 (1H, d, 8.4)
6'	6.94 (1H, dd, 1.8, 7.8)	6.85 (1H, d, 8.4, 1.8)	6.72 (1H, d, 8.4, 1.2)
2''-6''	7.03 (2H, d, 8.4)	6.97 (2H, d, 8.4)	6.93 (2H, d, 8.4)
3''-5''	6.70 (2H, d, 8.4)	6.68 (2H, d, 8.4)	6.62 (2H, d, 8.4)
OCH ₃	3.90 (3H, br s)	3.85 (3H, br s)	3.74 (3H, br s)

This NMR data were almost identical to those of hedycoropyran A and B [11], except for the small difference in the pyran ring. The large coupling constant between H-2 and H-3 ($J_{2,3} = 10.2$ Hz) indicated that H-2 and H-3

were in di-axial relationship, while H-4 was in an equatorial orientation due to the small coupling constant with H-3 ($J_{3,4} = 3.0$ Hz).¹⁰ In the NOESY spectrum of **1**, an evident cross-peak between H-2 and H-6 confirmed the axial-orientation of H-6. For the absolute configuration identification, an ECD experiment was recorded. The 2*R*-configuration was determined based on the negative Cotton effect at 225 nm.^{11,12} Consequently, (3*S*,4*S*,6*R*) absolute configurations were assigned. However, the configuration of C-6 could not be determined using the present spectroscopic data. Thus, compound **1** was determined to be (2*R*,3*S*,4*S*,6*R*)-6-(-1-hydroxy-2-(4-hydroxyphenyl)ethyl)-2-(4-hydroxy-3-methoxyphenyl)tetrahydro-2*H*-pyran-3,4-diol, for which the name zingiberpyran A is proposed.




Figure 2. Key COSY (—), HMBC (→) and NOESY (-----) correlations of compounds **1** and **2**.

Compound **2** was obtained as an amorphous pale-yellow powder. Its HRESIMS showed a pseudomolecular ion peak at m/z 741.2852 $[M + Na]^+$ compatible with the molecular formula $C_{40}H_{46}O_{12}$ of **2**. The NMR data of **2** were similar to those of **1** with two aromatic AA'BB' and ABX systems, five aliphatic methines, two methylenes and one methoxy group. The detailed 2D NMR analysis of **2** (COSY, NOESY, HSQC and HMBC) allowed to assign the 1H and ^{13}C NMR data as in Table 1. The appearance of the upfield-shifted resonance δ_C 58.4 (C-3) instead of δ_C 73.7 in **1** suggested that C-3 was not hydroxylated. Comparing with the NMR data of diarylheptanoids previously reported [13, 14], this chemical shift is compatible with the C–C connection. In combination with the molecular formula $C_{40}H_{46}O_{12}$ as deduced from the HRESIMS experiment, two identical moieties, A and B, were proposed and linked via a C-3(A)—C-3(B) bridge. This connection was also supported by HMBC analysis which revealed a clear cross-peak between the proton at δ_H 3.31 (H-3) and the “itself” carbon atom at δ_C 58.4 (C-3) (Figure 2 and S19). It is noted that HMBC experiment gives heteronuclear long-range proton–carbon correlations but not direct one-bond correlations. Thus, the coupling from the proton at δ_H 3.31 to the carbon atom at δ_C 58.4 was assigned for the correlation from H-3 (A) to C-3(B) and vice versa.

Table 2. ^{13}C NMR data of **1** and **2**

No	1 (in CD_3OD)	2 (in CD_3OD)	2 (in $\text{DMSO}-d_6$)
2	79.4	81.1	78.1
3	73.7	58.4	56.6
4	69.3	79.8	77.5
5	35.4	36.6	34.8
6	73.9	72.4	70.5
7	75.8	76.2	73.9
8	39.8	40.0	38.4
1'	133.5	135.8	134.7
2'	112.7	111.7	110.7
3'	148.7	149.2	147.4
4'	147.2	147.4	145.6
5'	115.7	116.3	115.2
6'	122.0	120.8	118.7
1''	131.1	131.4	130.1
2''-6''	131.4	131.3	129.9
3''-5''	116.0	116.0	114.6
4''	156.7	156.5	155.1
OCH_3	56.5	56.5	55.6

Similar to **1**, compound **2** exhibited the negative Cotton effect at 230 nm in the ECD spectrum, indicating 2*R*-configuration. For the relative configurations of the pyran ring, the coupling constants and NOESY correlations were analyzed. Because signals of H-3 and H-6 overlapped in the ^1H NMR spectrum of **2** recorded in CD_3OD , the NMR data in $\text{DMSO}-d_6$ of **2** is provided as well (see supplemental material). The smaller coupling constant between H-2 and H-3 ($J_{2,3} = 6.0$ Hz in CD_3OD and 4.8 in $\text{DMSO}-d_6$) indicated that H-2 and H-3 were in an axial-equatorial relationship but not di-axial in case of **1**. The NOE correlations of H-2/H-4, H-2/H-6 and H-4/H-6 confirmed the same orientation of H-2, H-4 and H-6. From these evidences, compound **2** was newly elucidated to be (2*R*,2'*R*,3*R*,3'*R*,4*S*,4'*S*,6*S*,6'*S*)-6,6'-bis(1-hydroxy-2-(4-hydroxyphenyl)ethyl)-2,2'-bis(4-hydroxy-3-methoxyphenyl)octahydro-2*H*,2'*H*-[3,3'-bipyran]-4,4'-diol, for which the name zingiberpyran B is proposed.

All the isolated compounds were evaluated for their anti-inflammatory effect via inhibition of nitric oxide production. In RAW264.7 cells stimulated by lipopolysaccharide (LPS), rhamnocitrin-3-*O*-rhamnoside (**3**) exhibited the strongest effect with the IC_{50} value of 49.6 ± 0.73 μM while zingiberpyran A (**1**) proved less active (IC_{50} 71.0 μM). The other compounds showed any remark inhibition at 100 μM .

Conclusions

Two new diarylheptanoid compounds named zingiberpyrans A and B, and four known compounds, vanillin, (+)- α -viniferin, rhamnocitrin-3-*O*-rhamnoside, and meranzin hydrate were isolated for the first time from the rhizomes of *Zingiber pellitum*. Their structures were elucidated by using NMR, HR-ESI-MS and CD spectral data. All isolated compounds were evaluated for the inhibition of NO production in the LPS-stimulated RAW264.7 cells. Compound **1** and **3** showed remarkable inhibitory activity with IC_{50} value of 71.0 and 49.6 μM , respectively.

Experimental Section

General:

Plant material

Zingiber pellium was collected from Binh Chau-Phuoc Buu Nature Reserve, Ba Ria - Vung Tau, Vietnam in January 2023 and was authenticated by one of the authors, Dr. Nguyen Quoc Binh. A voucher specimen (NPH1.2023) is deposited at the herbarium of the Institute of Ecology and Biological Resources.

Chemical and apparatus

Thin layer chromatography (TLC) was conducted on pre-coated silica gel (60 F254, Merck) and detected under ultraviolet light and sprayed with aqueous sulfuric acid 10%, vanillin 10% or ceric sulfate, then heated at 105°C until the spots appear clearly. Column chromatography (CC) was performed in silica gel (230-400 mesh ASTM, Merck), and Diaion HP 20 (Merck). HPLC system was used as Agilent HPLC 1100 series, coupling with detector DAD, and a semi-preparative process was performed on YMC-pack ODS-A 250x10 mm, 5µm, 20 nm column at a flow rate of 3.0 mL/min.

NMR spectra were recorded on a Bruker AvanceNEO 600 MHz and a Bruker 500MHz spectrometer using TMS-tetramethyl silane as an internal standard. HR-MS were collected on the Agilent 6530 Accurate-Mass Q-TOF LC/MS. Optical rotations were recorded on a Jasco P-2000 Digital Polarimeter.

Extraction and isolation

1.3 kg of dried powder of *Zingiber pellium* (ZP) was extracted exhaustively with methanol. Three methanol extractions (1.5L/time) were filtered, combined and concentrated under reduced pressure to obtain a concentrated total extract. This was partitioned with *n*-hexane and methanol to give the corresponding extracts: *n*-hexane extract (15.93 g) and methanol extract (39.13 g).

The methanol extract was loaded on the Diaion HP-20 CC with stepwise elution water, methanol 40%, methanol 100% and acetone 100% in order to collect 4 fractions M0, M40, M100 and A100 respectively. M100 fraction was chromatographed on silica gel CC and eluted with gradient elution of dichloromethane/methanol (10/1 – 1/1, v/v) to collect 8 fractions (from F1.1 to F1.8). Fraction F1.3 was separated by silica gel CC with hexane/ethyl acetate (4/1, v/v) followed by preparative HPLC (30-100% MeOH in H₂O in 120 min) to afford compound **5** (3.6 mg). Fraction F1.6 was separated on a column with silica gel (stationary phase) and dichloromethane/acetone (5/1, v/v) (mobile phase), to produce 8 fractions, denoted as F2.1 to F2.8. Fraction F2.6 was continually chromatographed on HPLC (40-80% MeOH in H₂O (HPLC grade) in 180 min) to yield compounds **1** (9.3 mg) and **2** (4.1 mg). Compound **6** was separated from fraction F2.8 by preparative HPLC eluting (30-80% MeOH in H₂O (HPLC grade) in 200 min). Compounds **3** (3.0 mg) and **4** (2.7 mg) were isolated from fraction F1.7 by semi-preparative HPLC eluting with gradient solvent from 30 to 70% MeOH in 180 min.

Zingiberpyran A (**1**): amorphous pale yellow powder; $[\alpha]_D^{28} = -19.6$ (c 0.10, MeOH); ECD (MeOH) $\lambda_{\max} (\Delta\epsilon)$ 202 (–14.8), 225 (–1.22); HR-ESI-MS $m/z = 399.1379$ $[M + Na]^+$ (calcd. 399.1420 for C₂₀H₂₄O₇Na). ¹H-NMR (600 MHz, CD₃OD) and ¹³C-NMR (150 MHz, CD₃OD), see Table 1 and 2.

Zingiberpyran B (**2**): amorphous pale yellow powder; $[\alpha]_D^{28} = -17.4$ (c 0.03, MeOH); ECD (MeOH) $\lambda_{\max} (\Delta\epsilon)$ 206 (–5.6), 230 (–4.96); HR-ESI-MS $m/z = 741.2852$ $[M + Na]^+$ (calcd. 741.2887 for C₄₀H₄₆NaO₁₂). ¹H-NMR (600 MHz, CD₃OD and DMSO-*d*₆) and ¹³C-NMR (150 MHz, CD₃OD and DMSO-*d*₆), see Table 1 and 2.

Inhibition assay for nitric oxide production

The inhibitory effect of four isolated compounds was determined using a previously reported procedure¹⁵ with dexamethasone used as a positive control (IC₅₀ 13.3 µM). All experiments were performed in triplicate.

Acknowledgements

This study was financially supported by the Vietnam Academy of Science and Technology (VAST) under the grant number VAST04.05/23-24.

Supplementary Material

Full experimental detail including NMR, HRESIMS and ECD spectra of compounds **1** and **2** can be found via the “Supplementary information” section of this article’s webpage.

References

1. Leong-Skornickova, J.; Binh, N. Q.; Dang, T. H.; Sida, O.; Rybkova, R.; Vuong, T.B. *Phytotaxa* **2015**, *219*, 201-220.
<https://doi.org/10.11646/phytotaxa.219.3.1>
2. Deng, M.; Yun, X.; Ren, S.; Qing, Z.; Luo, F. *Molecules* **2022**, *27*, 2826.
<https://doi.org/10.3390/molecules27092826>
3. Hanh, N. P.; Nguyen, S. K.; Thanh, B. V.; Binh, N. Q.; Ogunwande, I. J. *J. Essent. Oil Bear. Pl.* **2023**, *26*, 937-945.
<https://doi.org/10.1080/0972060X.2023.2252835>
4. Giang, P. M.; Son, P. T.; Konig, W. A. *J. Essent. Oil Bear. Pl.* **2011**, *14*, 494-497.
<https://doi.org/10.1080/0972060X.2011.10643607>
5. Van, H. T.; Dam, S. M.; Phan U. T. X.; Nguyen, T. N. A.; Nguyen, T. B. T.; Tran, T. L.; Luu, T. N.; Le, V. S.; Huynh, N. T. A. *Acta Univ. Agric. Silv. Mendeliana Brun.* **2022**, *70*, 273-281.
<https://doi.org/10.11118/actaun.2022.020>
6. Fukunaga, T.; Nishiya, K.; Kajikawa, I.; Watanabe, Y.; Suzuki, N.; Takeya, K.; Itokawa, H. *Chem. Pharm. Bull.* **1988**, *36*, 1180-1184.
<https://doi.org/10.1248/cpb.36.1180>
7. Sarker, S. D.; Habibi, B.; Sharifi, T.; Asnaashari, S.; Nahar, L.; Delazar, A. *Orient. Pharm. Exp. Med.* **2008**, *8*, 222-227.
<https://doi.org/10.3742/OPEM.2008.8.3.222>
8. Cicchetti, E.; Silvestre, V.; Fieber, W.; Sommer, H.; Remaud, G.; Akoka, S.; Chaintreau, A. *Flavour Fragr. J.* **2010**, *25*, 463-467.
<https://doi.org/10.1002/ffj.2006>
9. Kumar, D.; Gupta, N.; Ghosh, R.; Gaonkar, R. H.; Pal, B. C. *J. Funct. Foods* **2013**, *5*, 211-218.
<https://doi.org/10.1016/j.jff.2012.10.007>
10. Huitric, A. C.; Carr, J. B.; Trager, W. F.; Nist, B. J. *Tetrahedron* **1963**, *19*, 2145-2151.
[https://doi.org/10.1016/0040-4020\(63\)85029-2](https://doi.org/10.1016/0040-4020(63)85029-2)
11. Lin, Y. S.; Lin, J. H.; Chang, C. C.; Lee S. S. *J. Nat. Prod.* **2015**, *78*, 181-187.
<https://doi.org/10.1021/np500441r>
12. Yin, H.; Luo, J. G.; Kong, L. Y. *Phytochem. Lett.* **2013**, *6*, 403-406.
<https://doi.org/10.1016/j.phytol.2013.05.004>
13. Ali, M. S.; Tezuka, Y.; Banskota, A. H.; Kadota S. *J.Nat.Prod.* **2001**, *64*, 491-496.

<https://doi.org/10.1021/np000493l>

14. Lv, H.; She, G. *Nat. Prod. Commun.* **2010**, 5, 1687-1708.

<https://doi.org/10.1177/1934578X1000501035>

15. Hai, C. T.; Luyen, N. T.; Giang, D. H.; Minh, B. Q.; Trung, N. Q.; Chinh, P. T.; Hau, D. V.; Dat, N. T. *Chem. Pharm. Bull.* **2023**, 71, 451-453.

<https://doi.org/10.1248/cpb.c22-00779>

This paper is an open access article distributed under the terms of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>)